


Item ID Number: 00101
Author Simmon, Vincent F.
Corporate Author Stanford Research Institute, Menlo Park, California
Report/Article Title Evaluation of Selected Pesticides as Chemical Mutagens 'In Vitro' and 'In Vivo' Studies
Journal/Book Title
Year 1977
Month/Day May
Color 
Number of Images 252
Description Notes Contract No. 68-01-2458; EPA-600/1-77-028

Simmon, V.F.,
1977

Evaluation of selected pesticides as chemical
mutagens 'In vito' and 'In vivo' studies

U.S. DEPARTMENT OF COMMERCE
National Technical Information Service

PB-268 647

Evaluation of Selected Pesticides as Chemical Mutagens 'In vitro' and 'In vivo' Studies

Stanford Research Institute, Menlo Park, Calif

Simmon, V.F

Prepared for

Health Effects Research Lab, Research Triangle Park, N C

May 77

AEROMEDICAL LIBRARY

APR 21 1980

DOCUMENTS

TECHNICAL REPORT
 (Please read instructions on the reverse before completing)

1. REPORT NO. EPA-600/1-77-028	2.	3. RECIPIENT'S ACCESSION NO. PB 268647
4. TITLE AND SUBTITLE EVALUATION OF SELECTED PESTICIDES AS CHEMICAL MUTAGENS In Vitro and In Vivo Studies		5. REPORT DATE May 1977
7. AUTHOR(S) Vincent F. Simpson, Ann D. Mitchell and Ted A. Jorgenson		6. PERFORMING ORGANIZATION CODE
9. PERFORMING ORGANIZATION NAME AND ADDRESS Stanford Research Institute Menlo Park, California 94025		8. PERFORMING ORGANIZATION REPORT NO. LSU-3493
12. SPONSORING AGENCY NAME AND ADDRESS Health Effects Research Laboratory - RTP, NC Office of Research and Development U.S. Environmental Protection Agency Research Triangle Park, N.C. 27711		10. PROGRAM ELEMENT NO. 1EA615
		11. CONTRACT/GRANT NO. 68-01-2458
		13. TYPE OF REPORT AND PERIOD COVERED
		14. SPONSORING AGENCY CODE 600/11

15. SUPPLEMENTARY NOTES

16. ABSTRACT

Twenty pesticides being reviewed as part of the EPA Substitute Chemical Program were studied for mutagenic activity by several in vitro and in vivo test procedures. The pesticides reviewed were: monocrotophos, bromacil, cacodylic acid, captan, chlorpyrifos, dinoseb, DSMA, fenthion, folpet, azinphos-methyl, malathion, methomyl, monuron, MSMA, parathion, parathion-methyl, quintozene (PCNB), phorate, simazine, and trifluralin.

Ten of the twenty compounds were evaluated in vivo by the mouse dominant lethal test. All twenty compounds were tested in vitro. None of the ten compounds tested in the mouse produced a dominant lethal response. Ten of the twenty compounds were mutagenic in one or more in vitro assays. Two were mutagenic in all of the in vitro assays: captan and folpet.

17. KEY WORDS AND DOCUMENT ANALYSIS		
a. DESCRIPTORS	b. IDENTIFIERS/OPEN ENDED TERMS	c. COSATI Field/Group
pesticides in vivo analysis in vitro analysis mutagens		06 F 06 T
19. DISTRIBUTION STATEMENT RELEASE TO PUBLIC	19. SECURITY CLASS (This Report) UNCLASSIFIED 20. SECURITY CLASS (This page)	21. NO. 22. PRICE A12-A01

NOTICE

THIS DOCUMENT HAS BEEN REPRODUCED FROM THE BEST COPY FURNISHED US BY THE SPONSORING AGENCY. ALTHOUGH IT IS RECOGNIZED THAT CERTAIN PORTIONS ARE ILLEGIBLE, IT IS BEING RELEASED IN THE INTEREST OF MAKING AVAILABLE AS MUCH INFORMATION AS POSSIBLE.

RESEARCH REPORTING SERIES

Research reports of the Office of Research and Development, U.S. Environmental Protection Agency, have been grouped into nine series. These nine broad categories were established to facilitate further development and application of environmental technology. Elimination of traditional grouping was consciously planned to foster technology transfer and a maximum interface in related fields. The nine series are:

1. Environmental Health Effects Research
2. Environmental Protection Technology
3. Ecological Research
4. Environmental Monitoring
5. Socioeconomic Environmental Studies
6. Scientific and Technical Assessment Reports (STAR)
7. Interagency Energy-Environment Research and Development
8. "Special" Reports
9. Miscellaneous Reports

This report has been assigned to the ENVIRONMENTAL HEALTH EFFECTS RESEARCH series. This series describes projects and studies relating to the tolerances of man for unhealthful substances or conditions. This work is generally assessed from a medical viewpoint, including physiological or psychological studies. In addition to toxicology and other medical specialties, study areas include biomedical instrumentation and health research techniques utilizing animals — but always with intended application to human health measures.

This document is available to the public through the National Technical Information Service, Springfield, Virginia 22161.

EPA-600/1-77-028
May 1977

EVALUATION OF SELECTED PESTICIDES AS CHEMICAL MUTAGENS

In Vitro and In Vivo Studies

By

Vincent F. Simmon, Ann D. Mitchell, and Ted. A. Jorgenson
Stanford Research Institute
Menlo Park, California 94025

Contract No. 68-01-2458

Project Officer

Michael D. Waters
Environmental Toxicology Division
Health Effects Research Laboratory
Research Triangle Park, N.C. 27711

U.S. ENVIRONMENTAL PROTECTION AGENCY
OFFICE OF RESEARCH AND DEVELOPMENT
HEALTH EFFECTS RESEARCH LABORATORY
RESEARCH TRIANGLE PARK, N.C. 27711

DISCLAIMER

This report has been reviewed by the Health Effects Research Laboratory, U.S. Environmental Protection Agency, and approved for publication. Approval does not signify that the contents necessarily reflect the views and policies of the U.S. Environmental Protection Agency, nor does mention of trade names or commercial products constitute endorsement or recommendation for use.

FOREWORD

The many benefits of our modern, developing, industrial society are accompanied by certain hazards. Careful assessment of the relative risk of existing and new man-made environmental hazards is necessary for the establishment of sound regulatory policy. These regulations serve to enhance the quality of our environment in order to promote the public health and welfare and the productive capacity of our Nation's population.

The Health Effects Research Laboratory, Research Triangle Park, conducts a coordinated environmental health research program in toxicology, epidemiology, and clinical studies using human volunteer subjects. These studies address problems in air pollution, non-ionizing radiation, environmental carcinogenesis and the toxicology of pesticides as well as other chemical pollutants. The Laboratory develops and revises air quality criteria documents on pollutants for which national ambient air quality standards exist or are proposed, provides the data for registration of new pesticides or proposed suspension of those already in use, conducts research on hazardous and toxic materials, and is preparing the health basis for non-ionizing radiation standards. Direct support to the regulatory function of the Agency is provided in the form of expert testimony and preparation of affidavits as well as expert advice to the Administrator to assure the adequacy of health care and surveillance of persons having suffered imminent and substantial endangerment of their health.

This report describes the testing of a series of twenty technical grade pesticide chemicals for genotoxic properties by use of a battery of in vitro and in vivo methods. The battery includes tests for gene and chromosomal mutations and primary damage to DNA as measured by effects on DNA repair recombination. Since DNA is chemically similar in all species, test results from a variety of cells and organisms are relevant in assessing the potential genetic hazard of pesticide chemicals in humans.



John H. Knelson, M.D.
Director,
Health Effects Research Laboratory

ABSTRACT

Twenty pesticides being reviewed as a part of the EPA Substitute Chemical Program were studied for mutagenic activity by several in vivo and in vitro test procedures. Ten of the twenty compounds were evaluated in vivo by the mouse dominant lethal test. All twenty compounds were tested by the following in vitro procedures:

Unscheduled DNA synthesis (UDS) in human fibroblasts (WI-38 cells); reverse mutation in Salmonella typhimurium strains TA1535, TA1537, TA1538, and TA100 and in Escherichia coli WP2; mitotic recombination in the yeast Saccharomyces cerevisiae D3; and preferential toxicity assays in DNA repair-proficient and -deficient strains of E. coli (strains W3110 and p3478, respectively) and Bacillus subtilis (strains H17 and M45, respectively).

None of the ten compounds tested in the mouse produced a dominant lethal response.

Ten of the twenty compounds were mutagenic in one or more in vitro assays. Two were mutagenic in all of the in vitro assays: captan and folpet. In a heritable translocation study in mice, under the experimental procedures employed, captan at 5000 ppm in the diet of male mice for 8 consecutive weeks produced a heritable mutagenic event in F₁ generation male mice.

CONTENTS

LIST OF TABLES	111
SUMMARY	1
TWENTY PESTICIDES EVALUATED BY SRI FOR MUTAGENIC ACTIVITY	3
INTRODUCTION	4
DOMINANT LETHAL TEST IN THE MOUSE	6
General	6
Experimental	6
Animals and Chemicals	6
Determination of Acute Toxicity	7
Maximum Tolerated Dose Study	7
Treatment Levels	7
Administration of the Compounds	8
Test Groups	8
Necropsy and Evaluation	9
Results and Discussion	9
MAMMALIAN <u>IN VITRO</u> UNSCHEDULED DNA SYNTHESIS ASSAYS	12
General	12
Experimental	13
Cell Culture	13
Dilution of Compounds	13
Controls	13
UDS Assays	14
Interpretation of Results	15
MICROBIOLOGICAL ASSAYS	17
General	17
Experimental	18
<u>Salmonella typhimurium</u> Strains TA1535, TA1537, TA1538, and TA100	18
<u>Escherichia coli</u> WP2	19
<u>Escherichia coli</u> W3100/p3478 and <u>Bacillus subtilis</u> H17/M45	20
<u>Saccharomyces cerevisiae</u> D3	21
Aroclor 1254-Stimulated Metabolic Activation System	22
Results and Discussion	22
DISCUSSION	25
REFERENCES	26
APPENDIX A: Mutagenesis Studies of Pesticide Compounds - Mouse Heritable Translocation Test - Captan	171

TABLES

DOMINANT LETHAL TEST

1	Chi-Square Test of the Fertility Index - Monocrotophos	28
2	Average Implants per Pregnant Female - Monocrotophos	29
3	Average Dead Implants per Pregnant Female - Monocrotophos. . .	30
4	Chi-Square Test of the Death Index - Monocrotophos	31
5	Number of Dead Implants per Total Implants - Monocrotophos . .	32
6	Chi-Square Test of the Fertility Index - Bromacil.	33
7	Average Implants per Pregnant Female - Bromacil.	34
8	Average Dead Implants per Pregnant Female - Bromacil	35
9	Chi-Square Test of the Death Index - Bromacil.	36
10	Number of Dead Implants per Total Implants - Bromacil.	37
11	Chi-Square Test of the Fertility Index - Captan.	38
12	Average Implants per Pregnant Female - Captan.	39
13	Average Dead Implants per Pregnant Female - Captan	40
14	Chi-Square Test of the Death Index - Captan.	41
15	Number of Dead Implants per Total Implants - Captan.	42
16	Chi-Square Test of the Fertility Index - Folpet.	43
17	Average Implants per Pregnant Female - Folpet.	44
18	Average Dead Implants per Pregnant Female - Folpet	45
19	Chi-Square Test of the Death Index - Folpet.	46
20	Number of Dead Implants per Total Implants - Folpet.	47
21	Chi-Square Test of the Fertility Index - Azinphos-Methyl	48
22	Average Implants per Pregnant Female - Azinphos-Methyl	49
23	Average Dead Implants per Pregnant Female - Azinphos-Methyl. .	50
24	Chi-Square Test of the Death Index - Azinphos-Methyl	51
25	Number of Dead Implants per Total Implants - Azinphos-Methyl .	52
26	Chi-Square Test of the Fertility Index - Malathion	53
27	Average Implants per Pregnant Female - Malathion	54
28	Average Dead Implants per Pregnant Female - Malathion.	55
29	Chi-Square Test of the Death Index - Malathion	56
30	Number of Dead Implants per Total Implants - Malathion	57
31	Chi-Square Test of the Fertility Index - Parathion	58
32	Average Implants per Pregnant Female - Parathion	59
33	Average Dead Implants per Pregnant Female.	60
34	Chi-Square Test of the Death Index - Parathion	61
35	Number of Dead Implants per Total Implants - Parathion	62
36	Chi-Square Test of the Fertility Index - Parathion-Methyl. . . .	63
37	Average Implants per Pregnant Female - Parathion-Methyl. . . .	64
38	Average Dead Implants per Pregnant Female - Parathion-Methyl .	65
39	Chi-Square Test of the Death Index - Parathion-Methyl.	66
40	Number of Dead Implants per Total Implants - Parathion-Methyl.	67
41	Chi-Square Test of the Fertility Index - Quintozene (PCNB) . . .	68
42	Average Implants per Pregnant Female - Quintozene (PCNB) . . .	69

43	Average Dead Implants per Pregnant Female - Quintozene (PCNB)	70
44	Chi-Square Test of the Death Index - Quintozene (PCNB).	71
45	Number of Dead Implants per Total Implants - Quintozene (PCNB)	72
46	Chi-Square Test of the Fertility Index - Phorate.	73
47	Average Implants per Pregnant Female - Phorate.	74
48	Average Dead Implants per Pregnant Female - Phorate	75
49	Chi-Square Test of the Death Index - Phorate.	76
50	Number of Dead Implants per Total Implants - Phorate.	77

UDS ASSAYS

51	DNA Repair Synthesis Assay of Monocrotophos	78
52	DNA Repair Synthesis Assay of Monocrotophos with Metabolic Activation.	79
53	DNA Repair Synthesis Assay of Bromacil.	80
54	DNA Repair Synthesis Assay of Bromacil with Metabolic Activation.	81
55	DNA Repair Synthesis Assay of Cacodylic Acid.	82
56	DNA Repair Synthesis Assay of Cacodylic Acid with Metabolic Activation.	83
57	DNA Repair Synthesis Assay of Captan.	84
58	DNA Repair Synthesis Assay of Captan with Metabolic Activation.	85
59	DNA Repair Synthesis Assay of Chloropyrifos	86
60	DNA Repair Synthesis Assay of Chloropyrifos with Metabolic Activation.	87
61	DNA Repair Synthesis Assay of Dinoseb	88
62	DNA Repair Synthesis Assay of Dinoseb with Metabolic Activation.	89
63	DNA Repair Synthesis Assay of DSMA.	90
64	DNA Repair Synthesis Assay of DSMA with Metabolic Activation.	91
65	DNA Repair Synthesis Assay of Fenthion.	92
66	DNA Repair Synthesis Assay of Fenthion with Metabolic Activation.	93
67	DNA Repair Synthesis Assay of Folpet.	94
68	DNA Repair Synthesis Assay of Folpet with Metabolic Activation.	95
69	DNA Repair Synthesis Assay of Azinphos-Methyl	96
70	DNA Repair Synthesis Assay of Azinphos-Methyl with Metabolic Activation.	97
71	DNA Repair Synthesis Assay of Malathion	98
72	DNA Repair Synthesis Assay of Malathion with Metabolic Activation.	99
73	DNA Repair Synthesis Assay of Methomyl.	100
74	DNA Repair Synthesis Assay of Methomyl with Metabolic Activation.	101
75	DNA Repair Synthesis Assay of Monuron	102

76	DNA Repair Synthesis Assay of Monuron with Metabolic Activation.	103
77	DNA Repair Synthesis Assay of MSMA.	104
78	DNA Repair Synthesis Assay of MSMA with Metabolic Activation.	105
79	DNA Repair Synthesis Assay of Parathion	106
80	DNA Repair Synthesis Assay of Parathion with Metabolic Activation.	107
81	DNA Repair Synthesis Assay of Parathion-Methyl.	108
82	DNA Repair Synthesis Assay of Parathion-Methyl with Metabolic Activation	109
83	DNA Repair Synthesis Assay of Quintozene (PCNB)	110
84	DNA Repair Synthesis Assay of Quintozene (PCNB) with Metabolic Activation	111
85	DNA Repair Synthesis Assay of Phorate	112
86	DNA Repair Synthesis Assay of Phorate with Metabolic Activation.	113
87	DNA Repair Synthesis Assay of Simazine.	114
88	DNA Repair Synthesis Assay of Simazine with Metabolic Activation.	115
89	DNA Repair Synthesis Assay of Trifluralin	116
90	DNA Repair Synthesis Assay of Trifluralin with Metabolic Activation.	117

MICROBIOLOGICAL ASSAYS

91	<u>In Vitro Assays with <i>Salmonella typhimurium</i></u>	118
92	<u>Results of Assays with <i>Escherichia coli</i> WP2.</u>	138
93	<u>Microbial Inhibition in <i>Escherichia coli</i> and <i>Bacillus subtilis</i></u>	148
94	<u>In Vitro Assays with <i>Saccharomyces cerevisiae</i> D3 - Monocrotophos</u>	150
95	<u>In Vitro Assays with <i>Saccharomyces cerevisiae</i> D3 - Bromacil.</u>	151
96	<u>In Vitro Assays with <i>Saccharomyces cerevisiae</i> D3 - Cacodylic Acid.</u>	152
97	<u>In Vitro Assays with <i>Saccharomyces cerevisiae</i> D3 - Captan.</u>	153
98	<u>In Vitro Assays with <i>Saccharomyces cerevisiae</i> D3 - Chloropyrifos</u>	154
99	<u>In Vitro Assays with <i>Saccharomyces cerevisiae</i> D3 - Dinoseb</u>	155
100	<u>In Vitro Assays with <i>Saccharomyces cerevisiae</i> D3 - DMSA</u>	156
101	<u>In Vitro Assays with <i>Saccharomyces cerevisiae</i> D3 - Fenthion.</u>	157
102	<u>In Vitro Assays with <i>Saccharomyces cerevisiae</i> D3 - Folpet.</u>	158
103	<u>In Vitro Assays with <i>Saccharomyces cerevisiae</i> D3 - Azinphos-Methyl</u>	159
104	<u>In Vitro Assays with <i>Saccharomyces cerevisiae</i> D3 - Malathion</u>	160

105	<u>In Vitro Assays with <i>Saccharomyces cerevisiae</i> D3 -</u> Methomyl	161
106	<u>In Vitro Assays with <i>Saccharomyces cerevisiae</i> D3 -</u> Mouuron	162
107	<u>In Vitro Assays with <i>Saccharomyces cerevisiae</i> D3 -</u> MSMA	163
108	<u>In Vitro Assays with <i>Saccharomyces cerevisiae</i> D3 -</u> Parathion	164
109	<u>In Vitro Assays with <i>Saccharomyces cerevisiae</i> D3 -</u> Parathion-Methyl	165
110	<u>In Vitro Assays with <i>Saccharomyces cerevisiae</i> D3 -</u> Quintozene (PCNB)	166
111	<u>In Vitro Assays with <i>Saccharomyces cerevisiae</i> D3 -</u> Phorate	167
112	<u>In Vitro Assays with <i>Saccharomyces cerevisiae</i> D3 -</u> Simazine	168
113	<u>In Vitro Assays with <i>Saccharomyces cerevisiae</i> D3 -</u> Trifluralin	169
114	<u>In Vitro Mutagenesis with <i>Salmonella typhimurium</i>.</u>	170

ACKNOWLEDGMENTS

Stanford Research Institute wishes to thank the EPA project officers for their guidance and assistance during the course of this project. Dr. Robert E. McGaughy, Office of Research and Development, Washington, D.C., was Project Officer from the beginning of the project in June 1974 until July 1975. Project responsibility then was transferred to Dr. Ronald L. Baron, Health Effects Research Laboratory, Research Triangle Park, North Carolina, until January 1976. At that time, Dr. Michael D. Waters, now Chief of the Biochemistry Branch in the Environmental Toxicology Division, Health Effects Research Laboratory, Research Triangle Park, North Carolina, became Project Officer. Dr. Waters has continued in this project responsibility to the present.

INTRODUCTION

The Federal Insecticide, Fungicide, and Rodenticide Act designates the Environmental Protection Agency as the governmental body responsible for the safety of all pesticides used in the United States. More recently, the Federal Environmental Pesticide Control Act (PL 92-516) strengthened EPA's regulatory responsibilities in the area of pesticides to include intra- as well as inter-state commerce.

To be federally registered, a pesticide must have been determined not to be hazardous to health or to the environment when used according to its labeling restrictions. Thus, relative to new law as well as to specific directives included in Public Law 93-135, 1973, EPA now is conducting a thorough review of the implications of using alternate chemicals, including older registered pesticides, for pest control.

In the pesticide review process, EPA emphasizes development of scientific criteria for evaluating the safety of compounds substituted for those pesticides found to be hazardous. In addition to reviewing and evaluating the literature on pesticides and maintaining liaison with industry and academia, the strategy program includes laboratory studies to obtain additional data. One of these laboratory programs is directed toward gathering mutagenesis data on a selected number of compounds.

EPA's program is timely and responsive to one of the recommendations included in the President's Scientific Advisory Committee Report of September 1973, Chemicals and Health. In that document, the Committee recommended that "Regulatory agencies should take steps to insure that new scientific data raising the possibility of new or extended hazards from chemicals in use are subject to careful process of scientific review for merit interpretation."

Development of methods for evaluating the mutagenic hazard of chemical compounds has advanced markedly in the last few years. In contrast to the undefined empirical tests used a short time ago, procedures now

available can detect chromosome breaks and other genetic changes caused by chemical stress. Mutant strains of microorganisms in cell culture and mammalian fibroblast cells in tissue culture are effective in vitro systems for reliable detection of presumptive gene mutations, whereas the mammalian dominant lethal test is a recognized test for the assessment of chromosome damage to germinal cells.

Today many pesticide chemicals in commercial use have not been investigated adequately for their mutagenic hazard. With the public's increasing concern about possible pollution of our environment by chemicals, the widely used pesticides must be evaluated. In this project, SRI used test methods that are appropriate for these evaluations and that are in use by the scientific community.

Under contract to EPA, SRI examined 20 pesticides for mutagenic activity using a combination of in vivo and in vitro mutagenicity assay systems. The 20 pesticides tested and their sources are listed in the following two tables.

The assays used were the dominant lethal test in mice (only ten compounds); unscheduled DNA synthesis (UDS) in human fibroblasts (WI-38 cells); reverse mutation in Salmonella typhimurium strains TA1535, TA1537, TA1538, and TA100 and in Escherichia coli WP2; mitotic recombination in the yeast Saccharomyces cerevisiae D3; and preferential toxicity assays in DNA repair-proficient and -deficient strains of E. coli (strains W3110 and p3478, respectively) and Bacillus subtilis (strains H17 and M45, respectively).

Based on positive responses in both Tier I (in vitro test) and Tier II (Drosophila) mutagenic studies, it was recommended that a heritable translocation test (Tier III) in the mouse be conducted

to further assess the mutagenic potential of Captan. The results of these further studies are reported as Appendix A.

The experimental procedures and results for the mammalian dominant lethal test, the UDS assay, and the microbiological assays are described in the separate sections that follow.

IN VIVO AND IN VITRO MUTAGENESIS: SUMMARY DATA FOR EPA PESTICIDES

Positive Response, +; Negative Response, -

Pesticide	Mouse Dominant Lethal*	<u>Salmonella</u> <u>typhimurium</u> †		<u>Escherichia coli</u> WP2		<u>Saccharomyces cerevisiae</u>		<u>Escherichia coli</u>	<u>Sacillus aurtilis</u>	UDS	
		(His ⁺ Reversion)		(Try ⁺ Reversion)		(Mitotic Recombination)				(Relative Toxicity)	(Relative Toxicity)
		-MA	+MA	-MA	+MA	-MA	+MA			-MA	+MA
Monocrotophos	-	-	-	-	-	+	+	-	-	+	+
Bromacil	-	-	-	-	-	-	-	-	-	-	-
Sacodylic Acid	-	-	-	-	-	+	+	-	-	-	-
Captan	-	+	+	+	+	+	+	+	+	-	+
Chlorpyrifos	-	-	-	-	-	-	-	+	+	-	-
Dinoseb	-	-	-	-	-	-	-	+	+	-	-
DSMA	-	-	-	-	-	-	-	-	-	-	-
Fenthion	-	-	-	-	-	-	-	-	-	-	-
Folpet	-	+	+	+	+	+	+	+	+	-	+
† Azinphos-methyl	-	-	-	-	-	+	+	+	-	-	+
Malathion	-	-	-	-	-	-	-	-	-	-	-
Mathomyl	-	-	-	-	-	-	-	-	-	-	-
Monuron	-	-	-	-	-	-	-	-	+	-	+
MSMA	-	-	-	-	-	-	-	-	-	-	-
Parathion	-	-	-	-	-	-	-	-	-	+	-
Parathion-methyl	-	-	-	-	-	+	+	-	-	-	-
Quintozene (PCNB)	-	-	-	-	-	-	-	-	-	-	-
Phorate	-	-	-	-	-	-	-	-	-	-	-
Simazine	-	-	-	-	-	-	-	-	-	-	-
Trifluralin	-	-	-	-	-	-	-	-	-	-	-

* Only ten pesticides were tested by the dominant lethal procedure.

† See page 170.

‡ Marginally positive.

TWENTY PESTICIDES EVALUATED BY SRI FOR MUTAGENIC ACTIVITY

<u>Common Name*</u>	<u>Trade Name of Compound Tested</u>	<u>Manufacturer</u>	<u>Batch or Lot Number</u>	<u>Purity (%)</u>	<u>Supplier</u>
Monocrotophos	Azodrin-5	Shell Chemical Company	Batch H, 9-SCL-77	55.0	Manufacturer
Bromacil	Hyvar	E.I. DuPont de Nemours	T80619/40	95.9	Battelle
Cacodylic Acid	Phytar	Ansul Chemical Company	Phyton 138	65.6	Battelle
Captan	Orthoside 406	Chevron Chemical Company	5X640	Technical	Battelle
Chlorpyrifos	Dursban	Dow Chemical Company	MM-1114-1 (603-D1)	98.8	Battelle
Dinoseb	Premerge	Dow Chemical Company	MM 200554	97.7	Battelle
DSMA	Ansar	Ansul Chemical Company	8100	80.1	Battelle
Fenthion	Baytex	Chemogro	4-15-2026	96.0	Battelle
Folpet	Phaltan	Chevron Chemical Company	SX579	Technical	Battelle
Azinphos-methyl	Guthion	Chemogro	411-0229	Technical	Battelle
Malathion	Malathion	American Cyanamid Company	40216006.300	Technical	Battelle
Methomyl	Lannate	E.I. DuPont de Nemours	6602-82	99.0	Battelle
Monuron	Telvar	E.I. DuPont de Nemours	T-40817-20	97.0	Battelle
MSMA	Ansar	Ansul Chemical Company	170 H.C.	58.4	Battelle
Parathion	Niran	Monsanto Chemical Company	AD 1236	99.0	Battelle
Parathion-methyl	Methyl Parathion	Monsanto Chemical Company	AD 0659	80.0	Battelle
Quintozene (PCNB)	Terrachlor	Olin Mathieson Chemical Corporation	Technical	99.0	Battelle
Phorate	Thimet	American Cyanamid Company	MC85		Battelle
Simazine	Primatol	Ciba-Geigy Chemical Co.	FL-740846	97.7	Battelle
Trifluralin	Treflan	Eli Lilly & Company	X-26290	97.7	Battelle

* Common name as approved by the International Organization for Standardization.

DOMINANT LETHAL TEST IN THE MOUSE

General

In the dominant lethal test, the ten compounds under investigation were fed in the diet to proven male breeder mice for 7 weeks. After this period, each male was mated with two adult virgin females for 7 days; these females were then replaced by two others for another breeding. The sequence was continued for 8 weeks. This procedure emphasizes possible mutagenic effects on the male sperm, the normal female acting as a carrier to reveal in her offspring abnormalities that may have occurred in the male. We evaluated effects by examining the condition and state of fetal development during the middle to latter stages of gestation.

Experimental

Animals and Chemicals

Adult ICR/SIM mice from a closed, random-bred colony were used for the acute toxicity and maximum tolerated dose determinations as well as for the dominant lethal assay. These male and female mice were supplied by Simonsen Laboratories, Gilroy, California. The males were 3- to 4-month-old proven breeders, and the females were 10- to 12-week-old virgin stock.

At the direction of EPA, the Battelle Columbus Laboratories obtained the pesticides from the manufacturers and subsequently provided SRI with aliquots for the studies reported here. Each pesticide was a "technical" grade product (or equivalent) and was provided in sufficient quantity for us to complete all aspects of the experimental program. Excess supplies were refrigerated or frozen, should they be needed for future reference.

We investigated the solubility of each compound using water, propylene glycol, polyethylene glycol, corn oil, or carboxymethyl-cellulose to determine the most appropriate vehicle for administration. Compounds were administered orally, by gavage for the acute toxicity (LD₅₀) determinations, and via the diet for the maximum tolerated dose and dominant lethal studies.

Determination of Acute Toxicity

Although acute toxicity information on some of the compounds was available in the literature, we conducted confirmatory tests on all to obtain an LD₅₀ under our laboratory conditions and for the ICR/SIM strain of mouse. If no data were available, we conducted a preliminary range-finding test, followed by a determination of the oral LD₅₀.

Maximum Tolerated Dose Study

Based on the acute toxicity data and available information from the literature on dose levels known to cause adverse responses when administered in the diet, several dose levels were selected and administered in the diet to adult male mice for 2 weeks. Treated males then were caged with two adult virgin females each for 7 days; these females were replaced by two others weekly for 2 weeks. The females were examined daily for the presence of vaginal (mating) plugs. At midterm of pregnancy, the females were sacrificed and examined for total implants, as well as for early and late fetal deaths. For this work, we defined a maximum tolerated dose as that dietary level which may produce up to a 20% weight loss, mild but transient clinical signs, no inhibition of breeding performance, and no mortality. Thus, these initial studies provided information on changes in body weight, acceptability of the diet, clinical signs, mortality, and breeding performance.

Treatment Levels

For the dominant-lethal study, three dose levels were administered. The highest was the maximum tolerated dose or 5 g/kg (a maximum level

agreed on by EPA and SRI), whichever was lower. The intermediate and lower dosages were one-half and one-quarter of the highest dose, respectively.

Administration of the Compounds

Each pesticide was fed in the diet to adult male mice for 7 weeks. An appropriate amount of compound initially was dissolved or suspended in corn oil; then the compound-oil concentrate was added at a level of 3% to a finely ground commercial diet of known composition. The use of corn oil assured even distribution of the compound and prevented stratification of the test material in an otherwise dry diet. Diets were prepared at 2-week intervals and were refrigerated at 4°C until fed to the animals. Fresh diet was placed in the feed containers every other day to minimize the loss of compound through instability or volatility.

Test Groups

Two reference control groups were included in this project. One was run at the beginning of each of the two dominant lethal series, five pesticides being run concurrently. In this manner, reference breeding and implant data were obtained at two time periods, as was information on each shipment of research animals. Males in these groups were fed a finely ground commercial diet supplemented with corn oil at 3%. Control groups were treated in the same manner as the compound test groups.

Two positive control groups were run concurrently with each of the two series of five pesticide tests. For these groups, the known mutagen triethylenemelamine (TEM) was administered as a single intraperitoneal injection of 0.2 mg/kg approximately 2 hours before the first mating. A commercial pelleted diet was available at all times.

Each control and experimental test group contained 20 adult male mice. At the end of the 7-week compound treatment period, each male was allowed to breed with two virgin females over a period of 7 days. Females were replaced weekly for 8 weeks.

Necropsy and Evaluation

Females were sacrificed at midterm of pregnancy. A complete necropsy was performed to determine if an intercurrent infection was present; such a condition can induce preimplantation loss and early fetal deaths. At sacrifice, each female was scored for early fetal deaths, late fetal deaths, and living fetuses (all of which provide a total implant score).

The following parameters indicate effects in dominant lethal studies: Total implants (live fetuses plus early and late fetal deaths), total dead (early and late fetal deaths), and dead implants per total implants. Total implants and dead implants were analyzed for significance by the t-test.

The index of dead implants per total implants was analyzed statistically by the t-test on arcsine- (or angular) transformed data, as described in Experimental Design (Theory and Application).¹ This index was computed for each female. Other parameters analyzed were the fertility and death indices.

Results and Discussion

Single-dose oral acute toxicity data are as follows:

<u>Compound</u>	<u>LD50</u>
Monocrotophos	17 mg/kg
Bromacil	3.04 g/kg
Captan	> 15 g/kg
Folpet	> 10 g/kg
Azinphos-methyl	15 mg/kg
Malathion	1196 mg/kg
Parathion	17 mg/kg
Parathion-methyl	39 mg/kg
Quintozone (PCNB)	> 10 g/kg
Phorate	6.59 mg/kg

After evaluating the acute toxicity data and those from subsequent maximum tolerated dose studies, we selected the following dosage levels for the dominant lethal studies:

<u>Compound</u>	<u>Treatment Levels (mg/kg of Diet)</u>
Monocrotophos	15, 30, 60
Bromacil	1250, 2500, 5000
Captan	1250, 2500, 5000
Folpet	1250, 2500, 5000
Azinphos-methyl	20, 40, 80
Malathion	1250, 2500, 5000
Parathion	62.5, 125, 250
Parathion-methyl	20, 40, 80
Quintozene (PCNB)	1250, 2500, 5000
Phorate	5, 10, 20

Throughout the experiment, the biological criteria used to evaluate mutagenic effects in the mouse showed no consistent responses that could be attributed to treatment. Although we found occasional statistical differences between control and compound treated groups, they were random and did not suggest a time or dose-response effect.

Summary data on the fertility index, implantations per pregnant female, dead implants per pregnant female, death index, and number of dead implants per total implants are presented by compound as follows: Tables 1 through 5, Monocrotophos; Tables 6 through 10, Bromacil; Tables 11 through 15, Captan; Tables 16 through 20, Folpet; Tables 21 through 25, Azinphos-methyl; Tables 26 through 30, Malathion; Tables 31 through 35, Parathion; Tables 36 through 40, Parathion-Methyl; Tables 41 through 45, Quintozene (PCNB); and Tables 46 through 50, Phorate.

Two copies of a description of the statistical analysis procedures used for dominant lethal tests and computer printouts of the raw data and the statistical analyses are on file with the current Project Officer, Dr. Michael D. Waters, Environmental Toxicology Division, Health Effects Research Laboratory, EPA Environmental Research Center, Research Triangle Park, North Carolina 27711.

The following statistical procedures were used:

Chi-square test of the fertility index;

Armitage test for a linear trend in proportion for the fertility index based on dose levels, based on logarithms of the dose levels, and based on dose levels including the control group;

t-test of the number of implantations in pregnant females;

Regression fits of implantations on dose and log dose and with and without control group included;

t-test of the (Freeman-Tukey transformed) preimplantation losses in pregnant females;

t-test of the number of dead implants;

Chi-square test of the death index;

Armitage test for a linear trend in proportion for the death index, based on dose levels with and without control group included and based on logarithms of the dose levels;

Probit analysis of the proportion of pregnant females with one or more dead implants;

t-test of the (Freeman-Tukey transformed) number of dead implants (dead implants/total implants);

Control group analyses of variances for number of pregnant females, number of implantations per pregnant female, preimplantation loss per pregnant female, number of dead implants per pregnant female, ratio of dead implants to total implants per pregnant female; and

t-test of the number of corpora lutea in pregnant females.

Careful review and statistical evaluation of the data show that folpet, captan, parathion-methyl, parathion, phorate, malathion, bromacil, monocrotophos, quintozone (PCNB), and azinphos-methyl are not mutagenic in the mouse by the dominant lethal test.

MAMMALIAN IN VITRO UNSCHEDULED DNA SYNTHESIS ASSAYS

General

Many mutagenic and carcinogenic agents have been shown to induce unscheduled DNA synthesis (UDS) in an in vitro tissue culture system of mammalian cells. UDS is a form of mammalian repair synthesis that involves at least two processes. The first is interaction of the agent with DNA, resulting in damage of the DNA. The second, which follows, is incorporation of nucleotides to repair the DNA.

UDS may be considered a fairly universal system because it occurs in a wide variety of mammalian cell types and because it has been observed in all stages of the cell cycle (G_0 , G_1 , G_2 , and M) other than S, the normal DNA synthetic phase.^{2,3} (UDS is not observed during S-phase because the high level of incorporation of nucleotides during the scheduled DNA synthesis obscures the relatively low level of incorporation of nucleotides during unscheduled DNA synthesis.)

An additional feature of UDS is that it may detect a level of DNA damage higher than that revealed by examination of chromosomeal aberrations⁴ because some DNA repair results in little or no detectable change in chromosome morphology. For each compound tested, an in vitro metabolic activation system should be incorporated for a parallel series of UDS assays since some compounds may be ineffective in producing DNA damage unless they are first activated by a microsomal preparation from a mammalian liver homogenate.

The UDS system we have developed is unique in that, at the end of each assay, DNA is extracted from human diploid fibroblasts (WI-38 cells) so that the extent of repair may be expressed per unit of DNA. We have found that this UDS assay system affords sensitivity and precision without sacrificing efficiency or economy. Under separate contact, NCI approved our use and validation of this system for the prescreening of chemical carcinogens. With the approval of the EPA project officer, we used this system for testing the 20 substitute pesticides, with and without metabolic activation.

Experimental

Cell Culture

WI-38 cells grown in T-25 tissue culture flasks were used for the UDS assays. Replicate cultures of these cells were initiated in Eagle's Basal Medium (8ME) containing 10% (v/v) fetal calf serum and aureomycin, an antibiotic specific for PPLO*. For 1 to 2 weeks preceding the UDS assays, the cells were grown in medium containing 0.5% serum. This produced contact-inhibited cells in synchronous cultures in the G₁ phase of the mitotic cycle. To reduce further the possibility of incorporation of ³H-TdR by an occasional S-phase cell that might escape the contact-inhibition synchrony and thus obscure measurements of UDS, the cultures were preincubated for 1 hour with 10⁻² M hydroxyurea (HU) before each assay, and 10⁻² M HU was added during each subsequent step of the assays.

Dilution of Compounds

Chemicals to be tested were made up immediately before use and were diluted in appropriate solvents (water, ethanol, or DMSO), the final concentration of solvent being one that did not produce a cytotoxic effect after repeated testing. Sonification and pH adjustments were used to ensure maximum solubility or even suspension of the stock solutions of the compounds. The highest concentration was diluted further in solvent and then in culture medium to give several log dilutions of each compound. All compounds were in apparent solution and within the physiological pH range when tested, except as otherwise noted in the tables.

Controls

The positive controls were 4-nitroquinoline-N-oxide (4NQO), a compound that induces UDS in the absence of a metabolic activation system, and dimethylnitrosamine (DMN), a compound that induces UDS only with metabolic activation. The negative controls were the solvents diluted in culture medium.

* As an additional check against the presence of PPLO, which could incorporate tritiated thymidine (³H-TdR) and thus obscure measurements of UDS, stock cultures were analyzed monthly for the presence of PPLO. The results of these analyses were consistently negative.

UDS Assays

The contact-inhibited WI-38 cells were incubated at 37°C with log dilutions of the substitute pesticides and with 1 μ Ci/ml of ^3H -TdR (sp act, 6.7 Ci/mmole). For testing in the absence of metabolic activation, the cells were exposed simultaneously to the substitute pesticide and to ^3H -TdR for 3 hours. For testing with metabolic activation, the cells were exposed to the substitute pesticide, to ^3H -TdR, and to 500 mg/ml of the 9000 x g supernatant fraction of a liver homogenate from adult male Swiss-Webster mice, with appropriate cofactors,* for 1 hour; then the cells were incubated with only ^3H -TdR for an additional 4 hours. The shorter exposure time for metabolic activation testing was used to preclude cytotoxic effects of the liver homogenate preparation. Both approaches included a postincorporation incubation with unlabeled thymidine. DNA was extracted from the cells by a modification of the PCA-hydrolysis procedure;⁵ one aliquot of the DNA solution was used to measure the DNA content, after reaction with diphenylamine,⁶ and a second aliquot was used for scintillation counting measurements of the extent of incorporation of ^3H -TdR. Results were expressed as incorporated per unit of DNA and were compared with the background rate of incorporation.

We have defined as an acceptable assay one in which the response of the positive control compound is predicted, within the 95% confidence limits, by regressions of average dpm/ μ g DNA versus average dpm/ μ g for background. The regressions that follow are based on data that we have acquired in previous testing:

<u>Type of Testing</u>	<u>Regression[†]</u>	<u>Sample Size (n)</u>	<u>Correlation Coefficient (r)</u>
Without metabolic activation	$Y_1 = 696 + 17.45 (X)^\ddagger$	48	0.7668
With Metabolic activation	$Y_2 = 263 + 1.83 (X)^\ddagger$	13	0.9639

*Nicotinamide, 3.05 mg/ml; glucose-6-phosphate, 16.1 mg/ml; $\text{MgCl}_2 \cdot 6\text{H}_2\text{O}$, 5.08 mg/ml; NADP, 0.765 mg/ml.

[†]Regressions over a range of background dpm/ μ g DNA of 0 to 450.

[‡] Y_1 = Average dpm/ μ g DNA for 10^{-5} M 4NQO (positive control).

Y_2 = Average dpm/ μ g DNA for 5×10^{-2} M DMN (positive control).

X = Average dpm/ μ g DNA for background (negative control).

If the observed average level of incorporation for the positive control compound is outside the 95% confidence limits of the regression, we assume that some variation has occurred in the experimental procedure and repeat the test.

Interpretation of Results

In a report to the National Cancer Institute,⁷ we presented the results of tests performed without metabolic activation on 40 compounds of known carcinogenicity. We have analyzed these results using either the parametric One-Way Classification Analysis of Variance or the non-parametric Kruskal-Wallis One-Way Analysis of Variance, depending on which was more appropriate.* At the 99% confidence limits, all the ultimate carcinogens significantly elevate the incorporation of ³H-TdR into the DNA. The noncarcinogenic compounds, with one exception, fail to elevate significantly the incorporation of ³H-TdR at this level of confidence. Thus, the 99% confidence limits of these statistical analyses apparently can be used with reasonable accuracy to predict the biological significance of the response to a chemical.

The number of compounds we have tested with metabolic activation is insufficient to establish a correlation between statistical significance and biological significance. Therefore, we assumed that the 99% confidence levels of the analyses of variance used without metabolic activation also apply for testing with metabolic activation.

Results and Discussion

Tables 51 through 90 present the results of the UDS testing, with and without metabolic activation, of the 20 substitute pesticides. Tables 51 and 52, the DNA repair synthesis assays of monocrotophos, include detailed summaries of the cell culture and experimental conditions for these assays. The assays presented in the following tables (53 through 90) were conducted under similar conditions. In routine testing in the

*If there is reason to believe that the variances of each of the treatments in a test are equal (i.e., Bartlett's test of the variance is negative), the parametric analysis is the appropriate one. If the variances are not equal, the nonparametric analysis is the appropriate one.

absence of metabolic activation, six samples each are used for five log concentrations of each test compound and for the negative and positive controls. However, because of the expense of the metabolic activation preparations, for all compounds except bromacil we tested three replicate samples in the presence of metabolic activation and used three concentrations of the test compound (selected on the basis of the testing without metabolic activation).

Based on the criteria for positive responses, we observed significant increases in unscheduled DNA synthesis in the absence of metabolic activation after exposure of the cells to only two substitute pesticides, monocrotophos and parathion. In the presence of metabolic activation enzymes, significantly increased UDS was detected for five substitute pesticides: monocrotophos, captan, folpet, azinphos-methyl, and monuron.

Compared with those of negative controls, the levels of ^3H -TdR incorporation were greatly reduced in the absence of metabolic activation at the highest concentrations tested for captan, folpet, azinphos-methyl, and monuron, the same four compounds that induced UDS only in the presence of metabolic activation. The reduced levels of incorporation may be interpreted as cytotoxic effects or as inhibition of repair caused by the highest concentration of the test compounds. A similar effect was observed in the presence of metabolic activation for only one compound, captan, and this was observed at a higher concentration than had been tested without metabolic activation. Stich et al.⁶ have discussed the problem of cytotoxicity and possible inhibition of DNA repair systems by some chemicals and have stressed that, whereas such factors may obscure measurements of UDS, often a close relationship exists between concentrations that induce UDS and concentrations that are cytotoxic or that inhibit repair.

Because of the cytotoxic or inhibitory effects of the substitute pesticides, it should not be assumed without further testing that monocrotophos and parathion would be carcinogenic without metabolic activation or that the other four substitute pesticides that induced UDS in the presence of metabolic activation are procarcinogens. The positive UDS results indicate that these six substitute pesticides should be tested more extensively, with the testing to include evaluations of the effects of these chemicals in in vivo bioassays.

MICROBIOLOGICAL ASSAYS

General

SRI examined twenty pesticides for mutagenicity by in vitro microbiological assays with Salmonella typhimurium (TA1535, TA1537, TA1538, TA100), Escherichia coli WP2, repair-deficient and -proficient strains of Bacillus subtilis and E. coli, and with the yeast Saccharomyces cerevisiae D3. An Aroclor 1254-stimulated, rat-liver-homogenate metabolic activation system was included in each procedure, except the relative toxicity assays, to provide metabolic steps that the bacteria are either incapable of conducting or that they do not carry out under the assay conditions. The purpose of this study was to determine whether the compounds elicited a mutagenic response in microorganisms.

The assay procedure with S. typhimurium has been proven to be 85 to 90% accurate in detecting carcinogens as mutagens, and it has about the same accuracy in identifying chemicals that are not carcinogenic.⁹ The assay procedure with S. cerevisiae is about 50% accurate in detecting carcinogens as agents that increase mitotic recombination. E. coli WP2 and the microbial sensitivity assay are two additional methods of detecting mutagens. The combination of these four assay procedures significantly enhances the probability of detecting potentially hazardous chemicals.

Experimental

Salmonella typhimurium Strains TA1535, TA1537, TA1538, and TA100

The *S. typhimurium* strains used at SRI were obtained from Dr. Bruce Ames of the University of California at Berkeley.¹⁰⁻¹² All are histidine auxotrophs (*his*⁻) by virtue of mutations in the histidine operon. In addition to the mutations in the histidine operon, the indicator strains have mutations in the lipopolysaccharide coat (*rfa*⁻) and deletions that cover a gene involved in the repair of uv damage (*uvrB*⁻). The *rfa*⁻ mutation makes the strains more permeable to large molecules, thereby increasing their sensitivity to these molecules. The *uvrB*⁻ mutation decreases repair of some types of chemically damaged DNA and thereby enhances sensitivity to some mutagenic chemicals. Strain TA1535 is reverted to histidine prototrophy (*his*⁺) by many mutagens that cause base-pair substitutions. Strains TA1537 and TA1538 are reverted by many frameshift mutagens. TA1537 is more sensitive than TA1538 to mutation by some acridine and benzanthracenes, but the difference is quantitative rather than qualitative. TA100 is derived from TA1535 by the introduction of the R factor plasmid pKM101.¹³ The introduction of this plasmid, which confers ampicillin resistance to the strain, greatly enhances the sensitivity of the strain to some base-pair substitution mutagens. We have shown that mutagens such as benzyl chloride and 2-(2-furyl)-3-(5-nitro-2-furyl) acrylamide (known as AF2) can be detected in plate assays by TA100 but not by TA1535. The presence of this plasmid also makes strain TA100 sensitive to some frameshift mutagens--e.g., ICR-191, benzo(a)pyrene, aflatoxin B₁, and 7,12-dimethylbenz(a) anthracene.

All the indicator strains are stored at -80°C. For each experiment, an inoculum from frozen stock cultures is grown overnight at 37°C in a nutrient broth consisting of 1% tryptone and 0.5% yeast extract. After stationary overnight growth, the cultures are shaken for 3 to 4 hours to ensure optimal growth. Each culture is checked for sensitivity to crystal violet. The presence of the *rfa*⁻ mutation makes the indicator strains sensitive to this dye, whereas the parent strain, *rfa*⁺, is not sensitive to the dye. However, the mutation is reversible, leading to

the accumulation of rfa⁺ cells in the culture. Therefore, the cells must be tested routinely to ensure their sensitivity to crystal violet. Each culture also is tested by specific mutagens known to revert each test strain (positive controls).

To a sterile 13 x 100 mm test tube placed in a 43°C heating block, we add in the following order:

Assays in agar

- (1) 2 ml of 0.6% agar*
- (2) 0.1 ml of indicator organisms
- (3) 0.5 ml of metabolic activation mixture (optional)
- (4) Up to 100 µl of a solution of the test chemical.**

For negative controls, we use steps (1), (2), and (3) (optional) and 100 µl of the solvent used for the test chemical.

This mixture is stirred gently and then poured onto minimal agar plates.† After the soft agar has set, the plates are incubated at 37°C for 2 days. The number of his⁺ revertants (colonies that grow on plates lacking a sufficient amount of histidine to support colony formation) are counted and recorded. Some of the revertants are routinely tested to confirm that they are his⁺, require biotin, and are sensitive to crystal violet (rfa⁻).

Escherichia coli WP2

The E. coli WP2 (uvrA⁻) used in this project was given to us by Dr. D. McCalla.^{14,15} A procedure similar to the one used with Salmonella is used to measure the reversion of WP2 to tryptophan independence. However, instead of containing a trace of tryptophan in the top agar, the minimal agar plates contain 1.25 g of oxoid broth per liter to provide

* 0.6% agar contains 0.05 mM histidine and 0.05 mM biotin.

† Minimal agar plates consist of 15 g of agar, 20 g of glucose, 0.2 g of MgSO₄·7 H₂O, 2 g of citric acid monohydrate, 10 g of K₂HPO₄, and 3.5 g of NaH₂PO₄·H₂O per liter.

**Solvents used as appropriate include: water, dimethyl sulfoxide, ethanol, and benzene.

the trace of tryptophan required for enhancement of any mutagenic effect of the test chemical.

Alternatively, reversion of the mutated tryptophan gene, WP2 may undergo a forward mutation in a tryptophan tRNA gene to obtain tryptophan independence. We do not distinguish experimentally between the true revertants and the phenotypic revertants (although the latter tend to form smaller colonies).

Escherichia coli W3110/p3478 and Bacillus subtilis H17/M45

The E. coli strains W3110 and p3478 were obtained from Dr. H. Rosenkrantz.¹⁶ Strain p3478 is a polA⁻ derivative of strain W3110. It carries a single, revertable mutation in a gene for a DNA polymerase; Gross and Gross¹⁷ showed that this mutation is involved in DNA repair synthesis. This mutation increases the sensitivity of strain p3478 to chemicals that lead to alterations (damage) of the DNA. Therefore, we can assay for chemicals that damage DNA by comparing the relative sensitivity of the two strains (p3478 and W3110) to the test chemical.

The B. subtilis strains H17 and M45 were obtained from Dr. Kada.¹⁸ Strain H17 (rec⁺) is derived from H17 but is deficient in the genetic recombination mechanism necessary to repair DNA damage. Cells deficient in this repair mechanism are killed more easily by chemical mutagens than are wild-type cells (rec⁺). If the chemical is toxic to rec⁻ cells, but at the same concentration is not toxic to rec⁺ cells, the chemical probably is a mutagen.

Inoculums from frozen stocks are grown overnight in nutrient broth* at 37°C with shaking. To 2 ml of nutrient broth containing 0.6% agar is added 0.1 ml of the test culture. The suspension is mixed and poured into a petri dish containing nutrient broth and 2% agar.

After the soft agar has solidified, a sterile filter disc impregnated with the test chemical is placed in the center of the plate. The plates are incubated at 37°C for 16 hours, and the width of the zone of

* Tryptone, 1%, and 0.5% yeast extract, supplemented with 5 µg of thymine/ml to prevent selection of thy⁺ revertants.

toxicity or inhibition of growth is then measured. We usually must test several concentrations of chemical to detect accurately differences in the zones of growth inhibition because higher initial concentrations lead to steep concentration gradients that may reduce the differences in growth inhibition of the two strains.

The positive control for this assay is 1 ml of 1-phenyl-3,3-dimethyl-triazene placed on the disc. A zone of approximately 40-mm width is observed (52 and 61 mm, respectively). An additional control is 30 µg of chloramphenicol placed on a disc. Equal zones of inhibition are expected in all four strains (approximately 30 mm) since the toxicity of this chemical does not depend on a mechanism that leads to DNA damage. All assays are performed at least three times.

Saccharomyces cerevisiae D3

The yeast S. cerevisiae D3 is a diploid heterozygous for a mutation in an adenine-metabolizing enzymes.¹⁹ Cells homozygous for this mutation produce a red dye when grown on medium containing adenine. Adenine-requiring homozygotes can be generated from the heterozygotes by mitotic recombination. Many mutagens increase the frequency of mitotic recombination. Mitotic recombination is indicated by the development of colonies with red pigmentation, and the degree of conversion to this pigmented colony indicates the mutagenicity of a compound or its metabolite.²⁰

The Saccharomyces test strain from the liquid nitrogen is grown overnight at 30°C with aeration in 1.0% tryptone and 0.5% yeast extract. The cells are washed twice in 0.067M PO₄ buffer (pH 7.4) and resuspended in the same buffer at a concentration of 10⁸ cells/ml.

The in vitro yeast mitotic recombination assay in suspension consists of 5 x 10⁷ washed, stationary-phase yeast cells in 1 ml of 0.067M PO₄ buffer (pH 7.4) and 50 mg/ml of the test chemical (or a fraction of the concentration required to give 50% killing). The suspension is incubated at 30° for 4 hours. After incubation, the sample is diluted serially in sterile saline and plated on tryptone-yeast-agar plates.

Plates of a 10^{-3} dilution are incubated for 2 days at 30°C, followed by 2 days at 4°C to enhance the development of the red pigment indicative of adenine-negative homozygosity. To detect red colonies or red sectors, we scan the plates with a dissecting microscope at 10 x magnification. Plates of a 10^{-5} dilution are incubated for 2 days at 30°C for determination of the total number of colony-forming units.

The *in vitro* yeast itotic recombination assay in suspension with metabolic activation is conducted as above with the addition of the metabolic activation system to the incubation mixture.

Aroclor 1254-Stimulated Metabolic Activation System

Some carcinogenic mutagens (e.g., dimethylnitrosamine) are inactive unless they are converted to their active form by being metabolized. Ames et al.²¹ have described the metabolic activation systems we use. Adult male mice are given a single 500-mg/kg intraperitoneal injection of a polychlorinated biphenyl (Aroclor 1254).²² Four days after the injection, the animals' food is removed. On the fifth day, the mice are killed.

The liver are removed aseptically and placed in preweighed, sterile glass beakers. The organ weight is determined, and all subsequent operations to the metabolic activation step are conducted in an ice bath. The organ is washed in an equal volume of cold, sterile 0.15 M KCl (1 ml/g of wet organ), minced with sterile surgical scissors in three volume of 0.15 KCl, and homogenized with a Potter-Elvehjem apparatus. The homogenate is centrifuged for 10 minutes at 9000 x g, and the supernatant is removed and stored in liquid nitrogen. To the postmitochondrial supernate are added MgCl₂, KCl, glucose-6-phosphate, TPN, and sodium phosphate (pH 7.4).

Results and Discussion

All the pesticides submitted to SRI for examination were tested at least three times in the microbiological assays. The results presented here are an average of those experiments.

Table 91 presents the results of his reversion assays on agar with Salmonella typhimurium. In this histidine reverse mutation assay system, two pesticides--captan and folpet--were mutagenic. For each chemical, we observed an increase in the number of histidine-independent revertants on strains TA1535 and TA100 but not on strains TA98, TA1537, or TA1538. These results suggest that these pesticides can alkylate DNA, causing mutations of the base-pair substitution type. This conclusion is consistent with the mutagenic activity of these compounds in assays with E. coli WP2 (Table 92), which is sensitive to base-pair substitution mutagens. Although liver homogenate activation was not required for mutagenic activity, the mutagenic activity was enhanced somewhat with activation at some doses. A toxic effect (reduction of the number of mutants) was observed at doses of 100 μ g of each compound.

Table 92 presents the results of assays with E. coli WP2. Essentially, the results were identical to those obtained with S. typhimurium TA1535 and TA100; captan and folpet were mutagenic, but none of the other pesticides was mutagenic.

Table 93 presents the results of the assays for microbial inhibition in repair-deficient and-proficient strains of B. subtilis and E. coli. Folpet, captan, chloropyrifos, and dinoseb all gave toxic zones that were larger on the repair-deficient strains than on the repair-proficient strains, indicating a mutagenic response. Toxic chemicals that do not act by damaging DNA (e.g., chloramphenicol) should give equivalent zones of toxicity. However, many if not all mutagens damage DNA and, if the damage is not repaired, can result in cell death. Thus, a given concentration of mutagen may be toxic for a repair deficient strain but not for a strain the effectively repairs its DNA.

Tables 94 through 113 present the results of the assays for mitotic recombination in Saccharomyces cerevisiae D3. A positive response in this assay is indicated by an increase of more than threefold in the absolute number of mitotic recombinants per milliliter as well as in the relative number of mitotic recombinants per 10^5 survivors. Folpet,

captan, monocrotophos, cacodylic acid, and azinphos-methyl increased mitotic recombination significantly and are considered positive by these procedures. Methyl parathion gave a marginally positive response.

Our results indicate that 7 of the 20 pesticides examined give positive responses in one or more of the four microbiological assay procedures. Although a mutagenic response in a microorganism does not mean that a chemical is a mutagen in humans, the combination of four separate assay systems greatly enhances the probability of detecting potentially hazardous chemicals. Folpet and captan are mutagenic in all four assay procedures. Chlorpyrifos and dinoseb are positive in the microbial sensitivity. Monocrotophos, cacodylic acid, and azinphos-methyl are positive in the yeast assays.

DISCUSSION

Of the 20 pesticides tested for mutagenic activity, 9 were clearly mutagenic in one or more in vitro assays. Of these 9, 2 were mutagenic in all the in vitro assays, but none of them produced a dominant lethal response in the mouse. In the Salmonella assays, these chemicals caused base-pair substitution mutations but not frameshift mutations. The absence of activity in the dominant lethal assay may be due to a lack of sensitivity of the mouse to these types of compounds; for example, N-methyl-N'-nitro-N-nitrosoguanidine and other alkylating agents that cause base-pair substitution mutations do not all cause dominant lethality. Another explanation for the absence of activity may be that these pesticides did not reach the gonadal tissues in sufficient amounts to cause a mutagenic event. None of the other 6 pesticides was mutagenic in all the in vitro assays.

The combination of assays used in this program is one means of identifying those pesticides that may present a mutagenic health hazard. Those that show positive responses in several experimental systems should be evaluated more thoroughly before they are substituted for other pesticides already considered as a risk to the environment. Also apparent is that no one assay system is uniquely capable of detecting the spectrum of mutagenic events that different chemical structures may cause.

REFERENCES

1. W. T. Federer. *Experimental Design (Theory and Application)*. The Macmillan Company, 1955.
2. B. Djordjevic and L. Tolmach. Response of synchronized populations of HeLa cells to ultraviolet irradiation at selected stages of the generation cycle. *Radiat. Res.* 32, 327 (1967).
3. R. E. Reebussen and R. B. Painter. Radiation-stimulated DNA synthesis in cultured mammalian cells. *J. Cell Biol.* 29, 11 (1966).
4. H. Stich and B. A. Laishes. DNA repair and chemical carcinogens. In *Pathobiology Annual*. H. L. Joachim (ed.), Appleton-Century-Crofts, New York, 1973, pp. 341-376.
5. K. Elgjo, H. Hennings, D. Michael, and S. H. Yuspa. Natural synchrony of newborn mouse epidermal cells in vitro. *J. Invest. Derm.* 66, 292-296 (1966).
6. K. Burton. A study of the conditions and mechanism of the diphenylamine reaction for the colorimetric estimation of deoxyribonucleic acid. *Biochem. J.* 62, 315-323 (1956).
7. A. D. Mitchell. Potential Prescreens for Chemical Carcinogens: Unscheduled DNA Synthesis, Task 2. Final report prepared for National Cancer Institute under Contract N01/CP-33394, SRI Project LSU-2735 (April 1976).
8. H. F. Stich, D. Kieser, R.H.C. San, and B. A. Laishes. The Use of DNA repair in the identification of carcinogens, precarcinogens, and target tissue. In *Canadian Cancer Conference: Proceedings of the 1973 Canadian Cancer Conference*, Vol. 10. P. E. Scholefield (ed.), University of Toronto Press, Toronto, Canada, 1973, pp. 83-110.
9. S. Nishida, E. Choi, F. Yamasaki, and B. N. Ames. Detection of carcinogens as mutagens in the *Salmonella* microsome test: Assay of 300 chemicals. *Proc. Nat. Acad. Sci. USA* 72, 5135-5139.
10. B. N. Ames, E. G. Gurney, J. A. Miller, and H. Bartsch. Carcinogens as frameshift mutagens: Metabolites and derivatives of 2-acetylaminofluorene and other aromatic amine carcinogens. *Proc. Nat. Acad. Sci. USA* 69, 3128-3132 (1972).
11. B. N. Ames, F. D. Lee, and W. E. Durston. An improved bacterial test system for the detection and classification of mutagens and carcinogens. *Proc. Nat. Acad. Sci. USA* 70, 782-786 (1973).

12. B. N. Ames, E. W. Durston, K. Yamasaki, and F. D. Lee. Carcinogens are mutagens: A simple test system combining liver homogenates for activation and bacteria for detection. Proc. Nat. Acad. Sci. USA 70, 2281-2285 (1973).
13. J. McCann, N. E. Spingarn, J. Koberl, and B. N. Ames. The detection of carcinogens as mutagens: Bacterial tester strains with R factor plasmids. Proc. Nat. Acad. Sci. USA 72, 979-983 (1975).
14. B. A. Bridges. Simple bacterial systems for detecting mutagenic agents. Lab. Pract. 21, 413-416 (1972).
15. D. R. McCalla and D. Voutsinos. On mutagenicity of nitrofurans. Mutation Res. 26, 3-16 (1974).
16. M. D. Anderson, E. E. Slater, and H. S. Rosenkranz. Rapid detection of mutagens and carcinogens. Cancer Res. 31, 970-973 (1971).
17. J. Gross and M. Gross. Genetic analysis of an *E. coli* strain with a mutation affecting DNA polymerase. Nature 224, 1166-1168 (1969).
18. T. Kada. Mutagenicity testing of chemicals in microbial systems. New Methods in Environ. Chem. Toxicology 127-133 (November 1973).
19. F. K. Zimmermann and R. Schwaier. Induction of mitotic gene conversion with nitrous acid, 1-methyl-3-nitro-1-nitrosoguanidine and other alkylating agents in *Saccharomyces cerevisiae*. Mol. Gen. Genet. 100, 63-69 (1967).
20. D. J. Brusick and V. W. Mayer. New developments in mutagenicity screening techniques with yeast. Environ. Health Perspectives 6, 83-96 (1973).
21. L. D. Kier, E. Yamasaki, and B. N. Ames. Detection of mutagenic activity in cigarette smoke condensates. Proc. Nat. Acad. Sci. USA 71, 4159-4163 (1974).
22. B. N. Ames, J. McCann, and E. Yamasaki. Methods for Detecting Carcinogens and Mutagens with the *Salmonella*/Mammalian-Microsome Mutagenicity Test. Mut. Res. 31, 347-364 (1975).

Table 1

CHI SQUARE TEST OF THE FERTILITY INDEX - MONOCROTOPHOS
1 DEGREE OF FREEDOM

WEEK	VEHICLE CONTROL				74-10 15 MG/KG				74-10 30 MG/KG				74-10 60 MG/KG				TFM .2 MG/KG			
	N PRG	N MYD	FERT. INDEX	CHISQ	N PRG	N MYD	FERT. INDEX	CHISQ	N PRG	N MYD	FERT. INDEX	CHISQ	N PRG	N MYD	FERT. INDEX	CHISQ	N PRG	N MYD	FERT. INDEX	CHISQ
	MULTIPLE TREATMENT																			
1	28	40	.70	0.00	23	40	.57	.67	18	40	.45	4.14	16	40	.40	4.11*	29	40	.72	0.00
2	26	40	.65	0.00	31	40	.77	.98	20	40	.50	1.28	16	40	.40	4.06*	27	39	.69	.03
3	23	40	.57	0.00	30	40	.75	2.01	19	40	.47	.45	22	40	.55	0.00	32	40	.60	3.72
4	27	40	.67	0.00	25	40	.63	.05	22	40	.55	.84	15	40	.38	0.07*	27	40	.67	.06
5	24	40	.60	0.00	33	39	.65	4.79**	25	40	.63	0.00	21	40	.52	.20	30	40	.75	1.42
6	24	38	.63	0.33	27	40	.67	.03	23	40	.57	.08	25	38	.66	0.00	27	38	.71	.24
7	30	38	.79	0.00	25	40	.63	1.81	21	40	.52	4.91*	28	38	.74	.07	27	36	.75	.02
8	27	38	.71	0.00	28	40	.70	.02	20	40	.50	2.75	24	38	.63	.24	26	36	.72	.02

* SIGNIFICANT AT P LT 0.05
I INCREASED ABOVE CONTROL

Table 2
 AVERAGE IMPLANTS PER PREGNANT FEMALE - MONOCROTOPHOS

WEEK	CONTROL	74-10 15 MG/KG	74-10 30 MG/KG	74-10 60 MG/KG	TEM .2 MG/KG
MULTIPLE TREATMENT					
1	319/ 28=11.39	249/ 23=10.83	189/ 18=10.50	180/ 16=11.25	316/ 29=10.90
2	303/ 26=11.65	332/ 31=10.71	242/ 20=12.10	180/ 16=11.25	293/ 27=10.85
3	245/ 23=10.65	356/ 30=11.87	239/ 19=12.58**I	267/ 22=12.14 *I	348/ 32=10.87
4	309/ 27=11.44	314/ 25=12.56	274/ 22=12.45	166/ 15=11.07	266/ 27= 9.85*
5	274/ 24=11.42	372/ 33=11.27	270/ 25=10.80	248/ 21=11.81	354/ 30=11.80
6	302/ 24=12.58	327/ 27=12.11	275/ 23=11.96	255/ 25=10.20 **	273/ 27=10.11**
7	346/ 30=11.53	285/ 25=11.40	255/ 21=12.14	316/ 28=11.29	306/ 27=11.33
8	292/ 27=10.81	313/ 26=11.18	228/ 20=11.40	291/ 24=12.12	322/ 26=12.38**I

* SIGNIFICANT AT P LT 0.05
 ** SIGNIFICANT AT P LT 0.01
 I INCREASED ABOVE CONTROL

Table 3

AVERAGE DEAD IMPLANTS PER PREGNANT FEMALE - MONOCROTOPHOS

WEEK	CONTROL	74-10 15 MG/KG	74-10 30 MG/KG	74-10 60 MG/KG	TEM .2 MG/KG
MULTIPLE TREATMENT					
1	13/ 28= .46	28/ 23= 1.22	10/ 18= .56	10/ 16= .63	62/ 29= 2.14**
2	8/ 26= .31	3/ 31= .10 ^{AD}	10/ 20= .50	2/ 16= .13	77/ 27= 2.85**
3	9/ 23= .39	16/ 30= .53	9/ 19= .47	9/ 22= .41	87/ 32= 2.72**
4	2/ 27= .07	7/ 25= .28*	26/ 22= 1.18*	9/ 15= .60**	11/ 27= .41*
5	11/ 24= .46	22/ 33= .67	12/ 25= .48	9/ 21= .43	22/ 30= .73
6	21/ 24= .88	16/ 27= .59	14/ 23= .61	6/ 25= .24** ^{AD}	17/ 27= .63
7	30/ 30= 1.00	11/ 25= .44	4/ 21= .19	8/ 28= .29	11/ 27= .41
8	19/ 27= .70	24/ 28= .86	7/ 20= .35	9/ 24= .38	11/ 26= .42

* SIGNIFICANT AT P LT 0.05

** SIGNIFICANT AT P LT 0.01

D DECREASED BELOW CONTROL

Table 4

CHI-SQUARE TEST OF THE DEATH INDEX - MONOCROTOPHOS
1 DEGREE OF FREEDOM

WEEK	VEHICLE CONTROL				74-10 15 MG/KG				74-10 30 MG/KG				74-10 60 MG/KG				TEM .2 MG/KG				
	N	N	DEATH	CHISQ	N	N	DEATH	CHISQ	N	N	DEATH	CHISQ	N	N	DEATH	CHISQ	N	N	DEATH	CHISQ	
	WDI	PRG	INDEX		WDI	PRG	INDEX		WDI	PRG	INDEX		WDI	PRG	INDEX		WDI	PRG	INDEX		
MULTIPLE TREATMENT																					
1E	1	10	28	.36	0.00	11	23	.48	.35	7	18	.39	.01	6	16	.38	.04	25	29	.86	13.27 **
	2	7	26	.27	0.00	3	31	.10	1.84	7	20	.35	.07	2	16	.13	.52	26	27	.96	24.26 **
	3	9	23	.39	0.00	11	30	.37	.01	7	19	.37	.03	6	22	.27	.28	25	32	.79	7.05 **
	4	2	27	.07	0.00	7	25	.28	2.54	11	22	.50	9.20 **	9	15	.60	11.21 **	8	27	.70	3.07
	5	9	24	.38	0.00	11	33	.33	.00	12	25	.48	.21	8	21	.38	.07	10	30	.33	.00
	6	15	24	.63	0.00	13	27	.48	.56	11	23	.48	.52	5	25	.20	7.48	10	27	.37	2.36
	7	8	30	.27	0.00	8	25	.32	.02	4	21	.19	.09	7	28	.25	.02	9	27	.33	.07
	8	12	27	.44	0.00	15	28	.54	.17	7	20	.35	.12	8	24	.33	.27	10	26	.38	.63

** SIGNIFICANT AT P LT 0.01

Table 5

NUMBER OF DEAD IMPLANTS PER TOTAL IMPLANTS - MONOCROTOPHOS

WEEK	CONTROL	74-10 15 MG/KG	74-10 30 MG/KG	74-10 60 MG/KG	TEM .2 MG/KG
MULTIPLE TREATMENT					
1	13/ 310 ^a .04	26/ 249 ^a .11	10/ 189 ^a .05	10/ 180 ^a .06	62/ 316 ^a .20 ^{**}
2	8/ 303 ^a .03	3/ 332 ^a .01	10/ 242 ^a .04	2/ 180 ^a .01	77/ 293 ^a .26 ^{**}
3	9/ 245 ^a .04	14/ 356 ^a .04	9/ 239 ^a .04	9/ 267 ^a .03	87/ 348 ^a .25 ^{**}
4	2/ 309 ^a .01	7/ 314 ^a .02	26/ 274 ^a .09 ^{**}	9/ 166 ^a .05 ^{**}	11/ 266 ^a .04 [*]
5	11/ 274 ^a .04	22/ 372 ^a .06	12/ 270 ^a .04	9/ 248 ^a .04	22/ 356 ^a .06
6	21/ 302 ^a .07	16/ 327 ^a .05	14/ 275 ^a .05	6/ 255 ^a .02 ^{**D}	17/ 273 ^a .06
7	30/ 346 ^a .09	11/ 285 ^a .04	4/ 255 ^a .02	8/ 316 ^a .03	11/ 306 ^a .04
8	19/ 292 ^a .07	24/ 319 ^a .08	7/ 228 ^a .03	9/ 291 ^a .03	11/ 322 ^a .03

* SIGNIFICANT AT P LT 0.05

** SIGNIFICANT AT P LT 0.01

D DECREASED BELOW CONTROL

Table 6
 CHI-SQUARE TEST OF THE FERTILITY INDEX - BROMACIL
 1 DEGREE OF FREEDOM

WEEK	VEHICLE CONTROL				74-06 1250 MG/KG				74-06 2500 MG/KG				74-06 5000 MG/KG				TEM .2 MG/KG			
	N	N	FERT.	CHISQ	N	N	FERT.	CHISQ	N	N	FERT.	CHISQ	N	N	FERT.	CHISQ	N	N	FERT.	CHISQ
	PRG	MTD	INDEX		PRG	MTD	INDEX		PRG	MTD	INDEX		PRG	MTD	INDEX		PRG	MTD	INDEX	
MULTIPLE TREATMENT																				
1	26	40	.70	0.00	21	40	.52	1.90	23	40	.57	.87	29	40	.72	0.00	29	40	.72	0.00
2	26	40	.65	0.00	24	40	.60	.05	26	38	.68	.01	25	40	.63	0.00	27	39	.69	.03
3	23	40	.57	0.00	29	40	.72	1.37	19	38	.50	.19	25	40	.63	.05	32	40	.80	3.72
4	27	40	.67	0.00	25	40	.63	.05	22	38	.58	.41	26	40	.70	0.00	27	40	.67	.06
5	24	40	.60	0.00	31	40	.77	2.09	22	38	.58	.00	27	40	.67	.22	30	40	.75	1.42
6	24	38	.63	0.00	31	40	.77	1.30	25	38	.66	0.00	27	40	.67	.03	27	38	.71	.24
7	30	38	.79	0.00	29	40	.72	.16	22	38	.58	2.98	29	40	.72	.16	27	36	.75	.02
8	27	38	.71	0.00	28	39	.72	.03	24	38	.63	.24	27	40	.67	.01	26	36	.72	.02

33

Table 7
 AVERAGE IMPLANTS PER PREGNANT FEMALE - BROMACIL

WEEK	CONTROL	74-06 1250 MG/KG	74-06 2500 MG/KG	74-06 5000 MG/KG	TEM ,2 MG/KG
MULTIPLE TREATMENT					
1	319/ 28=11.39	235/ 21=11.19	278/ 23=12.09	328/ 23=11.31	316/ 29=10.90
2	303/ 26=11.65	253/ 24=10.54	284/ 26=10.92	304/ 25=12.16	293/ 27=10.85
3	245/ 23=10.65	335/ 29=11.55	225/ 19=11.84	319/ 25=12.76 ^{**I}	348/ 32=10.87
4	305/ 27=11.44	290/ 25=11.60	268/ 22=12.18	346/ 28=12.36	266/ 27= 9.85 *
5	274/ 24=11.42	351/ 31=11.65	261/ 22=11.86	323/ 27=11.96	356/ 30=11.87
6	302/ 24=12.58	368/ 31=11.87	294/ 25=11.76	276/ 27=10.30 ^{**}	273/ 27=10.11 ^{**}
7	346/ 30=11.53	355/ 29=12.24	253/ 22=11.50	357/ 26=12.31	306/ 27=11.33
8	292/ 27=10.81	318/ 28=11.36	277/ 24=11.54	307/ 27=11.37	322/ 26=12.38 ^{**I}

* SIGNIFICANT AT P LT 0.05
 ** SIGNIFICANT AT P LT 0.01
 I INCREASED ABOVE CONTROL

Table 8

AVERAGE DEAD IMPLANTS PER PREGNANT FEMALE - BROMACIL

WEEK	CONTROL	74-06 1250 MG/KG	74-06 2500 MG/KG	74-06 5000 MG/KG	TEM .2 MG/KG
MULTIPLE TREATMENT					
1	13/ 28= .46	8/ 21= .38	10/ 23= .43	17/ 29= .59	62/ 29= 2.14**
2	8/ 26= .31	12/ 24= .50	8/ 26= .31	5/ 25= .20	77/ 27= 2.85**
3	9/ 23= .39	6/ 29= .21	12/ 19= .63	10/ 25= .40	87/ 32= 2.72**
4	2/ 27= .07	7/ 25= .28	16/ 22= .73**	15/ 28= .54*	11/ 27= .41*
5	11/ 24= .46	12/ 31= .39	15/ 22= .68	8/ 27= .30	22/ 30= .73
6	21/ 24= .88	7/ 31= .23**D	16/ 25= .64	21/ 27= .78	17/ 27= .63
7	30/ 30= 1.00	11/ 29= .38	14/ 22= .64	14/ 29= .48	11/ 27= .41
8	19/ 27= .70	15/ 28= .54	8/ 24= .33	9/ 27= .33	11/ 26= .42

* SIGNIFICANT AT P < 0.05
 ** SIGNIFICANT AT P < 0.01
 D DECREASED BELOW CONTROL

35

Table 9
 CHI-SQUARE TEST OF THE DEATH INDEX - BROMACIL
 1 DEGREE OF FREEDOM

WEEK	VEHICLE CONTROL				74-06 1250 MG/KG				74-06 2500 MG/KG				74-06 5000 MG/KG				TEM .2 MG/KG				
	N	N	DEATH	CHISQ	N	N	DEATH	CHISQ	N	N	DEATH	CHISQ	N	N	DEATH	CHISQ	N	N	DEATH	CHISQ	
	WDI	PRG	INDEX		WDI	PRG	INDEX		WDI	PRG	INDEX		WDI	PRG	INDEX		WDI	PRG	INDEX		
MULTIPLE TREATMENT																					
36	1	10	28	.36	0.00	7	21	.33	.02	7	23	.30	.01	14	29	.48	.48	25	29	.86	13.27**
	2	7	26	.27	0.00	10	24	.42	.64	8	26	.31	0.00	5	25	.20	.06	26	27	.96	24.26**
	3	9	23	.39	0.00	6	29	.21	1.32	9	19	.47	.05	6	25	.24	.67	25	32	.78	7.09**
	4	2	27	.07	0.00	6	25	.24	1.02	11	22	.50	9.20**	14	28	.50	10.11**	8	27	.30	3.07
	5	9	24	.38	0.00	12	31	.39	.04	12	22	.55	.74	8	27	.30	.09	10	30	.33	.00
	6	15	24	.63	0.00	5	31	.16	10.65**D	13	25	.52	.21	10	27	.37	2.36	10	27	.37	2.76
	7	8	30	.27	0.00	7	29	.24	.01	10	22	.45	1.24	12	29	.41	.84	9	27	.33	.07
	8	12	27	.44	0.00	11	28	.39	.01	7	24	.29	.70	7	27	.26	1.30	10	26	.38	.03

** SIGNIFICANT AT P LT 0.01
 D DECREASED BELOW CONTROL

Table 10

NUMBER OF DEAD IMPLANTS PER TOTAL IMPLANTS - BROMACIL

WEEK	CONTROL	74-06 1250 MG/KG	74-06 2500 MG/KG	74-06 5000 MG/KG	TEM .2 MG/KG
MULTIPLE TREATMENT					
1	13/ 319= .04	8/ 235= .03	10/ 278= .04	17/ 328= .05	62/ 316= .20**
2	8/ 303= .03	12/ 253= .05	8/ 284= .03	5/ 304= .02	77/ 293= .26**
3	9/ 245= .04	6/ 335= .02 *D	12/ 225= .05	10/ 319= .03	87/ 348= .25**
4	2/ 309= .01	7/ 290= .02	16/ 268= .06**	15/ 340= .04**	11/ 266= .04*
5	11/ 274= .04	12/ 361= .03	15/ 261= .06	8/ 323= .02	22/ 356= .06
6	21/ 302= .07	7/ 368= .02 **D	16/ 294= .05	21/ 278= .08	17/ 273= .06
7	30/ 346= .09	11/ 355= .03	14/ 253= .06	14/ 357= .04	11/ 306= .04
8	19/ 292= .07	15/ 318= .05	8/ 277= .03	9/ 307= .03	11/ 322= .03

37

* SIGNIFICANT AT P LT 0.05

** SIGNIFICANT AT P LT 0.01

D DECREASED BELOW CONTROL

Table 11

CHI-SQUARE TEST OF THE FERTILITY INDEX - CAPTAN
1 DEGREE OF FREEDOM

WEEK	VEHICLE CONTROL				7*-02 1250 MG/KG				7*-02 2500 MG/KG				7*-02 5000 MG/KG				TEM .3 MG/KG				
	N	N	FERT.	CHISQ	N	N	FERT.	CHISQ	N	N	FERT.	CHISQ	N	N	FERT.	CHISQ	N	N	FERT.	CHISQ	
	PRG	MTD	INDEX		PRG	MTD	INDEX		PRG	MTD	INDEX		PRG	MTD	INDEX		PRG	MTD	INDEX		
MULTIPLE TREATMENT																					
38	1	23	40	.57	0.00	34	40	.85	6.10* ₁	31	40	.77	2.79	29	40	.72	1.37	29	40	.77	1.37
	2	27	39	.69	0.00	30	40	.75	.10	36	40	.90	4.07* ₁	30	40	.75	.10	27	39	.69	.06
	3	26	40	.65	0.00	23	40	.57	.21	30	40	.75	.54	31	40	.77	.98	32	40	.80	1.57
	4	27	40	.67	0.00	31	38	.82	1.35	33	40	.82	1.67	32	40	.80	1.03	27	40	.67	.06
	5	29	40	.72	0.00	26	38	.68	.02	30	40	.75	0.00	26	38	.68	.02	30	40	.75	0.00
	6	29	40	.72	0.00	27	38	.71	.01	34	40	.85	1.20	30	38	.79	.16	27	38	.71	.01
	7	29	40	.72	0.00	27	38	.71	.01	30	40	.75	0.00	26	38	.68	.02	27	36	.75	.00
	8	32	40	.80	0.00	33	38	.87	.26	34	40	.85	.09	27	38	.71	.43	26	36	.72	.28

* SIGNIFICANT AT FLT 0.05
1 INCREASED ABOVE CONTROL

Table 12
 AVERAGE IMPLANTS PER PREGNANT FEMALE - CAPTAN

WEEK	CONTROL	74-02 1250 MG/KG	74-02 2500 MG/KG	74-02 5000 MG/KG	TEM .2 MG/KG
MULTIPLE TREATMENT					
1	265/ 23=11.52	387/ 34=11.38	339/ 31=10.64	310/ 29=10.69	316/ 29=10.90
2	305/ 27=11.30	328/ 30=10.93	375/ 36=10.42	342/ 30=11.40	293/ 27=10.85
3	268/ 26=10.31	252/ 23=10.96	313/ 30=10.43	345/ 31=11.13	344/ 32=10.75
4	288/ 27=10.67	324/ 31=10.45	359/ 33=10.88	333/ 32=10.41	264/ 27= 9.85
5	334/ 29=11.52	282/ 26=10.85	348/ 30=11.60	298/ 26=11.46	356/ 30=11.87
6	323/ 29=11.14	312/ 27=11.56	361/ 34=10.62	340/ 30=11.33	273/ 27=10.11
7	323/ 29=11.14	304/ 27=11.44	329/ 30=10.97	309/ 26=11.88	304/ 27=11.33
8	381/ 32=11.91	382/ 33=11.58	383/ 34=11.26	301/ 27=11.15	322/ 26=12.38

39

* SIGNIFICANT AT P LT 0.05
 ** SIGNIFICANT AT P LT 0.01

Table 13
 AVERAGE DEAD IMPLANTS PER PREGNANT FEMALE - CAPTAN

WEEK	CONTROL	74-02 1250 MG/KG	74-02 2500 MG/KG	74-02 5000 MG/KG	TEM .2 MG/KG
MULTIPLE TREATMENT					
1	8/ 23* .35	17/ 34* .50	19/ 31* .61	15/ 24* .52	62/ 29* 2.14**
2	11/ 27* .41	15/ 30* .50	17/ 36* .47	14/ 30* .47	77/ 27* 2.85**
3	16/ 28* .42	16/ 23* .70	16/ 30* .53	15/ 31* .48	67/ 32* 2.72**
4	16/ 27* .59	12/ 31* .39	15/ 33* .45	11/ 32* .34	11/ 27* .41
5	15/ 24* .52	24/ 26* .92	9/ 30* .30	11/ 26* .42	22/ 30* .73
6	9/ 24* .31	13/ 27* .48	21/ 34* .62	18/ 36* .60	17/ 27* .63
7	21/ 29* .72	15/ 27* .56	14/ 30* .47	8/ 24* .31	11/ 27* .41
8	13/ 32* .41	12/ 33* .36	11/ 34* .32	7/ 27* .26	11/ 26* .42

40

* SIGNIFICANT AT P LT 0.05
 ** SIGNIFICANT AT P LT 0.01

Table 14
 CHI-SQUARE TEST OF THE DEATH INDEX - CAPTAN
 1 DEGREE OF FREEDOM

WEEK	VEHICLE CONTROL				74-02 1250 MG/KG				74-02 2500 MG/KG				74-02 5000 MG/KG				TEM .2 MG/KG				
	N WDI	N PRG	DEATH INDEX	CHISO	N WDI	N PRG	DEATH INDEX	CHISO	N WDI	N PRG	DEATH INDEX	CHISO	N WDI	N PRG	DEATH INDEX	CHISO	N WDI	N PRG	DEATH INDEX	CHISO	
MULTIPLE TREATMENT																					
17	1	8	23	.35	0.00	10	34	.29	.02	12	31	.39	.00	12	29	.41	.04	25	29	.46	12.49 **
	2	9	27	.33	0.00	8	30	.27	.07	12	36	.33	.07	10	30	.33	.08	26	27	.46	20.79 **
	3	10	25	.38	0.00	12	23	.52	.46	11	30	.37	.02	13	31	.42	.00	25	32	.70	7.85 **
	4	11	27	.41	0.00	8	31	.26	.86	7	33	.21	1.85	8	32	.25	1.02	8	27	.30	.32
	5	11	29	.38	0.00	10	26	.38	.06	7	30	.23	.27	11	26	.42	.00	10	30	.33	.01
	6	8	29	.28	0.00	9	27	.33	.03	15	34	.44	1.20	10	30	.33	.04	10	27	.37	.22
	7	12	29	.41	0.00	12	27	.44	.00	11	30	.37	.01	7	26	.27	.71	9	27	.33	.12
	8	11	32	.34	0.00	8	33	.24	.39	9	34	.26	.19	9	27	.19	1.15	10	26	.36	.00

** SIGNIFICANT AT PLI 0.01

Table 15
NUMBER OF DEAD IMPLANTS PER TOTAL IMPLANTS - CAPTAN

WEEK	CONTROL	74-02 1250 MG/KG	74-02 2500 MG/KG	74-02 5000 MG/KG	TEN 0.2 MG/KG
MULTIPLE TREATMENT					
1	8/ 265# .03	17/ 367# .04	19/ 339# .06	15/ 310# .05	62/ 316# .20**
2	11/ 305# .04	15/ 328# .05	17/ 375# .05	14/ 342# .04	77/ 293# .26**
3	16/ 268# .06	16/ 252# .06	16/ 313# .05	15/ 345# .04	87/ 348# .25**
4	10/ 288# .05	12/ 324# .04	15/ 359# .04	11/ 333# .03	11/ 266# .04
5	15/ 334# .04	24/ 282# .09	9/ 348# .03	11/ 298# .04	22/ 356# .06
6	9/ 323# .03	13/ 312# .04	21/ 361# .06*	10/ 340# .05	17/ 273# .06
7	21/ 323# .07	15/ 309# .05	14/ 329# .04	5/ 309# .03	11/ 366# .04
8	15/ 381# .03	12/ 382# .03	11/ 383# .03	7/ 301# .02	11/ 322# .03

* SIGNIFICANT AT P LT 0.05
** SIGNIFICANT AT P LT 0.01

Table 16

CHI-SQUARE TEST OF THE FERTILITY INDEX - FOLPET
1 DEGREE OF FREEDOM

43

WEEK	VEHICLE CONTROL				74-03 1250 MG/KG				74-03 2500 MG/KG				74-03 5000 MG/KG				TEM .2MG/KG			
	N PRG	N MTD	FERT. INDEX	CHISQ	N PRG	N MTD	FERT. INDEX	CHISQ	N PRG	N MTD	FERT. INDEX	CHISQ	N PRG	N MTD	FERT. INDEX	CHISQ	N PRG	N MTD	FERT. INDEX	CHISQ
MULTIPLE TREATMENT																				
1	23	40	.57	0.00	27	40	.67	.48	29	40	.72	1.37	34	40	.75	2.01	29	40	.72	1.37
2	27	39	.69	0.00	26	40	.70	.03	23	40	.57	.72	35	40	.88	2.90	27	39	.69	.06
3	26	40	.65	0.00	25	40	.63	0.00	31	40	.77	.98	32	40	.80	1.57	32	40	.80	1.57
4	27	40	.67	0.00	30	40	.75	.24	18	40	.45	3.25	30	40	.75	.24	27	40	.67	.06
5	29	40	.72	0.00	30	40	.75	0.00	26	40	.65	.23	35	40	.88	1.95	30	40	.75	0.00
6	29	40	.72	0.00	30	40	.75	0.00	28	40	.70	0.00	28	40	.73	0.00	27	38	.71	.01
7	29	40	.72	0.00	28	40	.70	0.00	27	40	.67	.06	31	40	.77	.07	27	40	.75	.00
8	32	40	.80	0.00	27	40	.67	1.03	29	40	.72	.24	30	40	.75	.07	26	36	.72	.28

Table 17
 AVERAGE IMPLANTS PER PREGNANT FEMALE - FOLPEL

WEEK	CONTROL	74-03 1250 MG/KG	74-03 2500 MG/KG	74-33 5000 MG/KG	124 25MG/KG
MULTIPLE TREATMENT					
1	265/ 23=11.52	308/ 27=11.41	324/ 29=11.17	328/ 30=10.93	316/ 29=10.90
2	305/ 27=11.30	304/ 28=10.86	257/ 23=11.17	390/ 35=11.14	293/ 27=10.85
3	262/ 26=10.31	271/ 25=10.84	334/ 31=10.77	327/ 32=10.22	348/ 32=10.87
4	268/ 27=10.67	294/ 30= 9.97	183/ 18=10.17	320/ 30=10.67	266/ 27= 9.85
5	334/ 29=11.52	333/ 30=11.10	318/ 26=12.23	399/ 35=11.40	356/ 30=11.87
6	323/ 29=11.14	334/ 30=11.13	294/ 26=10.50	312/ 28=11.14	273/ 27=10.11
7	323/ 29=11.14	307/ 28=10.96	311/ 27=11.52	356/ 31=11.48	305/ 27=11.33
8	361/ 32=11.91	301/ 27=11.15	345/ 29=11.90	346/ 30=11.53	322/ 26=12.38

* SIGNIFICANT AT P LT 0.05
 ** SIGNIFICANT AT P LT 0.01

Table 16
 AVERAGE DEAD IMPLANTS PER PREGNANT FEMALE - POLPET

WEEK	CONTROL	74-03 1250 MG/KG	74-03 2500 MG/KG	74-03 5000 MG/KG	TEM .2MG/KG
MULTIPLE TREATMENT					
1	8/ 23 ^a .35	10/ 27 ^a .37	25/ 29 ^a .86	13/ 30 ^a .43	62/ 29 ^a 2.14 ^{**}
2	11/ 27 ^a .41	15/ 28 ^a .64	20/ 23 ^a .87 [*]	21/ 35 ^a .60	77/ 27 ^a 2.85 ^{**}
3	16/ 26 ^a .62	13/ 25 ^a .52	19/ 31 ^a .61	11/ 32 ^a .34	87/ 32 ^a 2.72 ^{**}
4	16/ 27 ^a .59	16/ 30 ^a .53	8/ 18 ^a .44	7/ 30 ^a .23 ^{+D}	11/ 27 ^a .41
5	15/ 29 ^a .52	12/ 30 ^a .40	9/ 26 ^a .35	16/ 35 ^a .46	22/ 30 ^a .73
6	9/ 29 ^a .31	17/ 30 ^a .57	16/ 28 ^a .57	12/ 28 ^a .42	17/ 27 ^a .62
7	21/ 29 ^a .72	14/ 28 ^a .43	10/ 27 ^a .37	18/ 31 ^a .58	11/ 27 ^a .41
8	13/ 32 ^a .41	11/ 27 ^a .41	20/ 29 ^a .69	13/ 30 ^a .43	11/ 26 ^a .42

* SIGNIFICANT AT P LT 0.05
 ** SIGNIFICANT AT P LT 0.01
 D DECREASED BELOW CONTROL

45

Table 19
 CHI-SQUARE TEST OF THE DEATH INDEX - FOLPET
 1 DEGREE OF FREEDOM

WEEK	VEHICLE CONTROL				74-03 1250 MG/KG				74-03 2500 MG/KG				74-03 5005 MG/KG				TEM .2MG/KG				
	N	N	DEATH	CHISQ	N	N	DEATH	CHISQ	N	N	DEATH	CHISQ	N	N	DEATH	CHISQ	N	N	DEATH	CHISQ	
	#01	PRG	INDEX		#01	PRG	INDEX		#01	PRG	INDEX		#01	PRG	INDEX		#01	PRG	INDEX		
MULTIPLE TREATMENT																					
97	1	8	23	.35	0.00	8	27	.30	.41	13	29	.45	.20	10	30	.33	.03	25	29	.86	12.49**
	2	9	27	.33	0.00	13	28	.48	.51	12	23	.52	1.12	15	35	.43	.25	26	27	.96	20.79**
	3	10	26	.38	0.00	9	25	.36	.01	9	31	.29	.22	8	32	.25	.67	25	22	.78	7.85**
	4	11	27	.41	0.00	11	30	.37	.00	7	18	.39	.03	7	30	.23	1.27	8	27	.30	.32
	5	11	29	.38	0.00	10	30	.33	.01	8	26	.31	.47	12	35	.34	.90	10	30	.33	.01
	6	8	29	.28	0.00	11	30	.37	.42	11	28	.39	.43	9	28	.32	.01	10	27	.37	.22
	7	12	29	.41	0.00	8	28	.29	.54	9	27	.33	.12	13	31	.42	.05	9	27	.33	.12
	8	11	32	.34	0.00	9	27	.33	.04	10	29	.35	1.89	11	30	.37	.01	10	28	.38	.00

** SIGNIFICANT AT PLT 0.01

Table 20

NUMBER OF DEAD IMPLANTS PER TOTAL IMPLANTS - FOLPET

WEEK	CONTROL	74-03 1250 MG/KG	74-03 2500 MG/KG	74-03 5000 MG/KG	TEM .2MG/KG
MULTIPLE TREATMENT					
1	8/ 265* .03	10/ 308* .03	25/ 324* .08	13/ 328* .04	62/ 314* .20**
2	11/ 305* .04	18/ 304* .06	20/ 257* .08*	21/ 390* .05	77/ 293* .26**
3	16/ 268* .06	13/ 271* .05	19/ 334* .06	11/ 327* .03	87/ 348* .25**
4	16/ 288* .06	16/ 294* .05	8/ 183* .04	7/ 320* .02	11/ 266* .04
5	15/ 334* .04	12/ 333* .04	9/ 318* .02	16/ 399* .04	22/ 356* .06
6	9/ 323* .03	17/ 334* .05	16/ 294* .05	13/ 312* .04	17/ 273* .06
7	21/ 323* .07	12/ 307* .04	10/ 311* .03	18/ 356* .05	11/ 306* .04
8	13/ 381* .03	11/ 301* .04	20/ 345* .06	13/ 346* .04	11/ 322* .03

* SIGNIFICANT AT P LT 0.05

** SIGNIFICANT AT P LT 0.01

Table 21
 CHI-SQUARE TEST OF THE FERTILITY INDEX - AZINPHOS-METHYL
 1 DEGREE OF FREEDOM

WEEK	VEHICLE CONTROL				74-09 20 MG/KG				74-09 40 MG/KG				74-09 80 MG/KG				TEMP 12 MG/KG				
	N PRG	N MID	FERT. INDEX	CHISQ	N PRG	N MID	FERT. INDEX	CHISQ	N PRG	N MID	FERT. INDEX	CHISQ	N PRG	N MID	FERT. INDEX	CHISQ	N PRG	N MID	FERT. INDEX	CHISQ	
MULTIPLE TREATMENT																					
87	1	28	40	.70	0.00	34	40	.85	1.79	28	40	.70	.06	27	40	.67	0.00	29	40	.72	0.00
	2	26	40	.65	0.00	29	40	.72	.23	27	40	.67	0.00	27	40	.67	0.00	27	39	.69	.03
	3	23	40	.57	0.00	33	40	.82	4.82*†	29	40	.72	1.37	24	38	.63	.08	32	40	.80	3.72
	4	27	40	.67	0.00	30	40	.75	.24	29	40	.72	.06	21	36	.58	.35	27	40	.67	.06
	5	24	40	.60	0.00	29	40	.72	.89	26	40	.65	.05	20	34	.59	.02	30	40	.75	1.42
	6	24	39	.63	0.00	30	40	.75	.79	27	40	.67	.03	28	34	.82	2.41	27	38	.71	.24
	7	30	38	.79	0.00	27	40	.67	.70	25	40	.63	1.81	24	34	.71	.30	27	36	.75	.02
	8	27	35	.71	0.00	26	40	.65	.11	26	40	.65	.11	22	34	.65	.10	26	36	.72	.02

* SIGNIFICANT AT P LT 0.05
 † INCREASED ABOVE CONTROL

Table 22
 AVERAGE IMPLANTS PER PREGNANT FEMALE - AZINPHOS-METHYL

WEEK	CONTROL	74-09 20 MG/KG	74-09 40 MG/KG	74-09 80 MG/KG	TEM .2 MG/KG
MULTIPLE TREATMENT					
1	319/ 28=11.39	375/ 34=11.03	300/ 28=10.71	306/ 27=11.33	316/ 29=10.90
2	303/ 26=11.65	335/ 29=11.55	312/ 27=11.56	310/ 27=11.48	293/ 27=10.85
3	245/ 23=10.65	379/ 33=11.48	338/ 29=11.66	272/ 24=11.33	348/ 32=10.87
4	309/ 27=11.44	367/ 30=12.23	369/ 29=12.72 ^{**I}	247/ 21=11.76	266/ 27= 9.85 *
5	274/ 24=11.42	337/ 29=11.62	310/ 26=11.92	228/ 20=11.40	356/ 30=11.87
6	302/ 24=12.58	345/ 30=11.50 *	276/ 27=10.22 ^{**}	323/ 28=11.54 *	279/ 27=10.11 **
7	346/ 30=11.53	288/ 27=10.67	278/ 25=11.12	262/ 24=10.92	304/ 27=11.33
8	292/ 27=10.81	306/ 26=11.77	293/ 26=11.27	250/ 22=11.36	322/ 26=12.38 ^{**I}

* SIGNIFICANT AT P LT 0.05
 ** SIGNIFICANT AT P LT 0.01
 I INCREASED ABOVE CONTROL

49

Table 23

AVERAGE DEAD IMPLANTS PER PREGNANT FEMALE - AZINPHOS-METHYL

WEEK	CONTROL	74-04 20 MG/KG	74-05 40 MG/KG	74-09 80 MG/KG	TEM 12 MG/KG
MULTIPLE TREATMENT					
1	13/ 28 ^a .46	13/ 34 ^a .38	34/ 28 ^a 1.21 *	19/ 27 ^a .70	62/ 29 ^a 2.14 **
2	8/ 26 ^a .31	16/ 25 ^a .55	15/ 27 ^a .56	19/ 27 ^a .70**	77/ 27 ^a 2.85 **
3	9/ 23 ^a .39	31/ 33 ^a .94	8/ 29 ^a .28	11/ 24 ^a .46	87/ 32 ^a 2.72 **
4	2/ 27 ^a .07	21/ 30 ^a .70**	31/ 29 ^a 1.07 **	7/ 21 ^a .33*	11/ 27 ^a .41 *
5	11/ 24 ^a .46	15/ 29 ^a .52	6/ 26 ^a .23	11/ 20 ^a .55	22/ 30 ^a .73
6	21/ 24 ^a .69	16/ 36 ^a .33**D	17/ 27 ^a .63	9/ 28 ^a .32**D	17/ 27 ^a .63
7	30/ 30 ^a 1.00	8/ 27 ^a .30	8/ 25 ^a .32	19/ 24 ^a .79	11/ 27 ^a .41
8	19/ 27 ^a .70	13/ 26 ^a .50	10/ 26 ^a .38	6/ 22 ^a .36	11/ 26 ^a .42

* SIGNIFICANT AT P LT 0.05

** SIGNIFICANT AT P LT 0.01

D DECREASED BELOW CONTROL

Table 2-
 CHI-SQUARE TEST OF THE DEATH INDEX - AZINPHOS-METHYL
 1 DEGREE OF FREEDOM

WEEK	VEHICLE CONTROL				74-09 20 MG/KG				74-09 40 MG/KG				74-09 80 MG/KG				TSM .2 MG/KG			
	N	N	DEATH	CHISQ	N	N	DEATH	CHISQ	N	N	DEATH	CHISQ	N	N	DEATH	CHISQ	N	N	DEATH	CHISQ
	WDI	PRG	INDEX		WDI	PRG	INDEX		WDI	PRG	INDEX		WDI	PRG	INDEX		WDI	PRG	INDEX	
MULTIPLE TREATMENT																				
1	10	28	.36	0.00	12	34	.35	.05	14	28	.50	.66	11	27	.41	.01	25	29	.46	13.27**
2	7	26	.27	0.00	11	29	.38	.34	8	27	.30	.01	12	27	.44	1.09	26	27	.96	24.26**
3	9	23	.39	0.00	16	33	.48	.18	5	29	.17	2.11	8	24	.33	.01	25	32	.78	7.05**
4	2	27	.07	0.00	13	30	.43	7.70**	14	29	.48	9.53**	7	21	.33	3.65	8	27	.30	3.07
5	9	24	.38	0.00	13	29	.45	.07	6	26	.23	.64	7	20	.35	.02	10	30	.33	.06
6	15	24	.63	0.00	8	30	.27	5.61**D	9	27	.33	3.25	8	28	.29	4.73**D	10	27	.37	2.36
7	8	30	.27	0.00	7	27	.26	.06	7	25	.28	.04	10	24	.42	.76	9	27	.33	.07
8	12	27	.44	0.00	10	26	.38	.03	8	26	.31	.55	8	22	.36	.08	10	26	.38	.03

* SIGNIFICANT AT P LT 0.05
 ** SIGNIFICANT AT P LT 0.01
 D DECREASED BELOW CONTROL

51

Table 25

NUMBER OF DEAD IMPLANTS PER TOTAL IMPLANTS - AZINPHOS-METHYL

WEEK	CONTROL	74-09 20 MG/KG	74-09 40 MG/KG	74-09 80 MG/KG	TECH .2 MG/KG
MULTIPLE TREATMENT					
1	13/ 319= .04	13/ 375= .03	34/ 300= .11	19/ 306= .06	62/ 316= .20 **
2	5/ 303= .03	16/ 335= .05	15/ 312= .05	19/ 310= .06	77/ 293= .26 **
3	9/ 245= .04	31/ 379= .08	8/ 338= .02	11/ 272= .04	87/ 348= .25 **
4	2/ 309= .01	21/ 367= .06 **	31/ 369= .08 **	7/ 247= .03*	11/ 266= .04 *
5	11/ 276= .04	15/ 337= .04	6/ 310= .02	11/ 225= .05	22/ 356= .06
6	21/ 302= .07	10/ 345= .03 **D	17/ 276= .06	9/ 323= .03**D	17/ 273= .06
7	30/ 346= .09	8/ 288= .03	8/ 278= .03	19/ 262= .07	11/ 306= .04
8	19/ 292= .07	13/ 306= .04	10/ 293= .03	8/ 250= .03	11/ 322= .03

52

* SIGNIFICANT AT P LT 0.05
 ** SIGNIFICANT AT P LT 0.01
 D DECREASED BELOW CONTROL

Table 26

CHI-SQUARE TEST OF THE FERTILITY INDEX - MALATHION
1 DEGREE OF FREEDOM

WEEK	VEHICLE CONTROL				74-07 1250 MG/KG				74-07 2500 MG/KG				74-07 5000 MG/KG				TEM .2 MG/KG				
	N PRG	N MTD	FERT. INDEX	CHISQ	N PRG	N MTD	FERT. INDEX	CHISQ	N PRG	N MTD	FERT. INDEX	CHISQ	N PRG	N MTD	FERT. INDEX	CHISQ	N PRG	N MTD	FERT. INDEX	CHISQ	
MULTIPLE TREATMENT																					
53	1	28	40	.70	0.00	31	40	.77	.26	26	40	.65	.06	29	40	.72	0.00	29	40	.72	0.00
	2	26	40	.65	0.00	30	40	.75	.54	24	40	.60	.05	31	39	.79	1.40	27	39	.69	.03
	3	23	40	.57	0.00	23	38	.61	.00	24	40	.60	0.00	33	40	.52	4.82*I	32	40	.40	3.72
	4	27	40	.67	0.00	25	38	.66	.01	22	38	.58	.41	27	40	.67	.06	27	40	.67	.06
	5	24	40	.60	0.00	21	36	.58	.01	22	36	.61	.02	34	40	.85	5.08*I	30	40	.75	1.42
	6	24	38	.63	0.00	28	36	.70	1.26	19	34	.56	.15	35	40	.80	5.02*I	27	38	.71	.24
	7	30	38	.79	0.00	30	36	.83	.03	19	34	.56	3.39	32	40	.80	.03	27	26	.75	.02
	8	27	38	.71	0.00	30	36	.83	.96	16	34	.47	3.36	37	40	.92	4.72*I	26	36	.72	.02

** SIGNIFICANT AT P LT 0.05
I INCREASED ABOVE CONTROL

Table 27

AVERAGE IMPLANTS PER PREGNANT FEMALE - MALATHION

WEEK	CONTROL	74-07 1250 MG/KG	74-07 2500 MG/KG	74-07 5000 MG/KG	1EM .2 MG/KG
MULTIPLE TREATMENT					
1	319/ 28=11.39	346/ 31=11.16	298/ 26=11.46	321/ 29=11.07	316/ 29=10.90
2	333/ 26=11.65	363/ 30=12.10	267/ 24=11.12	356/ 31=11.48	293/ 27=10.85
3	273/ 23=10.65	281/ 23=12.22 *I	305/ 24=12.71 *I	402/ 33=12.18 *I	348/ 32=10.87
4	359/ 27=11.44	304/ 25=12.16	256/ 22=11.64	300/ 27=11.11	266/ 27= 9.85*
5	274/ 24=11.42	238/ 21=11.33	243/ 22=11.05	386/ 34=11.35	356/ 30=11.87
6	302/ 24=12.58	319/ 29=11.34 *	226/ 19=11.89	391/ 35=11.17 **	273/ 27=10.11**
7	346/ 30=11.53	337/ 30=11.23	235/ 19=12.37 *I	352/ 32=11.00	306/ 27=11.33
8	292/ 27=10.81	323/ 30=10.77	187/ 16=11.69	397/ 37=10.73	322/ 26=12.38**I

* SIGNIFICANT AT P LT 0.05

** SIGNIFICANT AT P LT 0.01

I INCREASED ABOVE CONTROL

Table 28

AVERAGE DEAD IMPLANTS PER PREGNANT FEMALE - MALATHION

WEEK	CONTROL	74-07 1250 MG/KG	74-07 2500 MG/KG	74-07 5000 MG/KG	TEM .2 MG/KG	
MULTIPLE TREATMENT						
55	1	13/ 28= .46	18/ 31= .58	6/ 26= .31	13/ 29= .45	62/ 29= 2.14**
	2	8/ 26= .31	12/ 30= .40	11/ 24= .46	11/ 31= .35	77/ 27= 2.85**
	3	9/ 23= .39	14/ 23= .61	14/ 24= .58	14/ 33= .42	87/ 32= 2.72**
	4	2/ 27= .07	14/ 25= .56**	8/ 22= .36*	7/ 27= .26	11/ 27= .41*
	5	11/ 24= .46	4/ 21= .19	8/ 22= .36	8/ 34= .24	22/ 30= .73
	6	21/ 24= .88	21/ 28= .75	20/ 19= 1.05	23/ 35= .66	17/ 27= .63
	7	30/ 38= 1.00	10/ 30= .33	6/ 19= .32	15/ 32= .47	11/ 27= .41
	8	19/ 27= .70	19/ 30= .63	7/ 16= .44	32/ 37= .86	11/ 26= .42

* SIGNIFICANT AT P LT 0.05

** SIGNIFICANT AT P LT 0.01

Table 29

CHI-SQUARE TEST OF THE DEATH INDEX - MALATNION
1 DEGREE OF FREEDOM

WEEK	VEHICLE CONTROL				74-07 1250 MG/KG				74-07 2500 MG/KG				74-07 5000 MG/KG				72M .2 MG/KG			
	N	N	DEATH	CHISQ	N	N	DEATH	CHISQ	N	N	DEATH	CHISQ	N	N	DEATH	CHISQ	N	N	DEATH	CHISQ
	#DI	PRG	INDEX		#DI	PRG	INDEX		#DI	PRG	INDEX		#DI	PRG	INDEX		#DI	PRG	INDEX	
MULTIPLE TREATMENT																				
1	10	28	.36	0.00	13	31	.42	.05	7	26	.27	.16	10	29	.34	.03	25	29	.86	13.27 **
2	7	26	.27	0.00	8	30	.27	.08	9	24	.38	.25	8	31	.26	.04	26	27	.96	24.26 **
3	9	23	.39	0.00	7	23	.30	.16	11	24	.46	.03	9	33	.27	.41	25	32	.78	7.05 **
4	2	27	.07	0.00	10	25	.40	6.04*	5	22	.23	1.24	5	27	.19	.66	8	27	.30	3.07
5	9	24	.39	0.00	3	21	.14	2.01	6	22	.27	.18	7	34	.21	1.26	10	30	.33	.00
6	15	24	.63	0.00	11	28	.39	1.93	6	19	.32	2.91	16	35	.46	1.01	10	27	.37	2.36
7	8	30	.27	0.00	7	36	.23	0.00	4	19	.21	.01	12	32	.38	.41	9	27	.33	.07
8	12	27	.44	0.00	14	36	.47	.01	7	16	.44	.07	16	37	.43	.03	10	26	.38	.03

* SIGNIFICANT AT P LT 0.05

** SIGNIFICANT AT P LT 0.01

Table 30

NUMBER OF DEAD IMPLANTS PER TOTAL IMPLANTS - MALATHION

WEEK	CONTROL	74-07 1250 MG/KG	74-07 2500 MG/KG	74-07 5000 MG/KG	TEM .2 MG/KG
MULTIPLE TREATMENT					
1	13/ 319= .04	18/ 346= .05	8/ 298= .03	13/ 321= .04	62/ 316= .20**
2	8/ 303= .03	12/ 363= .03	11/ 267= .04	11/ 356= .03	77/ 293= .26**
3	9/ 245= .04	14/ 281= .05	14/ 305= .05	14/ 402= .03	87/ 348= .25**
4	2/ 309= .01	14/ 304= .05**	8/ 256= .03	7/ 300= .02	11/ 266= .04*
5	11/ 274= .04	4/ 236= .02*D	8/ 243= .03	8/ 306= .02	22/ 356= .06
6	21/ 302= .07	21/ 318= .07	20/ 226= .09	23/ 391= .06	17/ 273= .06
7	30/ 346= .09	10/ 337= .03	6/ 235= .03	18/ 352= .04	11/ 306= .04
8	19/ 292= .07	19/ 323= .06	7/ 187= .04	32/ 397= .08	11/ 322= .03

* SIGNIFICANT AT P LT 0.05

** SIGNIFICANT AT P LT 0.01

D DECREASED BELOW CONTROL

57

Table 31
 CHI-SQUARE TEST OF THE FERTILITY INDEX - PARATHION
 1 DEGREE OF FREEDOM

WEEK	VEHICLE CONTROL				74-01 62.5 MG/KG				74-01 125. MG/KG				74-01 250. MG/KG				74-01 500 MG/KG			
	N PWS	N MTD	FERT. INDEX	CHISQ	N PWS	N MTD	FERT. INDEX	CHISQ	N PWS	N MTD	FERT. INDEX	CHISQ	N PWS	N MTD	FERT. INDEX	CHISQ	N PWS	N MTD	FERT. INDEX	CHISQ
MULTIPLE TREATMENT																				
1	23	40	.57	0.00	25	40	.63	.05	27	40	.67	.48	22	40	.55	0.00	29	40	.72	1.37
2	27	39	.65	0.00	32	40	.80	.71	23	40	.57	.72	23	40	.57	.72	27	39	.69	.06
3	26	40	.65	0.00	23	40	.57	.21	20	40	.50	1.28	26	40	.65	.05	32	40	.80	1.57
4	27	40	.67	0.00	25	40	.63	.05	23	40	.57	.48	28	40	.70	0.00	27	40	.67	.06
5	29	40	.72	0.00	25	40	.63	.51	25	40	.63	.51	30	40	.75	0.00	30	40	.75	0.00
6	29	40	.72	0.00	28	40	.70	0.00	23	40	.57	1.37	32	40	.80	.29	27	38	.71	.01
7	29	40	.72	0.00	29	40	.72	.06	21	40	.52	2.61	24	40	.62	.06	27	36	.75	.00
8	32	40	.80	0.00	33	40	.82	0.00	29	40	.72	.28	30	40	.75	.07	26	36	.72	.28

85

Table 32

AVERAGE IMPLANTS PER PREGNANT FEMALE - PARATHION

WEEK	CONTROL	74-01 62.5 MG/KG	74-01 125. MG/KG	74-01 250. MG/KG	TFM .2 MG/KG
MULTIPLE TREATMENT					
1	265/ 23=11.52	266/ 25=10.64	314/ 27=11.63	224/ 22=10.18*	315/ 29=10.90
2	305/ 27=11.30	337/ 32=10.53	250/ 23=10.47	238/ 23=10.35	297/ 27=10.85
3	268/ 26=10.31	257/ 23=11.17	246/ 20=12.30 **1	277/ 26=10.65	348/ 32=10.87
4	298/ 27=10.67	277/ 25=11.08	253/ 23=11.00	311/ 28=11.11	266/ 27= 9.85
5	334/ 29=11.52	299/ 25=11.96	285/ 25=11.40	339/ 30=11.30	356/ 30=11.87
6	323/ 29=11.14	322/ 28=11.50	259/ 23=11.26	331/ 32=10.34	277/ 27=10.11
7	323/ 29=11.14	322/ 29=11.10	236/ 21=11.24	339/ 29=11.69	304/ 27=11.33
8	381/ 32=11.91	393/ 33=11.91	336/ 29=11.59	355/ 30=11.83	322/ 26=12.38

* SIGNIFICANT AT P LT 0.05

** SIGNIFICANT AT P LT 0.01

1 INCREASED ABOVE CONTROL

59

Table 33

AVERAGE DEAD IMPLANTS PER PREGNANT FEMALE - PARATHION

WEEK	CONTROL		74-01 62.5 MG/KG		74-01 125. MG/KG		74-01 250. MG/KG		TEM 2 MG/KG		
MULTIPLE TREATMENT											
09	1	0/ 23=	.35	18/ 25=	.72	11/ 27=	.41	8/ 22=	.36	67/ 29=	2.14**
	2	11/ 27=	.41	14/ 32=	.44	10/ 23=	.43	17/ 23=	.74	77/ 27=	2.85**
	3	16/ 26=	.62	7/ 23=	.30	10/ 20=	.50	8/ 26=	.31	87/ 32=	2.72**
	4	14/ 27=	.59	9/ 25=	.36	5/ 23=	.22 ^D	42/ 28=	1.50	11/ 27=	.41
	5	15/ 29=	.52	17/ 25=	.52	14/ 25=	.56	11/ 30=	.37	22/ 30=	.73
	6	9/ 29=	.31	10/ 28=	.36	9/ 23=	.39	14/ 32=	.44	17/ 27=	.63
	7	21/ 29=	.72	15/ 29=	.52	23/ 21=	1.10	22/ 29=	.76	11/ 27=	.41
	8	13/ 32=	.41	13/ 33=	.39	9/ 29=	.31	6/ 30=	.20	11/ 26=	.42

* SIGNIFICANT AT P LT 0.05

** SIGNIFICANT AT P LT 0.01

D DECREASED BELOW CONTROL

Table 34

CHI-SQUARE TEST OF THE DEATH INDEX - PARATHION
1 DEGREE OF FREEDOM

WEEK	VEHICLE CONTROL				74-01 62.5 MG/KG				74-01 125. MG/KG				74-01 250. MG/KG				TFM .2 MG/KG				
	N	N	DEATH	CHISQ	N	N	DEATH	CHISQ	N	N	DEATH	CHISQ	N	N	DEATH	CHISQ	N	N	DEATH	CHISQ	
	MDI	PRG	INDEX		MDI	PRG	INDEX		MDI	PRG	INDEX		MDI	PRG	INDEX		MDI	PRG	INDEX		
MULTIPLE TREATMENT																					
61	1	8	23	.35	0.00	6	25	.24	.25	9	27	.33	.04	5	22	.23	.32	25	29	.86	12.49**
	2	9	27	.33	0.00	11	32	.34	.04	8	23	.35	.04	7	23	.30	.01	26	27	.96	20.74**
	3	10	26	.30	0.00	7	23	.30	.00	8	20	.40	.04	7	26	.27	.35	25	32	.78	7.89**
	4	11	27	.41	0.00	8	25	.32	.13	4	23	.17	2.21	10	20	.36	.01	8	27	.39	.32
	5	11	29	.38	0.00	7	25	.28	.23	9	25	.36	.02	8	30	.27	.42	10	30	.39	.01
	6	8	29	.28	0.00	9	28	.32	.01	7	23	.30	.01	10	32	.31	.00	10	27	.37	.22
	7	12	29	.41	0.00	12	29	.41	.07	12	21	.57	.66	11	29	.30	0.00	9	27	.39	.12
	8	11	32	.34	0.00	11	33	.33	.03	9	29	.31	.00	5	30	.17	1.70	10	26	.38	.00

** SIGNIFICANT AT PLT 0.01

Table 35

NUMBER OF DEAD IMPLANTS PER TOTAL IMPLANTS - PARTITION

WEEK	CONTROL	74-01 62.5 MG/KG	74-01 125. MG/KG	74-01 250. MG/KG	1E4 12 MG/KG
MULTIPLE TREATMENT					
1	8/ 265 ^a .03	18/ 266 ^a .07	11/ 314 ^a .04	8/ 224 ^a .04	62/ 316 ^a .20**
2	11/ 305 ^a .04	14/ 337 ^a .04	10/ 250 ^a .04	17/ 239 ^a .07	77/ 293 ^a .26**
3	16/ 268 ^a .06	7/ 257 ^a .03	10/ 246 ^a .04	8/ 277 ^a .03	87/ 349 ^a .25**
4	16/ 288 ^a .06	9/ 277 ^a .03	5/ 253 ^a .02 ^{BD}	42/ 311 ^a .14	11/ 266 ^a .04
5	15/ 334 ^a .04	13/ 299 ^a .04	14/ 285 ^a .05	11/ 339 ^a .03	27/ 356 ^a .06
6	9/ 323 ^a .03	10/ 322 ^a .03	9/ 254 ^a .03	14/ 331 ^a .04	17/ 273 ^a .06
7	21/ 323 ^a .07	15/ 322 ^a .05	23/ 236 ^a .10	22/ 334 ^a .06	11/ 304 ^a .04
8	13/ 381 ^a .03	13/ 393 ^a .03	9/ 336 ^a .03	6/ 355 ^a .02 ^{BD}	11/ 322 ^a .03

* SIGNIFICANT AT P LT 0.05

** SIGNIFICANT AT P LT 0.01

D DECREASED BELOW CONTROL

Table 36

CHI-SQUARE TEST OF THE FERTILITY INDEX - PARATHION-METHYL
1 DEGREE OF FREEDOM

WEEK	VEHICLE CONTROL				74-05 20 MG/KG				74-05 40 MG/KG				74-05 80 MG/KG				TEM 1.2 MG/KG			
	N PRG	N MTD	FERT. INDEX	CHISQ	N PRG	N MTD	FERT. INDEX	CHISQ	N PRG	N MTD	FERT. INDEX	CHISQ	N PRG	N MTD	FERT. INDEX	CHISQ	N PRG	N MTD	FERT. INDEX	CHISQ
MULTIPLE TREATMENT																				
1	23	40	.57	0.00	29	40	.72	1.37	24	40	.60	0.00	29	40	.72	1.37	29	40	.72	1.37
2	27	39	.69	0.00	33	40	.82	1.25	23	40	.57	.72	29	40	.72	.01	27	39	.69	.06
3	26	40	.65	0.00	30	40	.75	.94	32	40	.80	1.57	28	40	.70	.06	32	40	.80	1.57
4	27	40	.67	0.00	38	40	.95	8.21**I	26	37	.70	.00	26	39	.67	.03	27	40	.67	.06
5	29	40	.72	0.00	32	40	.80	.28	24	40	.60	.69	30	40	.75	0.00	30	40	.75	0.00
6	29	40	.72	0.00	33	40	.82	.65	30	40	.75	0.00	24	40	.60	.89	27	38	.71	.01
7	29	40	.72	0.00	36	40	.90	2.95	32	40	.80	.28	33	40	.82	.65	27	36	.75	.00
8	32	40	.80	0.00	32	40	.80	.08	35	40	.88	.37	34	40	.85	.09	26	36	.72	.28

** SIGNIFICANT AT P LT 0.01
I INCREASED ABOVE CONTROL

Table 37
 AVERAGE IMPLANTS PER PREGNANT FEMALE - PARATHION-METHYL

WEEK	CONTROL	74-05 20 MG/KG	74-05 40 MG/KG	74-05 80 MG/KG	TEW .2 MG/KG
MULTIPLE TREATMENT					
1	265/ 23=11.52	331/ 29=10.30*	268/ 24=11.17	303/ 29=10.45*	316/ 29=10.90
2	305/ 27=11.30	347/ 33=10.52	241/ 23=10.48	323/ 29=11.14	293/ 27=10.85
3	268/ 26=10.31	306/ 30=10.20	331/ 32=10.34	297/ 28=10.61	348/ 32=10.87
4	268/ 27=10.67	417/ 38=10.97	275/ 26=10.58	281/ 26=10.81	266/ 27= 9.85
5	334/ 29=11.52	364/ 32=12.16	300/ 24=12.50	368/ 30=12.27	356/ 30=11.87
6	323/ 29=11.14	366/ 33=11.09	334/ 30=11.13	279/ 24=11.62	273/ 27=10.11
7	323/ 29=11.14	401/ 36=11.14	357/ 32=11.16	399/ 33=12.09	306/ 27=11.33
8	381/ 32=11.91	347/ 32=10.84*	407/ 35=11.63	402/ 34=11.82	322/ 26=12.38

* SIGNIFICANT AT P LT 0.05
 ** SIGNIFICANT AT P LT 0.01

Table 38

AVERAGE DEAD IMPLANTS PER PREGNANT FEMALE - PARATHION-METHYL.

WEEK	CONTROL	74-05 20 MG/KG	74-05 40 MG/KG	74-05 80 MG/KG	TEM .2 MG/KG
MULTIPLE TREATMENT					
1	8/ 23= .35	9/ 29= .31	25/ 24= 1.04 *	16/ 29= .55	62/ 29= 2.14**
2	11/ 27= .41	10/ 33= .30	8/ 23= .35	9/ 29= .31	77/ 27= 2.85**
3	16/ 26= .62	8/ 30= .27 *D	22/ 32= .69	34/ 28= 1.21	87/ 32= 2.72**
4	16/ 27= .59	18/ 38= .47	5/ 26= .19 *D	9/ 26= .35	11/ 27= .41
5	15/ 29= .52	23/ 32= .72	17/ 24= .71	12/ 30= .40	22/ 30= .73
6	9/ 24= .31	20/ 33= .61	10/ 30= .33	6/ 24= .25	17/ 27= .63
7	21/ 29= .72	7/ 36= .19 *D	11/ 32= .34	16/ 33= .48	11/ 27= .41
8	13/ 32= .41	11/ 32= .34	18/ 35= .51	12/ 34= .35	11/ 26= .42

* SIGNIFICANT AT P LT 0.05

** SIGNIFICANT AT P LT 0.01

D DECREASED BELOW CONTROL

Table 39
CHI-SQUARE TEST OF THE DEATH INDEX - PARATHION-METHYL

WEEK	VEHICLE CONTROL				74-05 20 MG/KG				74-05 40 MG/KG				74-05 80 MG/KG				75-05 1.7 MG/KG				
	N	N	DEATH	CHISQ	N	N	DEATH	CHISQ	N	N	DEATH	CHISQ	N	N	DEATH	CHISQ	N	N	DEATH	CHISQ	
	MDI	PRG	INDEX		MDI	PRG	INDEX		MDI	PRG	INDEX		MDI	PRG	INDEX		MDI	PRG	INDEX		
MULTIPLE TREATMENT																					
99	1	8	23	.35	0.00	8	29	.28	.07	12	24	.50	.58	9	29	.31	.00	25	29	.86	12.49**
	2	9	27	.33	0.00	9	33	.27	.05	6	23	.26	.06	8	29	.28	.03	26	27	.96	20.79**
	3	10	26	.38	0.00	8	30	.27	.43	14	32	.44	.02	9	28	.32	.04	25	32	.78	7.55**
	4	11	27	.41	0.00	14	36	.37	.00	5	26	.19	1.98	7	26	.27	.00	8	27	.30	.32
	5	11	29	.38	0.00	11	32	.34	.00	10	24	.42	.00	10	30	.33	.01	10	30	.33	.01
	6	8	29	.28	0.00	11	33	.33	.05	8	30	.27	.05	6	24	.25	.01	10	27	.37	.22
	7	12	29	.41	0.00	6	36	.17	3.74	8	32	.25	1.18	13	33	.39	.01	9	27	.33	.12
	8	11	32	.34	0.00	8	32	.25	.30	7	35	.20	1.10	8	34	.24	.49	10	26	.38	.00

** SIGNIFICANT AT P LT 0.01

Table 40

NUMBER OF DEAD IMPLANTS PER TOTAL IMPLANTS - PARATHION-METHYL

WEEK	CONTROL	74-05 20 MG/KG	74-05 40 MG/KG	74-05 80 MG/KG	TEH .2 MG/KG
MULTIPLE TREATMENT					
1	8/ 265= .03	9/ 301= .03	25/ 268= .09*	16/ 303= .05	62/ 316= .20**
2	11/ 305= .04	10/ 347= .03	8/ 241= .03	9/ 323= .03	77/ 293= .26**
3	16/ 268= .06	8/ 306= .03	22/ 331= .07	34/ 297= .11	87/ 343= .25**
4	16/ 288= .06	18/ 417= .04	5/ 275= .02 ^D	9/ 281= .03	11/ 266= .04
5	15/ 334= .04	23/ 389= .06	17/ 300= .06	12/ 368= .03	22/ 356= .06
6	9/ 323= .03	20/ 366= .05	10/ 334= .03	6/ 279= .02	17/ 273= .06
7	21/ 323= .07	7/ 401= .02 ^D	11/ 357= .03	16/ 399= .04	11/ 306= .04
8	13/ 381= .03	11/ 347= .03	18/ 407= .04	12/ 402= .03	11/ 322= .03

* SIGNIFICANT AT P LT 0.05

** SIGNIFICANT AT P LT 0.01

D DECREASED BELOW CONTROL

67

Table 41
 CHI-SQUARE TEST OF THE FERTILITY INDEX - QUINTOZOLE (PCNS)
 1 DEGREE OF FREEDOM

WEEK	VEHICLE CONTROL				74-04 1250 MG/KG				74-08 2500 MG/KG				74-06 5000 MG/KG				TEM 1.2 MG/KG				
	N PRG	N MTD	FERT. INDEX	CHISQ	N PRG	N MTD	FERT. INDEX	CHISQ	N PRG	N MTD	FERT. INDEX	CHISQ	N PRG	N MTD	FERT. INDEX	CHISQ	N PRG	N MTD	FERT. INDEX	CHISQ	
MULTIPLE TREATMENT																					
89	1	28	40	.76	0.00	29	40	.72	0.00	30	40	.75	.06	31	40	.77	.26	29	40	.72	0.00
	2	26	40	.65	0.00	21	34	.62	.00	22	40	.55	.47	26	40	.65	.05	27	39	.69	.03
	3	23	40	.57	0.00	25	38	.66	.27	26	38	.68	.50	29	40	.72	1.37	32	40	.80	3.72
	4	27	40	.67	0.00	33	39	.87	3.09	30	40	.75	.2*	28	40	.70	0.00	27	40	.67	.06
	5	24	40	.60	0.00	19	38	.50	.44	28	40	.70	.49	33	40	.82	3.91 ¹	30	40	.75	1.42
	6	24	38	.63	0.00	23	38	.61	0.00	26	40	.65	.00	36	40	.90	6.47 ¹	27	38	.71	.24
	7	24	38	.79	0.00	27	38	.71	.26	27	40	.67	.78	28	40	.70	.42	27	36	.75	.02
	8	27	38	.71	0.00	30	38	.79	.28	31	40	.77	.15	30	40	.75	.02	26	36	.72	.02

* SIGNIFICANT AT P LT 0.05
 1 INCREASED ABOVE CONTROL

Table -2

AVERAGE IMPLANTS PER PREGNANT FEMALE - QUINTOZENE (PCNB)

WEEK	CONTROL	74-08 1250 MG/KG	74-08 2500 MG/KG	74-08 5000 MG/KG	TEM .2 MG/KG
MULTIPLE TREATMENT					
1	319/ 28=11.39	349/ 29=12.03	338/ 30=11.27	361/ 31=11.65	316/ 29=10.90
2	303/ 26=11.65	222/ 21=10.57	235/ 22=10.68	259/ 26= 9.96*	293/ 27=10.85
3	245/ 23=10.65	296/ 25=11.84	326/ 26=12.54 **I	353/ 29=12.17*I	348/ 32=10.87
4	309/ 27=11.44	392/ 33=11.88	349/ 30=11.63	317/ 28=11.32	266/ 27= 9.85*
5	274/ 24=11.42	214/ 19=11.26	342/ 28=12.21	376/ 33=11.39	356/ 30=11.87
6	302/ 24=12.58	267/ 23=11.61	291/ 26=11.19 **	400/ 36=11.11*	273/ 27=10.11**
7	346/ 30=11.53	267/ 27=10.63*	292/ 27=10.81	302/ 28=10.79	306/ 27=11.33
8	292/ 27=10.81	354/ 30=11.80	336/ 31=10.84	333/ 30=11.10	322/ 26=12.38**I

* SIGNIFICANT AT P LT 0.05

** SIGNIFICANT AT P LT 0.01

I INCREASED ABOVE CONTROL

Table 43
 AVERAGE DEAD IMPLANTS PER PREGNANT FEMALE - QUINTOZENE (XCNB)

WEEK	CONTROL	74-08 1250 MG/KG	74-08 2500 MG/KG	74-08 5000 MG/KG	TEM .2 MG/KG
MULTIPLE TREATMENT					
1	13/ 28= .46	11/ 29= .38	12/ 30= .40	11/ 31= .35	62/ 29= 2.14 **
2	8/ 26= .31	12/ 21= .57	15/ 22= .68	6/ 26= .23	77/ 27= 2.85 **
3	9/ 23= .39	13/ 25= .52	7/ 26= .27	15/ 29= .52	87/ 32= 2.72 **
4	2/ 27= .07	15/ 33= .45**	7/ 30= .23	10/ 29= .36*	11/ 27= .41 *
5	11/ 24= .46	10/ 19= .53	21/ 28= .75	16/ 33= .48	22/ 30= .73
6	21/ 24= .88	6/ 23= .35 *D	9/ 26= .35 **D	16/ 36= .44 *D	17/ 27= .63
7	30/ 30= 1.00	11/ 27= .41	16/ 27= .59	20/ 28= .71	11/ 27= .41
8	19/ 27= .70	17/ 30= .57	10/ 31= .32	13/ 30= .43	11/ 26= .42

70

* SIGNIFICANT AT P LT 0.05
 ** SIGNIFICANT AT P LT 0.01
 D DECREASED BELOW CONTROL

Table 44

CHI-SQUARE TEST OF THE DEATH INDEX - QUINTOZENE (PCNB)
1 DEGREE OF FREEDOM

WEEK	VEHICLE CONTROL				74-08 1250 MG/KG				74-08 2500 MG/KG				74-08 5000 MG/KG				TFM 2 MG/KG			
	N	N	DEATH	CHISQ	N	N	DEATH	CHISQ	N	N	DEATH	CHISQ	N	N	DEATH	CHISQ	N	N	DEATH	CHISQ
	WDI	PRG	INDEX		WDI	PRG	INDEX		WDI	PRG	INDEX		WDI	PRG	INDEX		WDI	PRG	INDEX	
MULTIPLE TREATMENT																				
1	10	28	.34	0.00	7	29	.24	.44	10	30	.33	.01	9	31	.24	.07	25	29	.86	13.27 **
2	7	26	.27	0.00	10	21	.48	1.35	9	22	.41	.51	2	26	.08	2.15	26	27	.96	24.26 **
3	9	23	.39	0.00	11	25	.44	.00	6	26	.23	.82	11	29	.38	.04	25	32	.78	7.05 **
4	2	27	.07	0.00	12	33	.36	5.44 *	6	30	.20	.97	7	28	.25	1.96	8	27	.30	3.07
5	9	24	.38	0.00	9	19	.47	.12	13	28	.46	.14	13	33	.39	.02	10	30	.33	.00
6	15	24	.63	0.00	6	23	.26	4.91 *D	8	26	.31	3.86 *D	12	36	.33	3.84 *D	10	27	.37	2.36
7	8	30	.27	0.00	10	27	.37	.31	14	27	.52	2.81	10	28	.36	.21	9	27	.33	.07
8	12	27	.44	0.00	13	30	.43	.03	9	31	.29	.89	12	30	.40	.00	10	26	.38	.03

* SIGNIFICANT AT P LT 0.05

** SIGNIFICANT AT P LT 0.01

D DECREASED BELOW CONTROL

Table 45

NUMBER OF DEAD IMPLANTS PER TOTAL IMPLANTS - QUINTOZENE (PCNB)

WEEK	CONTROL	74-08 1250 MG/KG	74-08 2500 MG/KG	74-08 5000 MG/KG	TFM .2 MG/KG
MULTIPLE TREATMENT					
1	13/ 319# .04	11/ 349# .03	12/ 338# .04	11/ 361# .03	62/ 316# .20**
2	8/ 303# .03	12/ 222# .05	15/ 235# .06	6/ 259# .02	77/ 293# .26**
3	9/ 245# .04	13/ 296# .04	7/ 326# .02 =D	15/ 353# .04	87/ 348# .25**
4	2/ 309# .01	15/ 392# .04**	7/ 349# .02	10/ 317# .03*	11/ 266# .04*
5	11/ 274# .04	10/ 214# .05	21/ 342# .06	16/ 376# .04	22/ 356# .06
6	21/ 392# .07	8/ 267# .02=D	9/ 291# .03 =D	16/ 400# .04	17/ 273# .06
7	30/ 346# .09	11/ 287# .04	16/ 292# .05	20/ 302# .07	11/ 306# .04
8	19/ 292# .07	17/ 354# .05	10/ 336# .03	13/ 333# .04	11/ 322# .03

72

* SIGNIFICANT AT P LT 0.05

** SIGNIFICANT AT P LT 0.01

D DECREASED BELOW CONTROL

Table 46

CHI-SQUARE TEST OF THE FERTILITY INDEX - PRORATE
1 DEGREE OF FREEDOM

MULTIPLE TREATMENT

WEEK	VEHICLE CONTROL				74-04 5 MG/KG				74-04 10 MG/KG				74-04 20 MG/KG				TEM 12 MG/KG			
	N PRG	N MTU	FERT. INDEX	CHISQ	N PRG	N MTU	FERT. INDEX	CHISQ	N PRG	N MTU	FERT. INDEX	CHISQ	N PRG	N MTU	FERT. INDEX	CHISQ	N PRG	N MTU	FERT. INDEX	CHISQ
1	23	40	.57	0.00	24	40	.57	.05	21	40	.52	.05	24	40	.60	0.00	29	40	.72	1.37
2	27	39	.69	0.00	30	40	.75	.10	23	40	.57	.72	29	40	.72	.01	27	39	.69	.06
3	26	40	.65	0.00	26	40	.65	.05	24	40	.60	.05	31	40	.77	.98	32	40	.80	1.57
4	27	40	.67	0.00	26	40	.65	0.00	26	40	.65	0.00	32	40	.80	1.03	27	40	.67	.06
5	29	40	.72	0.00	25	40	.63	.91	27	40	.67	.06	31	40	.77	.07	30	40	.75	0.00
6	29	40	.72	0.00	25	40	.63	.91	26	40	.65	.23	33	40	.82	.65	27	38	.71	.01
7	29	40	.72	0.00	30	40	.75	0.00	29	40	.72	.06	35	40	.88	1.95	27	36	.75	.00
8	32	40	.80	0.00	26	40	.65	1.97	29	40	.72	.28	30	40	.75	.07	26	30	.72	.28

Table 47
 AVERAGE IMPLANTS PER PREGNANT FEMALE - PROSTATE

WEEK	CONTROL	74-04 5	MG/KG	74-04 10	MG/KG	74-04 20	MG/KG	1EM	.2 MG/KG
MULTIPLE TREATMENT									
1	265/ 23=11.52	241/ 23=10.48*		221/ 21=10.52 *		269/ 24=11.21		316/ 29=10.90	
2	303/ 27=11.30	326/ 30=10.87		255/ 23=11.09		307/ 29=10.59		293/ 27=10.85	
3	288/ 26=10.31	299/ 26=11.50*1		257/ 24=10.71		341/ 31=11.03		348/ 32=10.87	
4	288/ 27=10.67	241/ 26=11.19		277/ 26=10.65		353/ 32=11.03		266/ 27= 9.85	
5	334/ 29=11.52	278/ 29=11.12		312/ 27=11.56		369/ 31=11.90		356/ 30=11.87	
6	323/ 29=11.14	305/ 25=12.20*1		296/ 26=11.35		369/ 33=11.79		273/ 27=10.11	
7	323/ 29=11.14	336/ 30=11.20		317/ 29=10.93		407/ 35=11.63		306/ 27=11.33	
8	301/ 32=11.91	319/ 28=12.12		335/ 29=11.55		324/ 30=10.80*		322/ 28=12.38	

* SIGNIFICANT AT P LT 0.05
 ** SIGNIFICANT AT P LT 0.01
 1 INCREASED ABOVE CONTROL

74

Table 48
 AVERAGE DEAD IMPLANTS PER PREGNANT FEMALE - PHORATE

WEEK	CONTROL	74-04 5 MG/KG	74-04 10 MG/KG	74-04 20 MG/KG	74-04 20 MG/KG	74-04 20 MG/KG
MULTIPLE TREATMENT						
1	8/ 23= .35	9/ 23= .39	10/ 21= .48	3/ 24= .13	62/ 29= 2.14 **	
2	11/ 27= .41	16/ 30= .53	4/ 23= .17	17/ 29= .59	77/ 27= 2.85 **	
3	16/ 26= .62	13/ 26= .50	12/ 24= .50	15/ 31= .48	87/ 32= 2.72 **	
4	16/ 27= .59	15/ 26= .58	26/ 26= 1.00	12/ 32= .38	11/ 27= .41	
5	15/ 29= .52	13/ 25= .52	11/ 27= .41	13/ 31= .42	22/ 30= .73	
6	9/ 29= .31	13/ 25= .52	7/ 26= .27	7/ 33= .21	17/ 27= .63	
7	21/ 29= .72	13/ 30= .43	22/ 29= .76	7/ 35= .20* ^D	11/ 27= .41	
8	13/ 32= .41	10/ 26= .38	26/ 29= .90	11/ 30= .37	11/ 26= .42	

* SIGNIFICANT AT P LT 0.05
 ** SIGNIFICANT AT P LT 0.01
 D DECREASED BELOW CONTROL

75

Table 49
 CHI-SQUARE TEST OF THE DEATH INDEX - PHORATE
 1 DEGREE OF FREEDOM

WEEK	VEHICLE CONTROL				74-04 5 MG/KG				74-04 10 MG/KG				74-04 20 MG/KG				TEM .2 MG/KG			
	N	N	DEATH	CHISQ	N	N	DEATH	CHISQ	N	N	DEATH	CHISQ	N	N	DEATH	CHISQ	N	N	DEATH	CHISQ
	WDI	PRG	INDEX		WDI	PRG	INDEX		WDI	PRG	INDEX		WDI	PRG	INDEX		WDI	PRG	INDEX	
MULTIPLE TREATMENT																				
1	8	23	.35	0.00	8	23	.45	.10	7	21	.33	.05	2	24	.08	3.45	25	24	.86	12.49 **
2	9	27	.33	0.00	13	30	.43	.29	2	23	.09	3.07	13	29	.45	.37	26	27	.96	20.74 **
3	10	24	.38	0.00	7	26	.27	.35	12	24	.50	.29	10	31	.32	.04	25	32	.78	7.85 **
4	11	27	.41	0.00	11	26	.42	.03	15	26	.58	.92	11	32	.34	.05	8	27	.30	.32
5	11	29	.38	0.00	12	25	.48	.22	7	27	.26	.46	11	31	.35	.01	10	30	.33	.01
6	8	24	.28	0.00	10	25	.40	.46	7	26	.27	.06	7	33	.21	.06	10	27	.37	.22
7	12	29	.41	0.00	9	30	.30	.21	15	29	.52	.28	5	35	.14	4.66 ^D	9	27	.33	.12
8	11	32	.34	0.00	9	26	.35	.07	14	24	.48	.71	6	30	.27	.15	10	26	.38	.00

76

* SIGNIFICANT AT P LT 0.05
 ** SIGNIFICANT AT P LT 0.01
 D DECREASED BELOW CONTROL

Table 50
 NUMBER OF DEAD IMPLANTS PER TOTAL IMPLANTS - PHORATE

WEEK	CONTROL	74-04 5 MG/KG	74-04 10 MG/KG	74-04 20 MG/KG	1cM .2 MG/KG
MULTIPLE TREATMENT					
1	8/ 265= .03	9/ 241= .04	10/ 221= .05	3/ 264= .01 *D	62/ 316= .20 **
2	11/ 305= .04	16/ 326= .05	4/ 255= .02	17/ 307= .06	77/ 293= .26 **
3	16/ 268= .06	13/ 299= .04	12/ 257= .05	15/ 341= .04	87/ 348= .25 **
4	16/ 288= .06	15/ 291= .05	26/ 277= .09	12/ 353= .03	11/ 266= .04
5	15/ 334= .04	13/ 278= .05	11/ 312= .04	13/ 369= .04	22/ 356= .06
6	9/ 323= .03	13/ 305= .04	7/ 295= .02	7/ 369= .02	17/ 273= .06
7	21/ 323= .07	13/ 336= .04	22/ 317= .07	7/ 407= .02 *D	11/ 306= .04
8	13/ 381= .03	10/ 315= .03	26/ 335= .08	11/ 324= .03	11/ 322= .03

* SIGNIFICANT AT P LT 0.05
 ** SIGNIFICANT AT P LT 0.01
 D DECREASED BELOW CONTROL

77

Table 51

**DNA REPAIR SYNTHESIS ASSAY OF Monocrotophos
(dpm/ μ g DNA)**

	Monocrotophos (M)						4NQO (M)
	0	10^{-7}	10^{-6}	10^{-5}	10^{-4}	10^{-3}	10^{-2}
Sample 1	65*	46	31	54	70	64	1354
2	35	42	9*	46	46	86	1273
3	39	39	25	40	36	69	1308
4	34	34	25	40	40	51	975
5	30	26	46	53	64	64	972
6	20	--†	--†	--†	37	92	1135
Mean	32	38	32	47	50	71	1169
SD	7	7	10	7	14	15	168
SE	6	3	5	3	6	6	69

* Sample deleted from calculations because of low DNA value.

† Only five samples used.

Cell culture and experimental conditions

T-25 flask cultures of passage 24 WI-38 cells were initiated in medium containing 10% serum. The medium was replaced with medium containing 0.5% serum on day 5 following initiation and subsequently on days 11 and 15.

The assay was conducted on day 22.

Hydroxyurea (10^{-3} M) preincubation = 1 hour.

Compound exposure time = 3 hours.

3 H-TdR added with compound.

3 H-TdR incorporation = 1 μ Ci/ml (S.A. = 6.7 Ci/mole), 3 hours.

Postincorporation incubation = medium containing TdR, 3/4 hour.

Cells were removed with 1N NaOH, 1 minute, 70°C.

DNA was extracted by the PCA-hydrolysis procedure and measured following reaction with diphenylamine.

Negative control and compound solvent = 0.5% EtOH.

Table 52

**DNA REPAIR SYNTHESIS ASSAY OF Monocrotophos
WITH METABOLIC ACTIVATION
(dpm/ μ g DNA)**

	<u>Monocrotophos (M)</u>				<u>DMN (M)</u>
	<u>0</u>	<u>10^{-4}</u>	<u>10^{-3}</u>	<u>10^{-2}</u>	<u>5×10^{-2}</u>
Sample 1	55	67	43	87	206
2	54	66	48	84	220
3	52	43	52	82	223
Mean	54	59	48	84	216
SD	2	14	4	2	9
SE	1	8	2	1	5

Cell culture and experimental conditions

T-25 flask cultures of passage 24 WI-38 cells were initiated in medium containing 10% serum. The medium was replaced with medium containing 0.5% serum on day 4 following initiation and subsequently on day 10. The assay was conducted on day 16.

Hydroxyurea (10^{-2} M) preincubation = 1 hour.

Compound exposure time = 1 hour, with the 9,000 g fraction of a mouse liver homogenate.

3 H-TdR added with compound.

3 H-TdR incorporation = 1 μ Ci/ml (S.A. = 6.7 Ci/mole), 4 hours.

Postincorporation incubation = medium containing TdR, $\frac{1}{2}$ hour.

Cells were removed with 1N NaOH, 10 minutes, 22°C.

DNA was extracted by the PCA-hydrolysis procedure and measured following reaction with diphenylamine.

Negative control and compound solvent = 0.5% EtOH.

Table 53

DNA REPAIR SYNTHESIS TESTING
OF BROMACIL
(dpm/ μ g DNA)

Sample	Bromacil (M)						4NQO (M)
	<u>0*</u>	<u>10⁻⁷</u>	<u>10⁻⁶</u>	<u>10⁻⁵</u>	<u>10⁻⁴†</u>	<u>10⁻³†</u>	<u>10⁻⁵</u>
1	195	207	201	248	148	129	2670
2	153	212	215	178	144	137	2850
3	212	156	121	179	152	105	2688
4	230	187	218	231	204	107	2702
5	217	182	184	240	144	80	2438
6	298	251	220	178	200	74	2662
Mean	218	199	193	209	165	87	2668
SD	48	32	38	34	28	50	134
SE	19	13	15	14	12	20	55

* Negative control and compound solvent = 0.5% DMSO.

† Slight precipitate observed at 10⁻³ M and 10⁻⁴ M

Table 54

DNA REPAIR SYNTHESIS ASSAY OF BROMACIL
WITH METABOLIC ACTIVATION
(dpm/ μ g DNA)

Sample	Bromacil (M)						DMN (M)
	<u>0</u>	<u>10^{-7}</u>	<u>10^{-6}</u>	<u>10^{-5}</u>	<u>10^{-4}</u>	<u>10^{-3}</u>	<u>5×10^{-2}</u>
1	113	191	91	102	120	161	400
2	102	112	117	102	110	178	397
3	141	174	110	102	149	131	---*
4	158	189	82	91	135	185	529
5	218	152	104	126	167	141	645
6	136	170	96	190	165	133	448
Mean	145	165	100	119	141	155	484
SD	41	30	13	37	23	24	105
SE	17	12	5	15	9	10	50

* Sample lost.

Table 55

DNA REPAIR SYNTHESIS ASSAY
OF CACODYLIC ACID
(dpm/ μ g DNA)

	Cacodylic Acid (M)						4NQO (M)
	<u>0*</u>	<u>10⁻⁷</u>	<u>10⁻⁶</u>	<u>10⁻⁵</u>	<u>10⁻⁴</u>	<u>10⁻³†</u>	<u>10⁻²</u>
Sample 1	61	41	47	--‡	31	36	1891
2	31	48	27	--‡	27	19	1661
3	32	59	25	63	30	45	2418
4	18	32	38	68	29	19	2245
5	25	39	--‡	23	22	38	1430
6	35	22	--‡	29	28	36	2275
Mean	34	40	34	46	28	32	1990
SD	15	13	10	23	3	11	387
SE	6	5	5	11	1	4	158

* Negative control and compound solvent = 0.5% DMSO.

† Slight lowering of pH at 10⁻³ M.

‡ Sample lost.

Table 56

DNA REPAIR SYNTHESIS ASSAY OF CACODYLIC ACID
WITH METABOLIC ACTIVATION
(dpm/ μ g DNA)

	Cacodylic Acid (M)				DMN (M)
	0^*	10^{-5}	10^{-4}	10^{-3}	5×10^{-2}
Sample 1	44	25	21	29	381
2	30	33	22	25	384
3	23	39	28	21	339
Mean	33	32	24	25	368
SD	11	7	4	4	25
SE	6	4	2	2	15

* Negative control and compound solvent = 0.5% EtOH.

Table 57

**DNA REPAIR SYNTHESIS ASSAY OF CAPTAN
(dpm/ μ g DNA)**

	Captan (M)						4NQO(M)
	0*	10 ⁻⁸	10 ⁻⁷	10 ⁻⁶	10 ⁻⁵	10 ⁻⁴	10 ⁻⁵
Sample 1	37	43	41	74	81	8	924
2	64	58	53	60	81	6	947
3	76	50	68	52	57	7	1106
4	76	60	73	37	51	5	801
5	63	61	56	50	72	5	760
6	66	44	66	82	65	6	884
Mean	64	51	59	59	68	6	904
SD	14	8	12	16	12	1	122
SE	6	3	5	7	5	0.4	50

* Negative control and compound solvent = 0.5% DMSO.

Table 58

DNA REPAIR SYNTHESIS ASSAY OF CAPTAN
WITH METABOLIC ACTIVATION
(dpm/ μ g DNA)

	Captan (M)				DMN (M)
	0*	10^{-5}	10^{-4}	10^{-3}	5×10^{-2}
Sample 1	30	53	89	7	--†
2	40	50	73	5	323
3	--†	48	71	5	384
Mean	35	50	77	6	353
SD	7	2	9	1	43
SE	5	1	5	0.6	31

* Negative control and compound solvent = 0.5% DMSO.

† Sample lost.

Table 59

DNA REPAIR SYNTHESIS ASSAY OF CHLOROPYRIFOS

(dpm/ μ g/DNA)

	Chloropyrifos						4NQO (M)
	0*	10 ⁻⁷	10 ⁻⁶	10 ⁻⁵	10 ⁻⁴	10 ⁻³	10 ⁻⁵
Sample 1	115	280 [†]	282 [†]	-- [‡]	61	109	1337
2	143 [†]	129	93	110	67	-- [‡]	-- [‡]
3	96	102	84	110	60	-- [‡]	1721
4	72	95	64	-- [‡]	52	68	1220
5	97	89	99	98	64	37	1208
6	78	98	85	86	79	49	1209
Mean	92	103	85	101	64	66	1339
SD	17	15	13	11	9	31	220
SE	7	7	6	5	4	15	98

* Negative control and compound solvent = 0.5% DMSO.

† Sample deleted from calculations because of low DNA value.

‡ Sample lost.

Table 60

DNA REPAIR SYNTHESIS ASSAY OF CHLOROPYRIFOS
WITH METABOLIC ACTIVATION
(dpm/ μ g DNA)

	Chloropyrifos (M)				DMN (M)
	0*	10^{-5}	10^{-4}	10^{-3}	5×10^{-2}
Sample 1	72	79	70	52	355
2	75	65	75	71	349
3	55	67	63	79	384
Mean	67	70	69	67	363
SD	10	8	6	14	18
SE	6	5	4	8	11

* Negative control and compound solvent = 0.5% DMSO.

Table 61

**DEN REPAIR SYNTHESIS ASSAY
OF DINOSEB
(dpm/ μ g DNA)**

	Dinoseb (M)					4NQO (M)
	0^*	10^{-7}	10^{-6}	10^{-5}	10^{-4}^\dagger	10^{-6}
Sample 1	115	101	67	103	106	1337
2	143 [‡]	101	68	100	100	-- [§]
3	96	112	54	116	79	1721
4	72	61	58	62	66	1220
5	97	57	63	60	67	1208
6	78	58	73	60	76	1209
Mean	92	82	64	84	82	1339
SD	17	26	7	25	17	220
SE	7	11	3	10	7	98

* Negative control and compound solvent = 0.5% DMSO.

† Suggestion of precipitate at 10^{-4} M.

‡ Sample deleted from calculations because of low DNA value.

§ Sample lost.

Table 62

DNA REPAIR SYNTHESIS ASSAY OF DINOSEB
WITH METABOLIC ACTIVATION
(dpm/ μ g DNA)

	Dinoseb (M)				DMN (M)
	0*	10^{-5}	10^{-4}	10^{-3}	5×10^{-2}
Sample 1	72	93	76	71	355
2	75	81	80	64	349
3	55	51	58	89	384
Mean	67	75	71	74	363
SD	10	22	12	13	18
SE	6	12	7	8	11

* Negative control and compound solvent = 0.5% DMSO.

Table 63

DNA REPAIR SYNTHESIS ASSAY OF DSMA
(dpm/ μ g DNA)

	DSMA						4NQO (M)
	0*	10^{-7}	10^{-6}	10^{-5}	10^{-4}	10^{-3}	10^{-6}
Sample 1	67	51	79	86	100	62	902
2	58	54	67	64	155	49	1241
3	44	36	99	64	35	58	1380
4	55	36	400 [†]	45	44	59	990
5	79	62	74	53	55	68	1087
6	105	32	75	107	89	69	971
Mean	68	45	79	70	80	61	1095
SD	22	12	12	23	45	7	182
SE	9	5	5	9	18	3	74

* Negative control and compound solvent = H₂O.

† Sample deleted from calculations because of low DNA value.

Table 64

DNA REPAIR SYNTHESIS ASSAY OF DSMA
WITH METABOLIC ACTIVATION
(dpm/ μ g DNA)

	DSMA (M)				DMN (M)
	0*	10^{-5}	10^{-4}	10^{-3}	5×10^{-3}
Sample 1	44	28	25	38	381
2	30	36	29	36	384
3	23	21	32	28	339
Mean	33	29	29	34	368
SD	11	8	4	5	25
SE	6	4	2	3	15

* Negative control and compound solvent = 0.5% EtOH.

Table 65

DNA REPAIR SYNTHESIS ASSAY
OF FENTHION
(dpm/ μ g DNA)

	Fenthion (M)					4NQO (M)
	0 [*]	10 ⁻⁶	10 ⁻⁵	10 ⁻⁴	10 ⁻³ [†]	10 ⁻⁵
Sample 1	89	65	154	37	64	2983
2	43	105	337 [‡]	34	67	2272
3	107	83	85	40	36	2552
4	62	46	54	63	63	4059
5	61	34	102	31	44	1728
6	94	51	44	-- [§]	33	1893
Mean	76	64	88	41	51	2583
SD	24	26	44	13	15	857
SE	10	11	18	5	6	350

* Negative control and compound solvent = 0.5% EtOH.

† Precipitate observed at 10⁻³ M.

‡ Sample deleted from calculations because of low DNA value.

§ Sample lost.

Table 66

**DNA REPAIR SYNTHESIS ASSAY OF FENTHION
WITH METABOLIC ACTIVATION
(dpm/ μ g DNA)**

	Fenthion (M)				DMN (M)
	<u>0*</u>	<u>10⁻⁵</u>	<u>10⁻⁴</u>	<u>10⁻³</u>	<u>5 x 10⁻²</u>
Sample 1	55	54	60	51	206
2	54	50	46	64	220
3	52	42	64	63	223
Mean	54	48	57	60	216
SD	2	6	10	7	9
SE	1	4	6	4	5

* Negative control and compound solvent = 0.5% EtOH.

Table 67

DNA REPAIR SYNTHESIS ASSAY OF FOLPET
(dpm/ μ g DNA)

	Folpet (M)						4NQO(M)
	0*	10^{-8}	10^{-7}	10^{-6}	10^{-5}	10^{-4}	10^{-5}
Sample 1	37	43	63	45	82	29	924
2	64	58	62	54	58	25	947
3	76	91	108	52	92	31	1106
4	76	83	91	92	65	26	801
5	63	107	72	70	85	31	760
6	66	60	84	104	107	29	884
Mean	64	73	80	71	82	28	904
SD	14	24	18	25	18	2	122
SE	6	10	7	10	7	1	50

* Negative control and compound solvent = 0.5% DMSO.

Table 68
DNA REPAIR SYNTHESIS ASSAY OF FOLPET
WITH METABOLIC ACTIVATION
(dpm/ μ g DNA)

	<u>Folpet (M)</u>				<u>DMN (M)</u>
	<u>0*</u>	<u>10⁻⁵</u>	<u>10⁻⁴</u>	<u>10⁻³</u>	<u>5 x 10⁻²</u>
Sample 1	30	49	63	49	--†
2	40	54	82	40	323
3	--†	54	98	58	384
Mean	35	52	81	49	353
SD	7	3	18	9	43
SE	5	2	10	5	31

* Negative control and compound solvent = 0.5% DMSO

† Sample lost.

Table 69

DNA REPAIR SYNTHESIS ASSAY
OF AZINPHOS-METHYL
(dpm/ μ g DNA)

	Azinophos-methyl (M)						4NQO (M)
	0*	10^{-7}	10^{-6}	10^{-5}	10^{-4}	$10^{-3}\dagger$	10^{-5}
Sample 1	102	145	99	264 [‡]	51	55	804
2	99	115	103	99	105 [‡]	39	924
3	77	96	71	82	55	33	629
4	97	125	100	80	85	34	856
5	85	93	— [§]	79	57	97	761
6	111	77	72	56	68	35	897
Mean	95	108	89	79	63	49	822
SD	12	25	16	15	14	25	87
SE	5	10	7	7	6	10	36

* Negative control and compound solvent = 0.5% DMSO.

† Precipitate observed at 10^{-3} M.

‡ Sample deleted from calculations because of low DNA value.

§ Sample lost.

Table 70

DNA REPAIR SYNTHESIS ASSAY
OF AZINPHOS-METHYL WITH METABOLIC ACTIVATION
(dpm/ μ g DNA)

	DMN (M)				
	0*	10^{-5}	10^{-4}	10^{-3}	5×10^{-2}
Sample 1	30	70	75	42	--†
2	40	66	65	55	323
3	--†	78	41	54	384
Mean	35	71	60	50	353
SD	7	6	18	8	43
SE	5	3	10	4	31

* Negative control and compound solvent = 0.5% DMSO.

† Sample lost.

Table 71

DNA REPAIR SYNTHESIS ASSAY OF MALATHION
(dpm/ μ g DNA)

Sample	Malathion (M)						4NQO (M)
	0*	10^{-7}	10^{-6}	10^{-5}	10^{-4}	10^{-3}	10^{-5}
1	123	110	128	144	90	33	1943
2	125	111	124	78	126	34	1626
3	106	86	130	74	67	23	1538
4	100	133	116	119	113	44	1264
5	114	116	127	132	91	39	1737
6	138	110	143	127	156	40	1651
Mean	118	111	128	112	107	35	1626
SD	14	15	9	29	31	7	225
SE	6	6	4	12	13	3	92

* Negative control and compound solvent = 0.5% EtOH.

Table 72

**DNA REPAIR SYNTHESIS ASSAY OF MALATHION
WITH METABOLIC ACTIVATION
(dpm/ μ g DNA)**

	Malathion (M)				DMN (M)
	0*	10^{-5}	10^{-4}	10^{-3}	5×10^{-2}
Sample 1	55	48	46	38	206
2	54	49	52	48	220
3	52	62	55	37	223
Mean	54	53	51	41	216
SD	2	8	5	6	9
SE	1	5	3	4	5

* Negative control and compound solvent = 0.5% EtOH.

Table 73

DNA REPAIR SYNTHESIS ASSAY
OF METHOMYL
(dpm/ μ g DNA)

	Methomyl (M)						4NQO (M)
	0*	10^{-7}	10^{-6}	10^{-5}	10^{-4}	10^{-3}	10^{-2}
Sample 1	135	95	116	117	126	88	1442
2	133	-- [†]	133	108	116	69	1544
3	165	100	129	97	122	69	1518
4	72	98	97	115	109	69	1385
5	104	85	103	116	109	70	1423
6	103	95	93	139	130	61	-- [†]
Mean	118	94	112	115	118	71	1462
SD	32	6	17	14	9	9	67
SE	13	3	7	6	4	4	30

* Negative control and compound solvent = 0.5% DMSO.

[†] Sample lost.

Table 74

DNA REPAIR SYNTHESIS ASSAY OF METHOMYL
WITH METABOLIC ACTIVATION
(dpm/ μ g DNA)

	Methomyl (M)				DMN(M)
	0*	10 ⁻⁵	10 ⁻⁴	10 ⁻³	5 x 10 ⁻³
Sample 1	44	26	22	24	381
2	30	25	20	25	384
3	23	22	26	30	339
Mean	33	25	23	26	368
SD	11	2	3	3	25
SE	6	1	2	2	15

* Negative control and compound solvent = 0.5% EtOH.

Table 75

DNA REPAIR SYNTHESIS ASSAY
OF MONURON
(dpm/ μ g DNA)

	Monuron (M)						4NQO (M)
	0*	10^{-7}	10^{-6}	10^{-5}	10^{-4}	$10^{-3}\dagger$	10^{-6}
Sample 1	135	113	86	97	83	48	1442
2	133	-- [‡]	93	88	72	47	1544
3	165	129	89	77	81	46	1518
4	72	-- [‡]	83	75	82	49	1385
5	104	118	92	77	84	45	1423
6	103	131	110	109	85	38	-- [‡]
Mean	118	123	92	87	81	46	1462
SD	32	8	9	14	5	4	67
SE	13	4	4	6	2	1	30

* Negative control and compound solvent = 0.5% DMSO.

† Precipitate observed at 10^{-3} M.

‡ Sample lost.

Table 76

**DNA REPAIR SYNTHESIS ASSAY OF MONURON
WITH METABOLIC ACTIVATION
(dpm/ μ g DNA)**

	Monuron (M)				DMN (M)
	0*	10^{-6}	10^{-4}	10^{-3}	5×10^{-2}
Sample 1	30	78	81	74	--†
2	40	88	68	74	323
3	--†	92	63	88	384
Mean	36	86	71	79	353
SD	7	7	9	8	43
SE	5	4	5	5	31

* Negative control and compound solvent = 0.5% DMSO.

† Sample lost.

Table 77

DNA REPAIR SYNTHESIS ASSAY OF MSMA
(dpm/ μ g DNA)

	MSMA (M)						4NQO (M)
	0*	10 ⁻⁷	10 ⁻⁶	10 ⁻⁵	10 ⁻⁴	10 ⁻³	10 ⁻⁵
Sample 1	67	38	108	60	93	68	902
2	58	84	93	53	52	39	1241
3	44	114	68	61	66	61	1380
4	55	235 [†]	50	63	66	67	990
5	79	75	66	75	48	60	1087
6	105	63	-- [‡]	50	59	75	971
Mean	68	73	77	60	64	62	1095
SD	22	29	23	9	16	12	182
SE	9	13	9	4	7	5	74

* Negative control and compound solvent = H₂O.

† Sample deleted from calculations because of low DNA value.

‡ Sample lost.

Table 78

DNA REPAIR SYNTHESIS ASSAY OF MSMA
WITH METABOLIC ACTIVATION
(dpm/ μ g DNA)

	MSMA (M)				DMN (M)
	0*	10^{-5}	10^{-4}	10^{-3}	5×10^{-3}
Sample 1	44	27	21	32	381
2	30	25	35	25	384
3	23	21	34	24	339
Mean	33	24	30	27	368
SD	11	3	8	4	25
SE	6	2	5	3	15

* Negative control and compound solvent = 0.5% EtOH.

Table 79

DNA REPAIR SYNTHESIS ASSAY
OF PARATHION
(dpm/ μ g DNA)

Sample	Parathion (M)						4NQO (M)
	<u>0*</u>	<u>10⁻⁷</u>	<u>10⁻⁶</u>	<u>10⁻⁵</u>	<u>10⁻⁴</u>	<u>10⁻³</u>	<u>10⁻⁵</u>
1	123	127	151	221	102	90	1943
2	125	129	155	129	94	104	1626
3	106	135	135	157	83	93	1538
4	100	124	200	137	72	116	1264
5	114	137	160	155	93	102	1737
6	138	145	210	126	71	93	1651
Mean	118	133	169	154	86	100	1626
SD	14	7	29	35	13	10	225
SE	6	3	12	14	5	4	92

* Negative control and compound solvent = 0.5% EtOH.

Table 80

DNA REPAIR SYNTHESIS ASSAY OF PARATHION
WITH METABOLIC ACTIVATION
(dpm/ μ g DNA)

	Parathion (M)				DMN (M)
	0*	10^{-5}	10^{-4}	10^{-3}	5×10^{-2}
Sample 1	44	33	42	20	381
2	30	33	32	23	384
3	23	35	30	30	339
Mean	33	34	35	24	368
SD	11	1	6	5	25
SE	6	1	4	3	15

*

Negative control and compound solvent = 0.5% EtOH.

Table 81

DNA REPAIR SYNTHESIS ASSAY
OF PARATHION-METHYL
(dpm/ μ g DNA)

	Parathion-Methyl (M)						4NQO (M)
	0*	10^{-7}	10^{-6}	10^{-5}	10^{-4}	10^{-3} †	10^{-2}
Sample 1	36	36	33	34	44	40	411 [†]
2	34	38	32	55	44	23	781
3	86	49	28	31	53	28	782
4	94	56	27	52	41	29	858
5	53	49	35	43	40	29	1296
6	112 [‡]	46	41	85	44	27	1103
Mean	61	46	33	50	44	28	964
SD	28	8	5	20	5	6	227
SE	13	3	2	8	2	3	102

* Negative control and compound solvent = 0.5% EtOH.

† Precipitate observed at 10^{-3} M.

‡ Sample deleted from calculations because of low DNA value.

Table 82

DNA REPAIR SYNTHESIS ASSAY OF PARATHION-METHYL
WITH METABOLIC ACTIVATION
(dpm/ μ g DNA)

	Parathion-Methyl (M)				DMN (M)
	<u>0*</u>	<u>10⁻⁵</u>	<u>10⁻⁴</u>	<u>10⁻³</u>	<u>5 x 10⁻²</u>
Sample 1	55	44	65	51	206
2	54	54	52	48	220
3	52	57	37	45	223
Mean	54	52	51	48	216
SD	2	6	14	3	9
SE	1	4	8	2	5

* Negative control and compound solvent = 0.5% EtOH.

Table 83

DNA REPAIR SYNTHESIS ASSAY OF QUINTOZENE (PCNB)

(dpm/ μ g DNA)

	PCNB (M)						4NQO (M)
	0*	10^{-7}	10^{-8}	10^{-6}	10^{-4}	$10^{-3}\dagger$	10^{-5}
Sample 1	61	35	33	21	27	23	1891
2	31	43	22	21	18	37	1681
3	32	44	31	23	18	19	2418
4	18	30	30	16	18	21	2245
5	25	38	27	32	22	27	1430
6	35	20	20	39	20	27	2275
Mean	34	35	27	25	20	26	1990
SD	15	9	5	8	4	6	387
SE	6	4	2	3	2	3	158

* Negative control and compound solvent = 0.5% DMSO.

† Precipitate observed at 10^{-3} M.

Table 84

**DNA REPAIR SYNTHESIS ASSAY OF QUINTOZENE (PCNB)
WITH METABOLIC ACTIVATION
(dpm/ μ g DNA)**

	PCNB (M)				DMN (M)
	0 *	10^{-5}	10^{-4}	10^{-3}	5×10^{-3}
Sample 1	72	75	95	106	355
2	75	79	71	72	349
3	55	84	73	49	364
Mean	67	79	80	76	363
SD	10	5	13	29	18
SE	6	3	8	17	11

* Negative control and compound solvent = 0.5% DMSO.

Table 85

DNA REPAIR SYNTHESIS ASSAY OF PHORATE

(dpm/ μ g DNA)

	Phorate (M)						4NQO (M)
	0 [*]	10 ⁻⁷	10 ⁻⁶	10 ⁻⁵	10 ⁻⁴	10 ⁻³ †	10 ⁻²
Sample 1	36	58	38	25	27	56	411 ‡
2	34	45	44	17	50	45	781
3	86	31	107	22	42	55	782
4	84	26	-- §	43	43	50	858
5	53	40	44	52	37	61	1295
6	112 ‡	56	26	77	39	59	1103
Mean	61	43	52	39	40	55	964
SD	28	13	32	23	8	6	227
SE	13	5	14	9	3	2	102

* Negative control and compound solvent = 0.5% EtOH.

† Precipitate observed at 10⁻³ M.

‡ Sample deleted from calculations because of low DNA value.

§ Sample lost.

Table 86

DNA REPAIR SYNTHESIS ASSAY OF PHORATE
WITH METABOLIC ACTIVATION
(dpm/ μ g DNA)

	Phorate (M)				DMN (M)
	0*	10^{-5}	10^{-4}	10^{-3}	5×10^{-2}
Sample 1	55	45	43	37	206
2	54	59	41	35	220
3	52	63	39	38	223
Mean	54	55	41	37	216
SD	2	10	2	1.4	9
SE	1	6	1	0.8	5

* Negative control and compound solvent = 0.5% EtOH.

Table 87

DNA REPAIR SYNTHESIS ASSAY
OF SIMAZINE
(dpm/ μ g DNA)

Sample	Simazine (M)						4NQO (M)
	<u>0*</u>	<u>10⁻⁷</u>	<u>10⁻⁶</u>	<u>10⁻⁵</u>	<u>10⁻⁴</u>	<u>10⁻³</u>	<u>10⁻⁵</u>
1	195	165	151	195	177	369	2670
2	153	91	171	190	193	208	2850
3	212	131	152	312	253	233	2688
4	230	138	146	290	281	165	2702
5	217	152	179	237	166	205	2435
6	298	113	213	305	161	--- [†]	2662
Mean	218	132	169	255	205	236	2668
SD	48	27	25	55	50	78	134
SE	19	11	10	22	20	32	55

* Media control and compound solvent = 0.5% DMSO.

† Sample lost.

Table 38

DNA REPAIR SYNTHESIS ASSAY OF SIMAZINE
WITH METABOLIC ACTIVATION
(dpm/ μ g DNA)

	Simazine (M)				DMN (M)
	0*	10^{-8}	10^{-6}	10^{-3}	5×10^{-2}
Sample 1	72	57	64	59	355
2	75	58	60	58	349
3	55	64	61	76	384
Mean	67	60	62	64	333
SD	10	4	2	10	18
SE	6	2	1	6	11

* Negative control and compound solvent = 0.5% DMSO.

Table 89

**DNA REPAIR SYNTHESIS ASSAY
OF TRIFLURALIN
(dpm/ μ g DNA)**

	Trifluralin (M)						4NQO (M)
	0*	10^{-7}	10^{-6}	10^{-5}	10^{-4}	10^{-3}	10^{-2}
Sample 1	56	42	26	53	51	71	639
2	68	29	--†	152‡	97	123	1125
3	51	78	48	68	156‡	112	570
4	45	51	76	89	78	83	894
5	50	47	79	97	56	80	663
6	29	41	59	57	43	53	986
Mean	50	48	58	73	62	87	812
SD	13	17	22	19	28	26	222
SE	5	7	10	9	12	11	91

* Negative control and compound solvent = 0.5% EtOH.

† Sample lost.

‡ Sample deleted from calculations because of low DNA value.

Table 90

DNA REPAIR SYNTHESIS ASSAY OF TRIFLURALIN
WITH METABOLIC ACTIVATION
(dpm/ μ g DNA)

	Trifluralin (M)				DMN (M)
	0*	10^{-5}	10^{-4}	10^{-3}	5×10^{-2}
Sample 1	72	67	79	51	355
2	75	58	64	61	349
3	55	74	79	--†	384
Mean	67	66	74	56	363
SD	10	8	9	7	18
SE	6	5	5	5	11

* Negative control and compound solvent = 0.5% DMSO.

† Sample lost.

Table 91

IN VITRO ASSAYS WITH SALMONELLA TYPHIMURIUM

<u>Compound</u>	<u>Metabolic Activation</u>	<u>µg Compound Added/Plate</u>	<u>Average Number of Histidine-Positive Revertants/Plate</u>			
			<u>TA100</u>	<u>TA1535</u>	<u>TA1537</u>	<u>TA1538</u>
Negative control	-		95	13	8	7
	+		113	13	10	10
Positive control, 4-o-tolylazo-1-toluidine	-	25				6
	+	25				183
Monocrotophos	-	1	87	9	10	11
	-	5	101	22	9	10
	-	10	93	14	9	13
	-	50	107	23	7	10
	-	100	97	13	11	9
	-	500	95	17	10	7
	-	1000	126	14	10	6
	+	1	101	16	14	12
	+	5	89	20	13	12
	+	10	75	16	12	11
	+	50	79	14	10	10
	+	100	71	16	15	11
	+	500	78	19	9	15
	+	1000	113	18	10	10

118

Monocrotophos

Table 91 (continued)

Compound	Metabolic Activation	µg Compound Added/Plate	Average Number of Histidine-Positive Revertants/Plate			
			TA100	TA1535	TA1537	TA1538
Negative control	-		145	22	25	16
	+		154	25	24	30
Positive controls						
β-Propiolactone	-	50 µl		756		
AF-2	-	0.05	372			
2-Anthramine	-	50				63
	+	50				338
Bromacil	-	1	120	23	24	26
	-	5	129	17	13	14
	-	10	123	31	22	13
	-	50	117	29	16	15
	-	100	136	40	18	19
	-	500	140	30	15	15
	-	1000	101	14	6	11
	+	1	118	35	21	16
	+	5	131	28	16	20
	+	10	157	33	21	19
	+	50	136	26	14	13
	+	100	138	30	12	15
	+	500	145	21	20	20
	+	1000	162	8	5	16

119

Bromacil

Table 91 (continued)

Compound	Metabolic Activation	µg Compound Added/Plate	Average Number of Histidine-Positive Revertants/Plate			
			TA100	TA1535	TA1537	TA1538
Negative control	-		56	15	12	7
	+		72	14	9	15
Positive control, 4-o-tolylazo-o-toluidine	-	25				10
	+	25				150
Cacodylic Acid	-	1	48	17	15	5
	-	5	42	15	15	11
	-	10	42	12	15	8
	-	50	39	18	10	8
	-	100	43	22	11	9
	-	500	44	15	11	9
	-	1000	44	16	8	8
	+	1	69	17	14	18
	+	5	53	15	15	8
	+	10	64	16	12	18
	+	50	50	15	11	12
	+	100	64	18	15	19
	+	500	54	21	14	15
	+	1000	59	14	13	13

150

Cacodylic Acid

Table 91 (continued)

<u>Compound</u>	<u>Metabolic Activation</u>	<u>ug Compound Added/Plate</u>	<u>Average Number of Histidine-Positive Revertants/Plate</u>			
			<u>TA100</u>	<u>TA1535</u>	<u>TA1537</u>	<u>TA1538</u>
Negative control	-		72	18	7	8
	+		98	14	3	25
Positive control, 4-o-tolylazo-o-toluidine	-	2		3		
	+	2		100		
Captan	-	1	211	29	2	7
	-	5	532	80	5	14
	-	10	822	76	0	16
	-	15	820	104	0	26
	-	25	720	80	0	5
	-	50	Killing	Killing	0	22
	+	1	141	20	2	19
	+	5	210	60	2	22
	+	10	285	113	2	26
	+	15	340	55	0	21
	+	25	330	71	0	46
+	50	704	143	1	44	

Table 91 (continued)

Compound	Metabolic Activation	µg Compound Added/Plate	Average Number of Histidine-Positive Revertants/Plate			
			TA100	TA1535	TA1537	TA1538
Negative control	-		92	18	12	16
	+		80	14	16	16
Positive control, 4-o-tolylazo-o-toluidine	-					15
	+					168
Chloropyrifos	-	1	66	20	12	22
	-	5	92	22	15	28
	-	10	65	14	21	24
	-	50	88	26	15	25
	-	100	67	20	17	22
	-	500	87	17	18	17
	-	1000	79	13	20	22
	+	1	67	11	15	14
	+	5	71	9	15	16
	+	10	87	11	15	20
	+	50	77	16	18	13
	+	100	77	11	14	16
	+	500	72	13	14	30
	+	1000	77	14	11	22

122

Chloropyrifos

Table 91 (continued)

<u>Compound</u>	<u>Metabolic Activation</u>	<u>µg Compound Added/Plate</u>	<u>Average Number of Histidine-Positive Revertants/Plate</u>			
			<u>TA100</u>	<u>TA1535</u>	<u>TA1537</u>	<u>TA1538</u>
Negative control	-		97	15	12	21
	+		80	15	16	17
Positive control, 4-o-tolylazo-o-toluidine	-	25				15
	+	25				168
Dinoseb	-	1	69	12	21	15
	-	5	59	12	16	14
	-	10	57	17	18	20
	-	50	67	17	17	17
	-	100	82	12	16	15
	-	500	91	7	17	19
	-	1000	Killing	Killing	Killing	Killing
	+	1	79	13	19	15
	+	5	82	14	18	14
	+	10	94	12	17	15
	+	50	83	14	15	17
	+	100	87	13	17	14
	+	500	104	12	10	7
	+	1000	Killing	Killing	Killing	Killing

Table 91 (continued)

Compound	Metabolic Activation	µg Compound Added/Plate	Average Number of Histidine-Positive Revertants/Plate			
			TA100	TA1535	TA1537	TA1538
Negative control	-		56	15	12	7
	+		72	14	9	15
Positive control, 4-o-tolylazo-o-toluidine	-	25				10
	+	25				250
DSMA	-	1	50	12	7	4
	-	5	56	10	10	5
	-	10	51	20	10	3
	-	50	66	15	11	8
	-	100	71	16	7	8
	-	500	53	13	12	7
	-	1000	43	12	7	9
	+	1	54	22	8	8
	+	5	85	7	8	15
	+	10	50	16	-	3
	+	50	55	17	11	5
	+	100	50	13	9	7
	+	500	60	15	5	10
	+	1000	53	14	2	8

Table 91 (continued)

Compound	Metabolic Activation	µg Compound Added/Plate	Average Number of Histidine-Positive Revertants/Plate			
			TA100	TA1535	TA1537	TA1538
Negative control	-		101	26	6	13
	+		102	26	3	24
Positive control, 4-o-tolylazo-o-toluidine	-	25				13
	+	25				78
Azinphos-methyl	-	1	66	39	4	10
	-	5	74	22	3	11
	-	10	74	23	2	9
	-	50	73	30	3	11
	-	100	75	49	4	10
	-	500	107	30	2	10
	-	1000	104	31	3	13
	+	1	76	23	1	20
	+	5	69	23	2	25
	+	10	68	24	3	28
	+	50	81	22	3	21
	+	100	65	24	2	19
	+	500	84	30	0	24
	+	1000	119	24	0	23

125

Azinphos-methyl

Table 91 (continued)

Compound	Metabolic Activation	µg Compound Added/Plate	Average Number of Histidine-Positive Revertants/Plate			
			TA100	TA1535	TA1537	TA1538
Negative control	-		94	36	10	12
	+		80	20	9	12
Positive control, 4-o-tolylazo-o-toluidine	-	25				15
	+	25				168
Fenthion	-	1	64	31	8	13
	-	5	97	34	11	17
	-	10	105	32	15	12
	-	50	112	36	15	14
	-	100	100	38	16	14
	-	500	107	42	10	14
	-	1000	90	32	6	12
	+	1	114	15	10	12
	+	5	97	17	12	10
	+	10	81	9	9	17
	+	50	90	16	12	21
	+	100	98	14	7	13
	+	500	86	22	8	10
	+	1000	89	20	10	15

Table 91 (continued)

Compound	Metabolic Activation	µg Compound Added/Plate	Average Number of Histidine-Positive Revertants/Plate			
			TA100	TA1535	TA1537	TA1538
Negative control	-		72	19	3	8
	+		93	15	7	20
Positive control, 4-o-tolylazo-o-toluidine	-	25				6
	+	25				183
Folpet	-	1	127	20	7	5
	-	5	150	35	0	8
	-	10	244	39	1	11
	-	25	300	48	0	14
	-	50	550	111	2	7
	-	100	286	110	0	2
	-	500	Killing	Killing	0	Killing
	-	1000	Killing	Killing	0	Killing
	+	1	112	20	2	30
	+	5	173	51	1	26
	+	10	241	79	5	35
	+	25	420	70	6	36
	+	50	720	218	10	45
	+	100	532	216	3	48
+	500	Killing	Killing	0	Killing	
+	1000	Killing	Killing	0	Killing	

127

Folpet

Table 91 (continued)

Compound	Metabolic Activation	µg Compound Added/Plate	Average Number of Histidine-Positive Revertants/Plate			
			TA100	TA1535	TA1537	TA1538
Negative control	-		89	10	8	7
	+		92	10	10	7
Positive control, 4-c-tolylazo-o-toluidine	-	25				6
	+	25				183
Malathion	-	1	54	8	11	3
	-	5	48	7	12	3
	-	10	85	7	10	5
	-	50	99	8	6	7
	-	100	81	7	7	5
	-	500	82	12	5	7
	-	1000	61	10	7	4
	+	1	65	5	6	9
	+	5	61	6	8	5
	+	10	99	9	7	4
	+	50	92	8	7	6
	+	100	75	9	6	5
	+	500	90	8	12	4
	+	1000	66	7	10	4

128

Malathion

Table 91 (continued)

Compound	Metabolic Activation	µg Compound Added/Plate	Average Number of Histidine-Positive Revertants/Plate			
			TA100	TA1535	TA1537	TA1538
Negative control	-		128	18	33	17
	+		149	14	20	22
Positive control, 4-o-tolylazo-o-toluidine	-	25				17
	+	25				206
Methomyl	-	1	123	19	28	14
	-	5	112	20	35	10
	-	10	98	16	28	18
	-	50	109	18	27	14
	-	100	110	21	34	24
	-	500	119	17	23	26
	-	1000	105	13	24	21
	+	1	145	12	18	15
	+	5	115	10	18	15
	+	10	126	12	21	13
	+	50	129	10	19	20
	+	100	132	13	20	19
	+	500	133	10	20	18
	+	1000	122	14	24	14

Table 91 (continued)

<u>Compound</u>	<u>Metabolic Activation</u>	<u>µg Compound Added/Plate</u>	<u>Average Number of Histidine-Positive Revertants/Plate</u>			
			<u>TA100</u>	<u>TA1535</u>	<u>TA1537</u>	<u>TA1538</u>
Negative control	-		128	17	17	17
	+		149	12	15	22
Positive control, 4-o-tolylazo-o-toluidine	-	25				6
	+	25				177
Monuron	-	1	126	15	15	19
	-	5	98	24	12	18
	-	10	108	17	12	22
	-	50	122	19	11	21
	-	100	114	19	15	29
	-	500	122	20	18	29
	-	1000	125	22	14	19
	+	1	125	10	15	21
	+	5	142	13	15	15
	+	10	119	17	13	18
	+	50	116	15	19	21
	+	100	108	15	16	19
	+	500	104	15	15	12
	+	1000	123	11	15	17

130

Monuron

Table 91 (continued)

<u>Compound</u>	<u>Metabolic Activation</u>	<u>µg Compound Added/Plate</u>	<u>Average Number of Histidine-Positive Revertants/Plate</u>			
			<u>TA100</u>	<u>TA1535</u>	<u>TA1537</u>	<u>TA1538</u>
Negative control	-		56	15	12	7
	+		72	14	9	15
Positive control, 4-o-tolylazo-o-toluidine	-	25				10
	+	25				250
MSMA	-	1	79	15	7	6
	-	5	69	15	13	4
	-	10	62	14	11	6
	-	50	52	17	11	6
	-	100	41	15	12	7
	-	500	53	17	8	5
	-	1000	48	13	12	5
	+	1	79	11	11	10
	+	5	64	15	8	10
	+	10	65	7	10	9
	+	50	67	7	14	7
	+	100	53	12	8	8
	+	500	66	14	7	8
	+	1000	68	10	10	10

Table 91 (continued)

Compound	Metabolic Activation	µg Compound Added/Plate	Average Number of Histidine-Positive Revertants/Plate			
			TA100	TA1535	TA1537	TA1538
Negative control	-		95	19	6	6
	+		114	21	13	7
Positive control, 4-o-tolylazo-o-toluidine	-	25				6
	+	25				177
Parathion	-	1	94	12	7	8
	-	5	138	12	7	7
	-	10	85	13	4	7
	-	50	98	13	4	8
	-	100	87	15	4	6
	-	500	110	13	4	8
	-	1000	107	14	3	13
	+	1	56	11	12	15
	+	5	75	16	7	15
	+	10	69	14	5	19
	+	50	76	14	6	8
	+	100	88	17	7	5
	+	500	105	15	8	9
	+	1000	103	12	6	12

Table 91 (continued)

<u>Compound</u>	<u>Metabolic Activation</u>	<u>µg Compound Added/Plate</u>	<u>Average Number of Histidine-Positive Revertants/Plate</u>			
			<u>TA100</u>	<u>TA1535</u>	<u>TA1537</u>	<u>TA1538</u>
Negative control	-		96	15	8	8
	+		118	17	11	11
Positive control, 4-o-tolylazo-o-toluidine	-	25				6
	+	25				177
Phorate	-	1	70	17	8	16
	-	5	65	15	8	11
	-	10	85	17	7	11
	-	50	65	17	5	8
	-	100	72	11	7	9
	-	500	70	14	6	6
	-	1000	58	14	7	9
	+	1	101	14	11	15
	+	5	103	11	8	10
	+	10	79	13	10	12
	+	50	89	15	9	11
	+	100	79	15	6	7
	+	500	59	16	6	11
	+	1000	70	19	8	5

Phorate

Table 91 (continued)

Compound	Metabolic Activation	µg Compound Added/Plate	Average Number of Histidine-Positive Revertants/Plate			
			TA100	TA1535	TA1537	TA1538
Negative control	-		98	10	8	25
	+		106	7	8	26
Positive control; 4-o-tolylazo-o-toluidine	-	25				22
	+	25				266
Simazine	-	1	83	7	15	22
	-	5	72	5	10	20
	-	10	73	7	20	20
	-	50	87	7	10	22
	-	100	85	8	12	17
	-	500	71	3	7	25
	-	1000	69	4	11	22
	+	1	84	9	11	22
	+	5	90	4	11	22
	+	10	82	8	16	15
	+	50	83	9	9	14
	+	100	87	8	11	15
	+	500	89	4	16	15
	+	1000	120	2	10	18

136

Simazine

Table 91 (concluded)

Compound	Metabolic Activation	µg Compound Added/Plate	Average Number of Histidine-Positive Revertants/Plate			
			TA100	TA1535	TA1537	TA1538
Negative control	-		96	15	17	14
	+		80	15	20	8
Positive control, 4-o-tolylazo-o-toluidine	-	25				15
	+	25				168
Trifluralin	-	1	73	11	18	15
	-	5	81	12	24	14
	-	10	74	16	29	14
	-	50	93	19	23	16
	-	100	86	18	25	15
	-	500	76	18	22	11
	-	1000	95	13	18	15
	+	1	72	10	13	9
	+	5	80	13	16	9
	+	10	78	14	18	14
	+	50	90	15	14	13
	+	100	81	8	16	15
	+	500	79	12	13	11
	+	1000	81	12	15	10

Table 92

RESULTS OF ASSAYS WITH ESCHERICHIA COLI WP2

<u>Compound</u>	<u>Metabolic Activation</u>	<u>µg of Compound Added per Plate</u>	<u>Average Number of Tryptophan- Positive Revertants per Plate</u>	
Negative control	-		68	
	+		73	
Positive control, AF-2	-	0.05	204	
	+	0.05	220	
Monocrotophos	-	1	89	
	-	10	83	
	-	50	76	
	-	100	77	
	-	500	61	
	-	1000	70	
	+	1	90	
	+	10	88	
	+	50	76	
	+	100	73	
	+	500	75	
	+	1000	95	
	Bromacil	-	1	65
		-	10	74
-		50	71	
-		100	70	
-		500	70	
-		1000	67	
+		1	71	
+		10	73	
+		50	66	
+		100	70	
+		500	70	
+		1000	71	

Table 92 (continued)

<u>Compound</u>	<u>Metabolic Activation</u>	<u>µg of Compound Added per Plate</u>	<u>Average Number of Tryptophan- Positive Revertants per Plate</u>	
Cacodylic Acid	-	1	111	
	-	10	102	
	-	50	89	
	-	100	82	
	-	500	91	
	-	1000	85	
	+	1	95	
	+	10	76	
	+	50	79	
	+	100	89	
	+	500	81	
	+	1000	85	
	Captan	-	1	124
		-	5	381
-		10	733	
-		15	1358	
-		25	1755	
-		50	2600	
+		1	89	
+		5	182	
+		10	423	
+		15	699	
+		25	955	
+		50	1712	

Table 92 (continued)

<u>Compound</u>	<u>Metabolic Activation</u>	<u>µg of Compound Added per Plate</u>	<u>Average Number of Tryptophan-Positive Revertants per Plate</u>
Chloropyrifos	-	1	57
	-	10	57
	-	50	49
	-	100	71
	-	500	52
	-	1000	42
	+	1	60
	+	10	73
	+	50	53
	+	100	49
	+	500	61
	+	1000	49
Dinoseb	-	1	64
	-	10	73
	-	50	58
	-	100	55
	-	500	44
	-	1000	Toxic
	+	1	73
	+	10	69
	+	50	63
	+	100	65
	+	500	49
	+	1000	Toxic

140

Table 92 (continued)

<u>Compound</u>	<u>Metabolic Activation</u>	<u>µg of Compound Added per Plate</u>	<u>Average Number of Tryptophan- Positive Revertants per Plate</u>	
DSMA	-	1	86	
	-	10	81	
	-	50	67	
	-	100	68	
	-	500	68	
	-	1000	71	
	+	1	69	
	+	10	62	
	+	50	67	
	+	100	73	
	+	500	81	
	+	1000	79	
	Fenthion	-	1	59
		-	10	63
-		50	62	
-		100	56	
-		500	70	
-		1000	71	
+		1	64	
+		10	50	
+		50	67	
+		100	64	
+		500	64	
+		1000	81	

Table 92 (continued)

<u>Compound</u>	<u>Metabolic Activation</u>	<u>ug of Compound Added per Plate</u>	<u>Average Number of Tryptophan- Positive Revertants per Plate</u>	
Folpet	-	1	65	
	-	5	162	
	-	10	170	
	-	25	424	
	-	50	720	
	-	100	1260	
	+	1	74	
	+	5	167	
	+	10	202	
	+	25	900	
	+	50	1680	
	+	100	1880	
	Azinphos-methyl	-	1	92
		-	10	87
-		50	83	
-		100	89	
-		500	68	
-		1000	88	
+		1	83	
+		10	74	
+		50	87	
+		100	86	
+		500	73	
+		1000	79	

Table 92 (continued)

<u>Compound</u>	<u>Metabolic Activation</u>	<u>µg of Compound Added per Plate</u>	<u>Average Number of Tryptophan-Positive Revertants per Plate</u>	
Malathion	-	1	62	
	-	10	54	
	-	50	60	
	-	100	60	
	-	500	54	
	-	1000	48	
	+	1	58	
	+	10	50	
	+	50	55	
	+	100	59	
	+	500	75	
	+	1000	64	
	Methomyl	-	1	61
		-	10	76
-		50	83	
-		100	57	
-		500	68	
-		1000	71	
+		1	70	
+		10	81	
+		50	78	
+		100	83	
+		500	63	
+		1000	74	

Table 92 (continued)

<u>Compound</u>	<u>Metabolic Activation</u>	<u>µg of Compound Added per Plate</u>	<u>Average Number of Tryptophan-Positive Revertants per Plate</u>	
Monuron	-	1	72	
	-	10	68	
	-	50	57	
	-	100	63	
	-	500	65	
	-	1000	63	
	+	1	60	
	+	10	59	
	+	50	47	
	+	100	71	
	+	500	50	
	+	1000	61	
	MSMA	-	1	55
		-	10	64
-		50	57	
-		100	76	
-		500	60	
-		1000	63	
+		1	55	
+		10	71	
+		50	73	
+		100	71	
+		500	61	
+		1000	72	

Table 92 (continued)

<u>Compound</u>	<u>Metabolic Activation</u>	<u>µg of Compound Added per Plate</u>	<u>Average Number of Tryptophan- Positive Revertants per Plate</u>	
Parathion	-	1	71	
	-	10	64	
	-	50	66	
	-	100	70	
	-	500	64	
	-	1000	64	
	+	1	69	
	+	10	53	
	+	50	76	
	+	100	57	
	+	500	72	
	+	1000	66	
	Parathion-methyl	-	1	53
		-	10	56
-		50	60	
-		100	68	
-		500	63	
-		1000	52	
+		1	64	
+		10	83	
+		50	60	
+		100	65	
+		500	53	
+		1000	71	

Table 92 (continued)

<u>Compound</u>	<u>Metabolic Activation</u>	<u>µg of Compound Added per Plate</u>	<u>Average Number of Tryptophan-Positive Revertants per Plate</u>	
Quintozene (PCNB)	-	1	54	
	-	10	59	
	-	50	70	
	-	100	60	
	-	500	57	
	-	1000	62	
	+	1	78	
	+	10	67	
	+	50	54	
	+	100	57	
	+	500	59	
	+	1000	62	
	Phorate	-	1	63
		-	10	64
-		50	65	
-		100	49	
-		500	71	
-		1000	60	
+		1	78	
+		10	86	
+		50	83	
+		100	73	
+		500	90	
+		1000	70	

Table 92 (concluded)

<u>Compound</u>	<u>Metabolic Activation</u>	<u>µg of Compound Added per Plate</u>	<u>Average Number of Tryptophan-Positive Revertants per Plate</u>	
Simazine	-	1	55	
	-	10	51	
	-	50	73	
	-	100	54	
	-	500	54	
	-	1000	53	
	+	1	64	
	+	10	66	
	+	50	72	
	+	100	56	
	+	500	71	
	+	1000	83	
	Trifluralin	-	1	75
		-	10	73
-		50	81	
-		100	86	
-		500	69	
-		1000	60	
+		1	58	
+		10	63	
+		50	63	
+		100	65	
+		500	70	
+		1000	70	

Table 93

MICROBIAL INHIBITION IN ESCHERICHIA COLI AND BACILLUS SUBTILIS

Compound	mg of Compound Added to Disc	Diameter of Zone of Inhibition (mm)			
		<u>E. coli</u>		<u>B. subtilis</u>	
		<u>W3110</u>	<u>p3478</u>	<u>H17</u>	<u>m45</u>
Positive control, 1-phenyl-3, dimethyltriazeno	1.0	37	52	40	61
Negative control, chloramphenicol	0.03	34.5	34	32	31
Monocrotophos	1	6	6	6	6
Bromacil	1.0	6.5	6.5	6.5	6.5
Cacodylic acid	1	6	6	6	6
Captan	0.1	6.5	11	9	19
Chloropyrifos	2.5	6	10	6	11
Dinoseb	1	10	17	8.5	11
DMSA	1	6	6	6	6
Fenthion	1	6	6	6	6
Folpet	0.1	6.5	10	6.5	7.5
Azinphos-methyl	1	6	6	6	6

Table 93 (concluded)

Compound	mg of Compound Added to Disc	Diameter of Zone of Inhibition (mm)			
		E. coli		B. subtilis	
		W3110	p3478	H17	m45
Malathion	1	6	6	6	6
Methomyl	1	6	6	6	6
Monuron	1	6	6	6	6
MSMA	1	6	6	6	6
Parathion	1	6	6	6	6
Parathion-methyl	1	6	6	6	6
Quintozene (PCNB)	1	6	6	6	6
Phorate	1	6	6	6	6
Simazine	1	6	6	6	6
Trifluralin	1	6	6	6	6

Table 94

IN VITRO ASSAYS WITH SACCHAROMYCES CEREVISIAE D3 - MONOCROTOPHOS

<u>Compound</u>	<u>Metabolic Activation</u>	<u>Percent Concentration (w/v or v/v)</u>	<u>Survivors</u>		<u>Mitotic Recombinants</u>	
			<u>Cells/ml (x 10⁻⁷)</u>	<u>Percent of Control</u>	<u>per ml (x 10⁻³)</u>	<u>per 10⁵ Survivors</u>
<u>EXPERIMENT 1</u>						
Negative control	-		5.7	100	4.5	7.9
	+		5.8	100	4.5	7.8
Positive control 1,2,3,4-Diepoxybutane	-	0.1	5.8	102	1,650	2,845
	+	0.1	4.6	79	1,435	3,120
Monocrotophos	-	5	5.7	100	44	77.2
	+	5	4.7	81	30	63.8
<u>EXPERIMENT 2</u>						
Negative control	-		9.1	100	8	8.8
	+		8.6	100	9.5	11.0
Monocrotophos	-	5	6.3	69	26	41.2
	+	5	4.8	56	40	83.3

Table 95

IN VITRO ASSAYS WITH SACCHAROMYCES CEREVISIAE D3 - BROMACIL

Compound	Metabolic Activation	Percent Concentration (w/v or v/v)	Survivors		Mitotic Recombinants	
			Cells/ml ($\times 10^7$)	Percent of Control	per ml ($\times 10^3$)	per 10^8 Survivors
EXPERIMENT 1						
Negative control	-		4.8	100	3	6.3
	+		4.7	100	3	6.4
Positive control						
1,2,3,4-Diepoxybutane	-	0.04	3.4	71	745	2191
	+	0.04	3.5	74	683	1951
Bromacil						
	-	0.005	5.0	104	3	6.0
	-	0.01	4.5	94	2	4.4
	-	0.05	5.0	104	3	6.0
	-	0.10	4.4	92	1	2.3
	-	0.50	2.0	42	1	10.0
	+	0.005	4.4	94	5	11.4
	+	0.01	3.9	83	3	7.7
	+	0.05	4.3	91	3	7.0
	+	0.10	3.8	81	1	2.6
	+	0.50	2.4	51	3	12.5
EXPERIMENT 2						
Negative control	-		4.5	100	5	11.1
	+		4.2	100	3	7.1
Positive control						
1,2,3,4-Diepoxybutane	-	0.04	4.5	100	870	1933
	+	0.04	4.2	100	653	1555
Bromacil						
	-	0.05	5.7	128	7	12.3
	-	0.10	5.4	120	5	9.3
	-	0.25	5.5	122	5	9.1
	-	0.50	4.9	108	1	2.0
	+	0.05	4.9	117	4	8.2
	+	0.10	4.7	112	5	10.6
	+	0.25	4.8	114	6	12.5
	+	0.50	4.9	117	1	2.0

Table 96

IN VITRO ASSAYS WITH SACCHAROMYCES CEREVISIAE D3 - CACODYLIC ACID

<u>Compound</u>	<u>Metabolic Activation</u>	<u>Percent Concentration (w/v or v/v)</u>	<u>Survivors</u>		<u>Mitotic Recombinants</u>	
			<u>Cells/ml (x 10⁻⁷)</u>	<u>Percent of Control</u>	<u>per ml (x 10⁻³)</u>	<u>per 10⁵ Survivors</u>
<u>EXPERIMENT 1</u>						
Negative control	-		7.4	100	7.5	10.1
	+		7.7	100	5	6.5
Cacodylic acid	-	5	5.6	76	20	35.7
	+	5	6.3	82	11	17.5
<u>EXPERIMENT 2</u>						
Negative control	-		7.1	100	3.5	4.9
	+		6.5	100	3	4.6
Cacodylic acid	-	5	15.5	217	1,187	766
	+	5	17.5	270	1,159	662

Table 97

IN VITRO ASSAYS WITH SACCHAROMYCES CEREVISIAE D3 - CAPTAN

<u>Compound</u>	<u>Metabolic Activation</u>	<u>Percent Concentration (w/v or v/v)</u>	<u>Survivors</u>		<u>Mitotic Recombinants</u>	
			<u>Cells/ml (x 10⁻⁷)</u>	<u>Percent of Control</u>	<u>per ml (x 10⁻³)</u>	<u>per 10⁶ Survivors</u>
<u>EXPERIMENT 1</u>						
Negative control	-		7.1	100	3.5	4.9
	+		6.5	100	3.0	4.6
Captan	-	0.003	6.0	84	205	342
	+	0.003	9.1	140	145	159
<u>EXPERIMENT 2</u>						
Negative control	-		7.5	100	1.5	2.0
	+		6.0	100	4	6.7
Captan	-	0.003	.77	10	37	481
	+	0.003	5.1	85	58	114

Table 98

IN VITRO ASSAYS WITH SACCHAROMYCES CEREVISIAE D3 - CHLOROPYRIFOS

<u>Compound</u>	<u>Metabolic Activation</u>	<u>Percent Concentration (w/v or v/v)</u>	<u>Survivors</u>		<u>Mitotic Recombinants</u>	
			<u>Cells/ml (x 10⁻⁷)</u>	<u>Percent of Control</u>	<u>per ml (x 10⁻³)</u>	<u>per 10⁵ Survivors</u>
<u>EXPERIMENT 1</u>						
Negative control	-		7.5	100	1.5	2.0
	+		6.0	100	4	6.7
Chloropyrifos	-	5	7.7	103	7	9.1
	+	5	8.0	133	10	12.5
<u>EXPERIMENT 2</u>						
Negative control	-		6.3	100	1.5	2.4
	+		7.4	100	3.5	4.7
Positive control, 1,2,3,4-diepoxybutane	-	0.1	3.8	60	1,045	2,750
	+	0.1	5.2	70	903	1,737
Chloropyrifos	-	5	7.9	125	3	3.8
	+	5	7.4	100	10	13.5

Table 99

IN VITRO ASSAYS WITH SACCHAROMYCES CEREVISIAE D3 - DINOSEB

<u>Compound</u>	<u>Metabolic Activation</u>	<u>Percent Concentration (w/v or v/v)</u>	<u>Survivors</u>		<u>Mitotic Recombinants</u>	
			<u>Cells/ml (x 10⁻⁷)</u>	<u>Percent of Control</u>	<u>per ml (x 10⁻³)</u>	<u>per 10⁵ Survivors</u>
<u>EXPERIMENT 1</u>						
Negative control	-		6.3	100	1.5	2.4
	+		7.4	100	3.5	4.7
Positive control, 1,2,3,4-diepoxybutane	-	0.1	3.8	60	1,045	2,750
	+	0.1	5.2	70	903	1,737
Dinoseb	-	0.2	6.2	98	9	14.5
	+	0.2	4.0	54	1	2.5
	-	0.3	.3	5	5	167
	+	0.3	2.4	32	5	20.8
<u>EXPERIMENT 2</u>						
Negative control	-		5.5	100	2.5	4.5
	+		5.2	100	2.0	3.8
Dinoseb	-	0.1	4.4	80	3	6.8
	+	0.1	4.0	77	4	10.0
	-	0.2	4.1	75	8	19.5
	+	0.2	4.3	83	8	18.6

Table 100

IN VITRO ASSAYS WITH SACCHAROMYCES CEREVISIAE D3 - DSMA

<u>Compound</u>	<u>Metabolic Activation</u>	<u>Percent Concentration (w/v or v/v)</u>	<u>Survivors</u>		<u>Mitotic Recombinants</u>	
			<u>Cells/ml (x 10⁻⁷)</u>	<u>Percent of Control</u>	<u>per ml (x 10⁻³)</u>	<u>per 10⁵ Survivors</u>
<u>EXPERIMENT 1</u>						
Negative control	-		7.4	100	7.5	10.1
	+		7.7	100	5	6.5
DSMA	-	4.5	1.1	15	0	
	+	4.5	5.2	68	0	
<u>EXPERIMENT 2</u>						
Negative control	-		7.1	100	3.5	4.9
	+		6.5	100	3	4.6
DSMA	-	5	4.7	66	3	6.4
	+	5	3.4	52	7	20.6

Table 101

IN VITRO ASSAYS WITH SACCHAROMYCES CEREVISIAE D3 - FENTHION

<u>Compound</u>	<u>Metabolic Activation</u>	<u>Percent Concentration (w/v or v/v)</u>	<u>Survivors</u>		<u>Mitotic Recombinants</u>	
			<u>Cells/ml (x 10⁻⁷)</u>	<u>Percent of Control</u>	<u>per ml (x 10⁻³)</u>	<u>per 10⁵ Survivors</u>
<u>EXPERIMENT 1</u>						
Negative control	-		6.3	100	1.5	2.4
	+		7.4	100	3.5	4.7
Positive control, 1,2,3,4-diepoxybutane	-	0.1	3.8	60	1,045	2,750
	+	0.1	5.2	70	903	1,737
Fenthion	-	5	6.6	105	9	13.6
	+	5	7.5	101	5	6.7
<u>EXPERIMENT 2</u>						
Negative control	-		7.5	100	1.5	2.0
	+		6.0	100	4	6.7
Fenthion	-	5	7.8	104	4	5.1
	+	5	7.1	118	6	8.5

Table 102

IN VITRO ASSAYS WITH SACCHAROMYCES CEREVISIAE D3 - FOLPET

<u>Compound</u>	<u>Metabolic Activation</u>	<u>Percent Concentration (w/v or v/v)</u>	<u>Survivors</u>		<u>Mitotic Recombinants</u>	
			<u>Cells/ml (x 10⁻⁷)</u>	<u>Percent of Control</u>	<u>per ml (x 10⁻³)</u>	<u>per 10⁵ Survivors</u>
<u>EXPERIMENT 1</u>						
Negative control	-		7.5	100	1.5	2.0
	+		6.0	100	4	6.7
Folpet	-	0.003	4.0	53	119	298
	+	0.003	3.8	63	65	171
<u>EXPERIMENT 2</u>						
Negative control	-		6.3	100	1.5	2.3
	+		7.4	100	3.5	4.7
Folpet	-	0.003	9.5	151	89	94
	+	0.003	9.1	123	82	90

Table 103

IN VITRO ASSAYS WITH SACCHAROMYCES CEREVISIAE D3 - AZINPHOS-METHYL

<u>Compound</u>	<u>Metabolic Activation</u>	<u>Percent Concentration (w/v or v/v)</u>	<u>Survivors</u>		<u>Mitotic Recombinants</u>	
			<u>Cells/ml (x 10⁻⁷)</u>	<u>Percent of Control</u>	<u>per ml (x 10⁻³)</u>	<u>per 10⁵ Survivors</u>
<u>EXPERIMENT 1</u>						
Negative control	-		5.7	100	4.5	7.9
	+		5.8	100	4.5	7.8
Positive control 1,2,3,4-Diepoxybutane	-	0.1	5.8	102	1,650	2,845
	+	0.1	4.6	79	1,435	3,120
Azinphos-methyl	-	4.5	5.3	93	15	28.3
	+	4.5	5.8	100	15	25.9
<u>EXPERIMENT 2</u>						
Negative control	-		9.1	100	8	8.8
	+		8.6	100	9.5	11.0
Azinphos-methyl	-	5	5.7	63	68	119.3
	+	5	6.2	72	80	129

Table 104

IN VITRO ASSAYS WITH SACCHAROMYCES CEREVISIAE D3 - MALATHION

<u>Compound</u>	<u>Metabolic Activation</u>	<u>Percent Concentration (w/v or v/v)</u>	<u>Survivors</u>		<u>Mitotic Recombinants'</u>	
			<u>Cells/ml (x 10⁻⁷)</u>	<u>Percent of Control</u>	<u>per ml (x 10⁻³)</u>	<u>per 10⁵ Survivors</u>
<u>EXPERIMENT 1</u>						
Negative control	-		5.7	100	4.5	7.9
	+		5.8	100	4.5	7.8
Positive control, 1,2,3,4-diepoxybutane	-	0.1	5.8	102	1,650	2,845
	+	0.1	4.6	79	1,435	3,120
Malathion	-	5	7.8	137	11	14.1
	+	5	6.3	109	7	11.1
<u>EXPERIMENT 2</u>						
Negative control	-		9.1	100	8	8.8
	+		8.6	100	9.5	11.0
Malathion	-	5	8.1	89	13	16.0
	+	5	7.6	88	8	10.5

Table 105

IN VITRO ASSAYS WITH SACCHAROMYCES CEREVISIAE D3 - METHOMYL

<u>Compound</u>	<u>Metabolic Activation</u>	<u>Percent Concentration (w/v or v/v)</u>	<u>Survivors</u>		<u>Mitotic Recombinants</u>		
			<u>Cells/ml (x 10⁻⁷)</u>	<u>Percent of Control</u>	<u>per ml (x 10⁻³)</u>	<u>per 10⁵ Survivors</u>	
<u>EXPERIMENT 1</u>							
Negative control	-		6.6	100	4.5	6.8	
	+		5.8	100	2.5	4.6	
Positive control, 1,2,3,4-diepoxybutane	-	0.1	1.8	27	266	1,478	
	+	0.1	1.5	29	184	1,227	
Methomyl	-	2.0	5.0	76	4	8.0	
	+	2.0	2.7	50	0		
	-	3.0	3.7	56	8	21.6	
	+	3.0	4.1	76	6	14.6	
	-						
	+						
<u>EXPERIMENT 2</u>							
Negative control	-		5.5	100	2.5	4.5	
	+		5.2	100	2.0	3.8	
Methomyl	-	3	4.7	85	13	31.9	
	+	3	4.4	85	10	22.7	

Table 106

IN VITRO ASSAYS WITH SACCHAROMYCES CEREVISIAE D3 - MONURON

<u>Compound</u>	<u>Metabolic Activation</u>	<u>Percent Concentration (w/v or v/v)</u>	<u>Survivors</u>		<u>Mitotic Recombinants</u>	
			<u>Cells/ml (x 10⁻⁷)</u>	<u>Percent of Control</u>	<u>per ml (x 10⁻³)</u>	<u>per 10⁵ Survivors</u>
<u>EXPERIMENT 1</u>						
Negative control	-		6.6	100	4.5	6.8
	+		5.4	100	2.5	4.6
Positive control, 1,2,3,4-diepoxybutane	-	0.1	1.8	27	266	1,478
	+	0.1	1.5	29	184	1,227
Monuron	-	5	3.5	53	3	8.6
	+	5	3.8	70	1	2.6
<u>EXPERIMENT 2</u>						
Negative control	-		5.5	100	2.5	4.5
	+		5.2	100	2.0	3.8
Monuron	-	5	6.9	125	2	2.9
	+	5	6.2	119	9	14.5

Table 107

IN VITRO ASSAYS WITH SACCHAROMYCES CEREVISIAE D3 - MSMA

<u>Compound</u>	<u>Metabolic Activation</u>	<u>Percent Concentration (w/v or v/v)</u>	<u>Survivors</u>		<u>Mitotic Recombinants</u>	
			<u>Cells/ml (x 10⁻⁷)</u>	<u>Percent of Control</u>	<u>per ml (x 10⁻³)</u>	<u>per 10⁵ Survivors</u>
<u>EXPERIMENT 1</u>						
Negative control	-		7.4	100	7.5	10.1
	+		7.7	100	5	6.5
MSMA	-	5	4.3	58	1	2.3
	+	5	5.4	70	3	5.6
<u>EXPERIMENT 2</u>						
Negative control	-		7.1	100	3.5	4.9
	+		6.5	100	3	4.6
MSMA	-	5	4.9	69	10.2	20.8
	+	5	5.8	89	10.4	17.9

Table 108

IN VITRO ASSAYS WITH SACCHAROMYCES CEREVISIAE D3 - PARATHION

<u>Compound</u>	<u>Metabolic Activation</u>	<u>Percent Concentration (w/v or v/v)</u>	<u>Survivors</u>		<u>Mitotic Recombinants</u>	
			<u>Cells/ml (x 10⁻⁷)</u>	<u>Percent of Control</u>	<u>per ml (x 10⁻³)</u>	<u>per 10⁵ Survivors</u>
<u>EXPERIMENT 1</u>						
Negative control	-		5.7	100	4.5	7.9
	+		5.8	100	4.5	7.8
Positive control, 1,2,3,4-diepoxybutane	-	0.1	5.8	102	1,650	2,845
	+	0.1	4.6	79	1,435	3,120
Parathion	-	5	6.5	114	3	4.6
	+	5	5.8	100	5	8.6
<u>EXPERIMENT 2</u>						
Negative control	-		9.1	100	8	8.8
	+		8.6	100	9.5	11.0
Parathion	-	5	8.8	96	4	4.5
	+	5	8.2	95	5	6.1

Table 109

IN VITRO ASSAYS WITH SACCHAROMYCES CEREVISIAE D3 - PARATHION-METHYL

<u>Compound</u>	<u>Metabolic Activation</u>	<u>Percent Concentration (w/v or v/v)</u>	<u>Survivors</u>		<u>Mitotic Recombinants</u>	
			<u>Cells/ml (x 10⁻⁷)</u>	<u>Percent of Control</u>	<u>per ml (x 10⁻³)</u>	<u>per 10⁵ Survivors</u>
<u>EXPERIMENT 1</u>						
Negative control	-		5.7	100	4.5	7.9
	+		5.8	100	4.5	7.8
Positive control, 1,2,3,4-diepoxybutane	-	0.1	5.8	102	1,650	2,845
	+	0.1	4.6	79	1,435	3,120
Parathion-methyl	-	5	7.7	135	16	20.8
	+	5	5.4	93	15	27.8
<u>EXPERIMENT 2</u>						
Negative control	-		9.1	100	8	8.8
	+		8.6	100	9.5	11.0
Parathion-methyl	-	5	7.4	81	19	25.7
	+	5	7.2	84	25	34.7

Table 110

IN VITRO ASSAYS WITH SACCHAROMYCES CEREVISIAE D3 - QUINTOZENE (PCNB)

Compound	Metabolic Activation	Percent Concentration (w/v or v/v)	Survivors		Mitotic Recombinants	
			Cells/ml (x 10 ⁻⁷)	Percent of Control	per ml (x 10 ⁻³)	per 10 ⁵ Survivors
<u>EXPERIMENT 1</u>						
Negative control	-		5.7	100	4.5	7.9
	+		5.8	100	4.5	7.8
Positive control, 1,2,3,4-diepoxybutane	-	0.1	5.8	102	1,650	2,845
	+	0.1	4.6	79	1,435	3,120
Quintozene (PCNB)	-	2	3.7	65	3	8.1
	+	2	4.2	72	4	9.5
<u>EXPERIMENT 2</u>						
Negative control	-		9.1	100	8	8.8
	+		8.6	100	9.5	11.0
Quintozene (PCNB)	-	1	5.8	64	4	6.9
	+	1	7.0	81	7	10.0
	-	2	6.8	75	3	4.4
	+	2	7.5	87	10	13.3

Table 111

IN VITRO ASSAYS WITH SACCHAROMYCES CEREVISIAE D3 - PHORATE

<u>Compound</u>	<u>Metabolic Activation</u>	<u>Percent Concentration (w/v or v/v)</u>	<u>Survivors</u>		<u>Mitotic Recombinants</u>	
			<u>Cells/ml (x 10⁻⁷)</u>	<u>Percent of Control</u>	<u>per ml (x 10⁻³)</u>	<u>per 10⁵ Survivors</u>
<u>EXPERIMENT 1</u>						
Negative control	-		9.1	100	8	8.8
	+		8.6	100	9.5	11.0
Phorate	-	5	8.7	96	9	10.3
	+	5	7.5	87	3	4.0
<u>EXPERIMENT 2</u>						
Negative control	-		5.7	100	4.5	7.9
	+		5.8	100	4.5	7.8
Phorate	-	5	7.5	132	4	5.3
	+	5	7.2	124	7	9.7

Table 112

IN VITRO ASSAYS WITH SACCHAROMYCES CEREVISIAE D3 - SIMAZINE

<u>Compound</u>	<u>Metabolic Activation</u>	<u>Percent Concentration (w/v or v/v)</u>	<u>Survivors</u>		<u>Mitotic Recombinants</u>	
			<u>Cells/ml (x 10⁻⁷)</u>	<u>Percent of Control</u>	<u>per ml (x 10⁻³)</u>	<u>per 10⁵ Survivors</u>
<u>EXPERIMENT 1</u>						
Negative control	-		6.6	100	4.5	6.8
	+		5.4	100	2.5	4.6
Positive control, 1,2,3,4-diepoxybutane	-	0.1	1.8	27	266	1,478
	+	0.1	1.5	29	184	1,227
Simazine	-	5	3.8	58	3	7.9
	+	5	2.0	37	1	5.0
<u>EXPERIMENT 2</u>						
Negative control	-		5.5	100	2.5	4.5
	+		5.2	100	2	3.8
Simazine	-	5	7.0	127	7	10.0
	+	5	7.0	135	4	5.7

Table 113

IN VITRO ASSAYS WITH SACCHAROMYCES CEREVISIAE D3 - TRIFLURALIN

<u>Compound</u>	<u>Metabolic Activation</u>	<u>Percent Concentration (w/v or v/v)</u>	<u>Survivors</u>		<u>Mitotic Recombinants</u>	
			<u>Cells/ml (x 10⁻⁷)</u>	<u>Percent of Control</u>	<u>per ml (x 10⁻³)</u>	<u>per 10⁵ Survivors</u>
<u>EXPERIMENT 1</u>						
Negative control	-		7.5	100	1.5	2.0
	+		6.0	100	4	6.7
Trifluralin	-	5	8.4	112	5	5.9
	+	5	6.0	100	2	3.3
<u>EXPERIMENT 2</u>						
Negative control	-		6.3	100	1.5	2.4
	+		7.4	100	3.5	4.7
Positive control, 1,2,3,4-diepoxybutane	-	0.1	3.8	60	1,045	2,750
	+	0.1	5.2	70	903	1,737
Trifluralin	-	5	8.7	138	7	8.0
	+	5	8.4	114	3	3.6

Table 114

IN VITRO MUTAGENESIS WITH SALMONELLA TYPHIMURIUMSUMMARY DATA FOR EPA PESTICIDES
Positive Response, +; Negative Response, -

Pesticide	TA100		TA1535		TA1537		TA1538	
	- Metabolic Activation	+ Metabolic Activation	- Metabolic Activation	+ Metabolic Activation	- Metabolic Activation	+ Metabolic Activation	- Metabolic Activation	+ Metabolic Activation
Monocrotophos	-	-	-	-	-	-	-	-
Bromacil	-	-	-	-	-	-	-	-
Cacodylic Acid	-	-	-	-	-	-	-	-
Captan	+	+	+	+	-	-	-	-
Chlorpyrifos	-	-	-	-	-	-	-	-
Dinoseb	-	-	-	-	-	-	-	-
DSMA	-	-	-	-	-	-	-	-
Fenthion	-	-	-	-	-	-	-	-
Folpet	+	+	+	+	-	-	-	-
Azinphos-methyl	-	-	-	-	-	-	-	-
Malathion	-	-	-	-	-	-	-	-
Methomyl	-	-	-	-	-	-	-	-
Monuron	-	-	-	-	-	-	-	-
MSMA	-	-	-	-	-	-	-	-
Parathion	-	-	-	-	-	-	-	-
Parathion-methyl	-	-	-	-	-	-	-	-
Quintozene (PCNB)	-	-	-	-	-	-	-	-
Phorate	-	-	-	-	-	-	-	-
Simazine	-	-	-	-	-	-	-	-
Trifluralin	-	-	-	-	-	-	-	-

APPENDIX A

MUTAGENESIS STUDIES OF PESTICIDE COMPOUNDS

MOUSE HERITABLE TRANSLOCATION TEST

CAPTAN

SUMMARY

SRI conducted a heritable translocation study of Captan in mice to investigate whether heritable mutagenic events occur when the compound is ingested repeatedly over an extended period.

For 8 weeks, adult male mice were administered Captan in their diet; 60 mice received 2500 ppm, and 61 received 5000 ppm. A control group of 60 adult male mice received an untreated diet during this time. A positive control group containing 66 adult male mice was treated as a control group for 4 weeks and then received the known mutagen triethylenemelamine (TEM) in the drinking water for 4 weeks. After treatment, all males were bred with two virgin females each to produce an F_1 generation, the males of which were raised to maturity. Selected (200 per group) F_1 males were bred to three virgin females each, and presumptive translocates were rebred to three additional females each. A third breeding was conducted with selected nonbreeder and/or presumptive males.

Evaluation of the data on fertility, breeding, and litter size distribution for F_0 and F_1 generations does not suggest the presence of translocation heterozygotes in control or Captan-treated male mice. Data on dead implants and rebreeding did, however, suggest the presence of translocation heterozygotes in the group treated with 5000 ppm Captan.

Meiotic cell preparations of the testes of the presumptive males were evaluated cytogenetically. Normal meiotic chromosomes were found in the following numbers of F_1 males derived from the group specified: 8 of 8 controls, 8 of 8 from the 2500 ppm Captan group, 8 from the 5000 ppm Captan group, and 2 of 2 from the 5000 ppm Captan-treated group derived from traumatized F_0 females. Five of 5 TEM-treated F_1 males and 1 of 8 from the 5000 ppm Captan-treated males showed reciprocal translocations.

The results of this study show that under the experimental procedures employed, Captan at 5000 ppm in the diet of male mice for 8 consecutive weeks can produce a heritable mutagenic event in F₁ generation male mice.

INTRODUCTION

The EPA is reviewing and evaluating the health hazard of pesticides and of substitute candidate pesticides according to available data. Additionally, the Agency is obtaining supplemental laboratory data. The objective is to enable the EPA to select those chemicals that are minimally hazardous when used according to labeling restrictions. SRI is participating in this Substitute Chemical Program by investigating the mutagenic potential of selected materials by in vitro and in vivo procedures.

Captan has been shown to respond in a positive manner in Salmonella typhimurium, Escherichia coli WP2, Saccharomyces cerevisiae, E. coli (relative toxicity), Bacillus subtilis, WI-38 unscheduled DNA synthesis (UDS) with metabolic activation, and Drosophila melanogaster experiments. It was not positive in a mouse dominant-lethal test. Based on positive responses in both Tier I (in vitro test) and Tier II (Drosophila) mutagenic studies, it was recommended that a heritable translocation test (Tier III) in the mouse be conducted to further assess the mutagenic potential of Captan.

In this study, young adult male ICR/SIM mice from a closed, random-bred colony were administered Captan in the diet for 8 weeks. After treatment, each male was mated to two virgin females to produce an F_1 generation, the males of which were raised to maturity and bred to three virgin females each. Pregnant females were evaluated against pre-determined selection criteria for identification of suspect F_1 males, which were rebred and evaluated again. Presumptive F_1 males were examined cytogenetically.

Through this procedure, a heritable mutagenic response can be detected. Potential mutagenic effects were identified by examination of fetuses during the middle to later stages of gestation. Cytogenetic

examinations were made of meiotic cell preparations of the testes from suspect males for confirmation of findings obtained from the breeding studies.

Reported here are the results of the heritable translocation study of Captan.

MOUSE HERITABLE TRANSLOCATION TEST

Background

Human populations frequently are exposed to man-made chemicals, often at barely detectable levels, for extended periods. To evaluate the genetic hazards of such chemicals, a prudent approach is to study them in mammalian systems so as to maximize detection of a mutagenic response. The study reported here was such an investigation of Captan for its potential to produce heritable genetic defects.

Chemical induction of chromosomal aberrations in the mouse is a valuable and important experimental aid in understanding the many genetic defects due to chromosomal anomalies in humans. To date, mammalian evaluations of chemically induced chromosomal aberrations have been attempted with the dominant-lethal test and cytogenetic studies of somatic and germinal cells. Although these procedures can provide useful information, they do not measure heritable genetic effects, the most important mutagenic occurrences that are permanent and transmissible. A need exists for a method to reliably identify compounds that cause heritable chromosomal aberrations in mammalian systems. The mouse translocation procedure appears to be such a system.

A well-defined translocation test will demonstrate the fertility of an F_1 male population derived from F_0 males treated with a test agent. Confirmation of a nonbreeder, sterile, or partially sterile response can be obtained by cytological examination of the germ cells from suspected males. Sterility and partial sterility are closely correlated with the induction of translocation heterozygotes.

The procedure used in conducting this translocation test was based on experimental techniques described by Leonard and DeKnudt,¹ Cattanach et al.,² Falconer et al.,³ and Generoso.⁴ We modified this approach, in consultation with government and industry scientists actively engaged in mutagenesis research.

Materials and Methods

Animals

Male and female ICR/SIM mice were purchased from Simonsen Laboratories, Gilroy, California. The F₀ males were 8 to 10 weeks old. The females used in the breeding phases were 10- to 12-week-old virgin stock.

Chemical Supply

A supply of Captan sufficient for all aspects of the experimental program was received from Battelle Columbus Laboratory and EPA-RTP. Lot number SX-640, Chevron Chemical Company, was used for all treatment periods. The excess material has been placed in storage in case it is needed for future reference.

Dosage Selection and Compound Administration

SRI and EPA staff selected the two dosage levels of Captan to be used in this experimental program. For 8 weeks, Captan was fed in the diet at 2500 and 5000 ppm.

An appropriate amount of Captan was dissolved and/or suspended in corn oil. Then the compound-oil concentrate was added at a level of 3% to a finely ground commercial diet (Purina) of known composition. The use of corn oil assured even distribution of Captan and prevented its stratification in an otherwise dry diet. Diets prepared at 2-week intervals were refrigerated at 4°C until fed to the animals. The diet was replaced in the feed containers twice weekly to minimize the possibility of compound loss. Body weights and food consumption were recorded weekly during the 8-week exposure period.

Reference Control

Males in the reference control group were fed the Purina diet with only corn oil added at a level of 3%. These mice were treated in the same manner as those in the compound test groups. Body weights were recorded weekly, as was food consumption.

Positive Control

For the positive control group, the known mutagen triethylenemelamine (TEM) was administered in the drinking water at 0.32 mg/liter for 2 weeks and then at 0.124 mg/liter for 2 weeks. TEM treatment was initiated after the males had been on the control diet for 4 weeks. Body weights and food consumption were recorded weekly. TEM is one of the chemical mutagens that have the demonstrated effect of inducing translocations in the F_1 progeny of F_0 treated males.

Genetic Tests

After 8 weeks of treatment, the males in each treatment group were mated to two adult virgin females each. After 1 week, each female was housed individually and allowed to deliver its litter. The F_0 males were discarded. All litters were raised to weaning age, at which time the females were discarded. The F_1 males were raised to maturity. At maturity (10 to 12 weeks of age), 200 F_1 males from each experimental group were selected randomly and housed individually.

Three adult virgin females were housed with each F_1 male for the first breeding. They were examined daily for the presence of vaginal plugs. These females were sacrificed 14 days after mating, and a uterine analysis was performed for determination of the number of total, live, and dead implants. Males bred to females that produced litters fitting our criteria for presumptive classification as sterile, partially sterile, or nonbreeder were rebred to three new virgin females each. The same evaluation was made for the second breeding.

Our criteria for presumptive classification of a male as "partially sterile," "sterile," or "nonbreeder" are:

• "Partially Sterile" Male

- If all 3 females are pregnant, each must have 9 or fewer live implants, with at least 1 having 6 or fewer live implants.
- If only 2 of 3 females are pregnant, both must have 9 or fewer live implants, with 1 having 6 or fewer live implants.

- If only 1 of 3 females is pregnant, this female must have 6 or fewer live implants.
- "Sterile" Male
 - None of 3 females pregnant--previously identified by presence of vaginal plug.
- "Nonbreeder" Male
 - None of 3 females pregnant--not previously identified by presence of vaginal plug.

Any F_1 male that did not fit one of these descriptions was considered "normal" and was discarded. For each F_1 male in the control and compound-treated groups suspected of being a translocate or nonbreeder after 2 or 3 breedings, a cytogenetic evaluation was made of meiotic cell preparations of its testes. Five males from the positive control group were also subjected to cytogenetic evaluation.

Evaluation of Breeding Data

F_1 males were identified as sterile, partially sterile, or non-breeders by the methods outlined above. Individual data were totaled to give the number of observed (presumptive) translocations per treatment group, using a data base of 600 to 800 females per group. Also, for an accurate review of such findings, the F_0 breeding and litter data were thoroughly evaluated. The various measured evaluated included percentage of pregnancies, average litter size, average number of males and females, average number of males with females having zero to five or more dead implants, average number of females having zero to five or more dead implants, percentage of females with plugs, and percentage of pregnancies with and without plugs.

Meiotic Cell Cytogenetic Studies

Cytogenetic examinations were made of the testes of 31 F_1 mice, with the two testes from each mouse being examined separately. The procedures

used for the cytogenetic preparations are as follows. CO₂ was used to sacrifice the mice. The testes were removed, weighed, and placed in an isotonic solution of 2.2% sodium citrate. The tunica of each testis was punctured to release the tubules, which were then rolled on a glass plate to release the cell contents into the isotonic solution. The resulting cell suspension was centrifuged at 800 rpm for 5 minutes; the supernatant was removed, and each pellet of cells was resuspended in 5 ml of 1% sodium citrate hypotonic and held at room temperature for 15 minutes. The cells were centrifuged again at 800 rpm for 5 minutes and the supernatant was discarded. The cells were then treated with Carnoy's fixative (3 parts methyl alcohol and 1 part glacial acetic acid) to give a total volume of 5 ml, and immediately centrifuged again at 800 rpm for 5 minutes. This procedure was performed twice. Then the cells, suspended in an appropriate amount of fixative, were dropped onto clean, wet microscope slides and allowed to air-dry. The slides were stained with 2% buffered Giemsa for 5 minutes. Coverslips were attached with Permount. The slides were coded to preclude bias on the part of the scorers.

RESULTS AND DISCUSSION

General

Table 1 presents the average body weights for mice in the various groups. The body weights of the control, TEM, and 2500 ppm Captan group were within normal limits and comparable throughout the experiment. The 5000 ppm Captan-treated group showed a depressed body weight for 5 weeks before demonstrating a recovery trend during Weeks 6 to 8. This body weight depression appeared to be due to the inability of the male mice to acclimate to such a high level of compound in their diet.

Table 2 summarizes the average food consumption by treatment group. Both Captan-treated groups (2500 and 5000 ppm) showed a lower average weekly food intake than did the control and TEM-treated groups.

During the week immediately following the 1-week F_0 generation mating, one animal rack holding some females that had been mated with the mice given 5000 ppm Captan was accidentally tipped and some cages were spilled onto the floor, resulting in our inability to identify which males had been mated with these females. There were, however, sufficient numbers of F_1 generation males from those females that were not traumatized to allow us to randomly select 200 F_1 males for use in subsequent F_1 generation breedings. In addition, we held all females that had been traumatized by the tipping of the rack and maintained them throughout the remainder of the study as a separate group. From this separate group we selected 50 F_1 males (at least one per female retained) for use in the F_1 generation breedings and evaluations.

F_0 Generation

Information on the breeding performance, litter size, sex distribution, and clinical effects of the F_0 generation should be included in the evaluation of translocation data, because it may provide valuable reference data.

Table 3 summarizes the breeding and litter performance of the F_0 generation. No adverse effects were observed in the control and 2500 ppm Captan groups. The two 5000 ppm Captan groups showed a reduced pregnancy rate; the rate for the traumatized females was 29% below that of the controls. Litter sizes for the 5000 ppm groups were slightly below control values. As expected, the TEM group had a reduced pregnancy rate and a litter size 47% below the control level.

Table 4 presents litter-size distributions of live young from the F_0 generation mating. Although the distribution patterns for the control and Captan-treated animals were within normal ranges for our strain of mouse, there was evident a definite pattern of decreasing litter size and increasing variance and standard deviations between the different experimental groups. As expected, the TEM-treated animals showed the classic shift toward smaller litters. Figure 1 graphically presents the data on the F_0 generation litter-size distribution.

F_1 Generation

Table 5 summarizes breeding data from the first mating of the F_1 generation male mice. In the females mated with TEM males and with males from the 5000 ppm Captan group of traumatized F_0 mothers, there were 10% fewer females with mating plugs and an increased percentage of nonpregnant females in comparison with control values; also, these two male groups had an increased percentage of males with no pregnant females. Results from males in the 2500 and 5000 ppm Captan groups were within normal limits for this strain of mouse and comparable with values from control males.

Litter-size distributions of live implants derived from the first mating of F_1 generation males are presented in Table 6. Responses of control and Captan groups were within normal limits and readily comparable. The TEM group showed approximately a 13% reduction in litter size. Mean litter sizes were 11.72 for the control group, 11.73 for the 2500 ppm Captan group, 11.56 for the 5000 ppm Captan group, 11.88 for the 5000 ppm (traumatized F_0 female) group, and

10.24 for the TEM-treated group. The data on the F_1 generation litter-size distribution are presented graphically in Figure 2.

Tables 7 and 8 summarize the data on dead implants per F_1 male and dead implants per female, respectively. The 2500 and 5000 ppm Captan groups showed a slight increase in total dead implants for both males and females at the 4, 5, and >5 levels when compared with controls. TEM animals showed significant increases in dead implants for both males and females.

Table 9 summarizes the breeding results by treatment of those F_1 males classified as presumptive sterile, partially sterile, or non-breeders after three breedings. Table 10 identifies these F_1 males individually by number and treatment.

Details of the breeding and rebreeding data for presumptive F_1 males are presented in Table 11. In the reference control group, 24 of the 200 males were considered as presumptive translocates. When rebred, 12 males remained in this classification. A third mating of selected questionable and/or nonbreeder males reduced the number of presumptive males to eight: 1 nonbreeder, 1 presumptive sterile, and 6 partially sterile (3 of which were questionable partially sterile).

For the TEM group, 83 of 200 F_1 males were identified as presumptive mutants after the first breeding. When rebred, 53 still met the original criteria. A third breeding reduced this number to 49; 4 continued to be nonbreeders, 14 were presumptive sterile, and 31 were partially sterile (6 of which were questionable partially sterile).

In the 2500 ppm Captan group, 30 of 200 F_1 males were identified as presumptive mutants after the first breeding. When rebred, 11 still met the criteria: 5 were nonbreeders, 1 was a presumptive sterile, and 2 were partially sterile (1 of which was questionable partially sterile).

The 5000 ppm Captan group also had 30 of 200 F_1 males identified as presumptive mutants after the first breeding. The second mating reduced this number to 9. After a third breeding, 8 males still met the

original criteria: 2 were nonbreeders, 1 was a presumptive sterile, and 5 were partially sterile.

For the group of F_1 males derived from traumatized F_0 females and males treated with 5000 ppm Captan, 12 of 50 F_1 males were identified as presumptive mutants after the first mating. A second breeding reduced this number to 4 and a third breeding further reduced to 2 the number of F_1 males that still met the original criteria; both were partially sterile, with one of them being questionable partially sterile.

The data on the F_0 and F_1 generations' fertility, breeding, and litter-size distribution as well as the data on the F_1 generation's dead implants and rebreeding show that Captan tends to induce dose-related effects on the reproductive performance of male mice. The data also suggest the presence of translocation heterozygotes in the 5000 ppm Captan group.

Review of the data on dead implants, breeding, and rebreeding for the F_1 generation of the TEM-treated group showed, as expected, the potential for the presence of translocation heterozygotes in 24.5% of the F_1 males.

Cytogenetic Studies

Table 12 presents the findings from the cytogenetic evaluation of meiotic cell preparations from F_1 males in the Captan groups characterized as nonbreeder, presumptive sterile, or partially sterile. Also, eight control males and 5 of 49 TEM males were evaluated.

Whenever possible, 25 spermatocytes per testis were scored. The slides were decoded only after all scoring was completed. The results are summarized as follows:

- All eight males examined in the control group were cytogenetically normal.
- The five TEM males all showed positive reciprocal translocations.
- All eight males in the 2500 ppm Captan group were cytogenetically normal.
- Seven males in the 5000 ppm Captan group were cytogenetically normal; however, the eighth male (No. 657) showed as a positive reciprocal translocation.

- The two males examined in the 5000 ppm Captan group derived from traumatized F_0 females were cytogenetically normal.

Discussion

Increased use of the translocation procedure has revealed that a meaningful relationship exists between the incidence of dead implants in F_1 matings and the occurrence of a heritable translocation event. Previous experiments at SRI and at Oak Ridge National Laboratories have demonstrated this correlation. The following paragraphs discuss occurrences of dead implants in this study.

When females had a total implant count of less than six or when all their implants were identified as dead, we generally considered this to be a result of first breeding or of some factor other than compound treatment (such as background incidence) and excluded those females from this evaluation. Tables 13 through 17 present the total, dead, and live implantation data for suspect translocates of the control, the TEM group, and the three Captan groups.

The control group (Table 13) showed a normal implant distribution, with the exception of one F_1 male (No. 122) for whom the number of implants was high. In the TEM group (Table 14), the expected increase in dead implants and the resultant decrease in live implants occurred, although the numbers of total implants were generally normal. This pattern occurred during all breeding periods.

The 2500 ppm Captan group (Table 15) contained seven males in the first breeding with females having high dead implant counts but normal live litter size, according to the criteria. Also, 3 males showed high dead implant counts during the first breeding, with live implant counts fitting the criteria for partially sterile males. This increase in dead implant occurrence was not repeated in subsequent breeding, and all F_1 males were classified as normal after the breeding phases.

Five F_1 males in the 5000 ppm Captan group (Table 16) showed an increase in dead implants during the first breeding. In the rebreeding of these males, only male No. 657 continued to show the increase in dead implants and the resultant decrease in live implants. All other F_1 males in this group had a normal distribution of dead and live implants for all breedings.

Table 17 presents the implant data for F_1 males derived from traumatized females in the 5000 ppm Captan group. With the exception of an occasional female showing an increase in dead implants, the distribution of total, dead, and live implants was normal.

The numbers of total implantations were generally within normal limits for all experimental groups in all breedings.

REFERENCES

1. A. Leonard and G. H. DeKnuft. *Mutation Res.* 9, 127 (1970).
2. B. M. Cattanaach, C. E. Pollard, and J. H. Isaacson. *Mutation Res.* 6, 297 (1968).
3. D. S. Falconer, D. M. Slizynski, and C. Auerbach. *J. Genet.* 51, 81 (1952).
4. W. M. Generoso. *Proc. Second Annual Environmental Mutagen Soc.*, p. 9 (March 1971).

Table 1

AVERAGE BODY WEIGHTS IN GRAMS FOR MICE RECEIVING
VARIOUS LEVELS OF CAPTAN IN THEIR DIETS

<u>Week of Test</u>	<u>Control</u>	<u>TEM</u>	<u>Dietary Levels of Captan (ppm diet)</u>	
			<u>2500</u>	<u>5000</u>
Initial	33.8	33.4	33.8	33.6
1	32.5	32.9	33.3	31.8
2	33.7	34.6	33.7	31.4
3	34.8	35.4	34.4	32.1
4	34.6	35.8	34.4	31.4
5	34.8	35.9	36.0	33.0
6	37.3	38.1	37.5	35.3
7	37.7	38.9	39.3	35.8
8	39.2	39.8	38.9	36.9

Table 2

AVERAGE FOOD CONSUMPTION FOR MICE RECEIVING
 VARIOUS LEVELS OF CAPTAN IN THEIR DIETS
 (Grams of Food Consumed/Mouse/Day)

<u>Week of Test</u>	<u>Control</u>	<u>TEM</u>	<u>Dietary Levels of Captan (ppm diet)</u>	
			<u>2500</u>	<u>5000</u>
1	4.01	4.05	3.74	3.25
2	4.82	5.00	4.72	4.38
3	4.91	5.08	4.90	4.40
4	5.28	5.33	4.93	4.45
5	5.08	5.55	5.44	4.95
6	5.36	5.56	4.95	4.75
7	5.81	6.14	5.63	5.27
8	5.55	5.74	5.13	4.83

Table 3

TRANSLOCATION STUDY OF CAPTAN
F₀ GENERATION MICE
SUMMARY OF BREEDING AND LITTER DATA

<u>Parameter</u>	<u>Control</u>	<u>TEM</u>	<u>2500 ppm</u>	<u>5000 ppm</u>	<u>5000 ppm^a</u>
Number of F ₀ males	60	66	60	31	30
Number of F ₀ females	120	132	119	61	59
Number pregnant	93	77	91	38	33
Percent pregnant	77.5	58.3	76.5	62.3	55.9
Number of nonbreeder males	8	19	8	5	--
Percent nonbreeders	13.3	28.8	13.3	16.1	--
Average live litter size	12.30	6.55	12.16	11.68	11.33
Average number of males weaned/litter	5.84	3.26	5.68	6.29	5.25

^aGroup of females accidentally tipped off rack following 1 week of mating with 5000 ppm-treated mates. These traumatized females, and their offspring, most of which could not be identified back to a particular F₀ male, were considered as a separate group for the remainder of the study.

Table 4

TRANSLOCATION STUDY OF CAPTAN
 MOUSE LITTER-SIZE DISTRIBUTION OF LIVE YOUNG
 DERIVED FROM F₀ GENERATION ADULTS

Litter Size	Control	TEM	Captan (ppm diet)		
			2500	5000	5000 ^a
1	0	1	0	0	0
2	0	2	0	1	0
3	0	3	0	0	0
4	0	8	0	0	1
5	0	10	1	0	0
6	0	15	2	1	1
7	1	12	0	0	1
8	2	8	3	0	0
9	9	8	3	1	3
10	6	2	8	6	3
11	8	3	9	5	7
12	18	1	24	8	5
13	22	0	20	9	8
14	18	1	10	6	1
15	6	0	6	1	3
16	3	0	3	0	0
17	0	0	2	0	0
18	0	0	0	0	0
Mean (μ)	12.30	6.55	12.16	11.68	11.33
Variance (σ^2)	3.89	5.84	4.97	5.73	6.09
Standard Deviation (σ)	1.97	2.42	2.23	2.39	2.47

^aTraumatized F₀ females.




FIGURE 1 LITTER SIZE DISTRIBUTION F_0 GENERATION

Table 5

TRANSLOCATION STUDY OF CAPTAN
F₁ GENERATION MICE
SUMMARY OF BREEDING DATA--FIRST BREEDING

Parameter	Control	TEM	Captan (ppm)		
			2500	5000	5000 ^a
Number of F ₁ males	200	200	200	200	50
Number of females	600	600	600	600	150
Number of mating plugs	450	376	442	439	95
Percent mating plugs	75	63	74	73	63
Number pregnant	492	383	472	477	112
Percent pregnant	82	64	79	80	75
Number pregnant with plug	434	326	419	413	94
Percent pregnant with plug	88	85	89	87	84
Number pregnant without plug	58	57	53	64	18
Percent pregnant without plug	12	15	11	13	16
Number not pregnant	108	217	128	123	38
Percent not pregnant	18	36	21	20	25
Number not pregnant with plug	16	50	23	26	1
Percent not pregnant with plug	15	23	18	21	3
Males with no pregnant females	13	39	14	9	7
Percent males with no pregnant females	6.5	19.5	7.0	4.5	14.0

^aTraumatized F₀ females.

Table 6

TRANSLOCATION STUDY OF CAPTAN
 MOUSE LITTER SIZE DISTRIBUTION OF LIVE YOUNG
 DERIVED FROM F₁ GENERATION ADULTS--FIRST BREEDING

Litter Size	Control	TEM	Captan (ppm)		
			2500	5000	5000 ^a
1	5	4	4	3	0
2	4	9	0	3	0
3	2	12	5	1	0
4	4	17	6	5	1
5	5	21	4	6	1
6	8	11	5	8	4
7	7	8	6	11	3
8	14	17	13	19	4
9	35	25	35	17	2
10	37	31	40	53	16
11	73	53	77	77	14
12	80	61	75	85	15
13	89	52	87	93	17
14	72	27	54	55	14
15	32	21	36	19	13
16	16	8	14	11	5
17	7	1	7	7	1
18	1	1	2	3	0
19	0	1	0	0	0
20	0	0	1	0	0
Mean (μ)	11.72	10.24	11.73	11.56	11.88
Variance (σ^2)	8.13	13.59	8.68	7.40	7.20
Standard Deviation (σ)	2.85	3.69	2.95	2.72	2.68

^aTraumatized F₀ females.




FIGURE 2 LITTER SIZE DISTRIBUTION F₁ GENERATION — FIRST MATING

Table 7

TRANSLOCATION STUDY OF CAPTAN
SUMMARY OF DEAD IMPLANTS PER F₁ MALE

Number of Males with Females Having	Control	TEM	Captan (ppm)		
			2500	5000	5000 ^a
0 Dead implants	68	37	69	49	20
1 Dead implant	54	36	44	68	11
2 Dead implants	30	25	31	36	7
3 Dead implants	18	16	13	18	2
4 Dead implants	8	9	18	9	0
5 Dead implants	2	6	4	3	2
> 5 Dead implants	7	32	7	8	1

^aTraumatized F₀ females.

Table 8

TRANSLOCATION STUDY OF CAPTAN
SUMMARY OF DEAD IMPLANTS PER FEMALE

<u>Number of Dead Implants/Female</u>	<u>Control</u>	<u>TEM</u>	<u>Captan (ppm)</u>		
			<u>2500</u>	<u>5000</u>	<u>5000^a</u>
0	320	182	295	284	79
1	109	83	113	130	23
2	39	35	37	41	9
3	18	16	16	9	0
4	3	11	7	3	0
5	0	11	2	2	0
> 5	3	45	2	8	1
Total Pregnant Females	492	383	472	477	112

^aTraumatized F₀ female.

Table 9
 TRANSLOCATION STUDY OF CAPTAN
 SUMMARY OF PRESUMPTIVE TRANSLOCATION F₁ MALES AFTER THREE BREEDINGS

	<u>Control</u>	<u>TEM</u>	<u>Captan (ppm)</u>		
			<u>2500</u>	<u>5000</u>	<u>5000^a</u>
Total number of F ₁ males	200	200	200	200	50
Number of nonbreeder males	1	4	5	2	0
Number of presumptive sterile males	1	14	1	1	0
Number of partially sterile males	6(3?)	31(6?)	2(1?)	5	2(1?)

^aTraumatized F₀ females

Table 10

TRANSLOCATION STUDY OF CAPTAN
INDIVIDUAL IDENTIFICATION OF PRESUMPTIVE F₁ MALES
AFTER THREE BREEDINGS

<u>Treatment</u>	<u>Partially Sterile</u>	<u>Presumptive Sterile</u>	<u>Nonbreeder</u>
Control	16? 65 122? 164? 189 190	72	220
TEM	215 216 232 235? 238 245 262 264 269 280 281 290 292 299? 300 314 315 327? 344 345 350 359 360? 361 371 375 376 388 390 399? 400?	202 231 251 256 268 273 288 317 321 326 339 343 369 389	220 241 391 396

(Continued)

Table 10
(Concluded)

<u>Treatment</u>	<u>Partially Sterile</u>	<u>Presumptive Sterile</u>	<u>Nonbreeder</u>
Captan	480?	474	449
2500 ppm	526		479
			496
			523
			583
Captan	635	736	634
5000 ppm	657		733
	760		
	779		
	780		
Captan	805		
5000 ppm ^a	837?		

^aTraumatized F₀ females.

Table 11

TRANSLOCATION STUDY OF CAPTAN
BREEDING AND REBREEDING SUMMARY OF PRESUMPTIVE F₁ MALES
(Live Implants Only)

Treatment	F ₁ Male No.	First Breeding (3 Females)			Second Breeding (3 Females)			Third Breeding (3 Females)		
Control	11	**	-	-	13	8	(15) [†]	-	-	-
	16	-	-	(13)	-	-	-	-	-	-
	17	-	-	-	-	-	-	-	-	-
	24	-	-	-	0*	11	-	-	-	-
	39	9	0	-	0	-	-	12	11	11
	43	-	-	-	-	-	-	10	-	-
	50	4	7	6	12	11	12	-	-	-
	59	0	-	-	10	9	-	-	-	-
	65	4	0	5	3	6	2	-	-	-
	67	-	-	-	14	11	-	-	-	-
	72	-	-	-	0	-	-	-	-	-
	86	-	-	-	(12)	-	-	-	-	-
	102	-	-	(11)	9	5	16	12	12	4
	122	6	2	13	7	7	14	-	-	-
	129	-	-	-	-	-	-	14	-	-
	139	-	-	-	14	-	-	-	-	-
	144	-	-	-	9	10	-	-	-	-
	152	3	9	9	10	8	13	-	-	-
	164	-	-	-	-	-	(9)	-	-	-
	179	-	-	-	10	-	-	-	-	-
189	0	1	0	0	0	0	0	1	0	
190	1	4	6	1	0	0	0	1	1	
192	7	(14)	-	12	11	-	-	-	-	
200	-	-	-	13	-	-	-	-	-	
	Totals		24			12			8	
TEM	202	0	0	0	0	0	0	-	-	-
	215	-	-	-	7	4	-	-	-	-
	216	-	-	(9)	-	-	-	4	-	-
	220	-	-	-	-	-	-	-	-	-
	226	-	-	-	12	2	-	-	-	-
	227	0	4	2	12	4	8	-	-	-
	228	-	(9)	-	14	12	11	-	-	-
	229	7	0	9	14	14	13	-	-	-
	230	-	-	-	11	10	-	-	-	-

(Continued)

*"0" indicates a plug was observed for a female that was not pregnant.

**"- " indicates a plug was not detected and the female was not pregnant.

[†]"(")" indicates all implants were in early stages of development and impossible to determine if they were live or dead upon gross observation.

Table 11
(Continued)

Treatment	F1 Male No.	First Breeding (3 Females)			Second Breeding (3 Females)			Third Breeding (3 Females)		
TEM	231	0*	-**	-	0	0	0			
	232	3	4	1	8	2	5			
	233	-	-	-	11	15	6			
	235	8	7	6	7	7	13			
	237	5	9	8	10	10	14			
	238	3	0	1	1	2	(P)†			
	239	-	-	-	13	-	-			
	241	-	-	-	-	-	-	-	-	-
	244	-	-	-	10	13	11			
	245	(P)	0	1	1	0	3			
	251	0	0	0	0	0	0			
	253	3	10	-	14	11	-			
	254	4	-	-	13	14	11			
	256	0	-	-	-	-	-	-	-	-
	258	0	-	-	13	0	-			
	262	2	8	5	5	4	5			
	264	3	5	-	(P)	5	9			
	267	-	-	-	13	(P)	12			
	268	-	-	-	-	-	-	0	-	-
	269	-	-	-	6	7	-			
	270	0	-	-	8	-	-	9	-	-
	273	-	-	-	0	-	-	-	-	-
	280	9	7	2	8	8	5			
	281	(P)	6	-	4	1	2			
	283	9	9	8	14	0	-			
	286	-	-	-	-	-	-	10	-	-
	288	0	0	0	0	0	0			
	290	5	5	4	7	5	0			
	292	6	3	0	3	3	8			
	294	8	(18)†	9	10	12	11			
	299	8	-	-	9	4	9			
	300	6	(15)	-	8	5	2			
	302	9	-	-	13	12	15			
	314	-	-	-	2	1	4			
	315	0	3	0	0	0	0			
	317	0	0	0	0	0	-			
	318	4	4	3	9	4	11			
	321	0	0	-	0	0	-			
	322	9	-	-	11	11	-			

(Continued)

*"0" indicates a plug was observed for a female that was not pregnant.

**"- " indicates a plug was not detected and the female was not pregnant.

†"()" indicates all implants were in early stages of development and impossible to determine if they were live or dead upon gross observation.

†† "(P)" indicates female was pregnant but had no live implants.

Table 11
(Continued)

Treatment	F1 Male No.	First Breeding (3 Females)			Second Breeding (3 Females)			Third Breeding (3 Females)		
TEM	326	**	-	-	0*	-	-	-	-	-
	327	0	-	-	0	(10)†	-	1	(9)	-
	333	0	6	-	10	0	-	-	-	-
	334	-	-	-	-	-	-	12	12	-
	339	0	-	-	0	0	0	-	-	-
	343	0	0	0	0	0	0	-	-	-
	344	3	5	-	5	4	5	-	-	-
	345	6	4	5	4	4	3	-	-	-
	346	-	-	-	12	11	-	-	-	-
	349	-	-	(12)	10	8	10	-	-	-
	350	4	3	2	Ⓟ††	0	1	-	-	-
	355	-	-	-	13	0	-	-	-	-
	356	-	-	-	11	-	(15)	-	-	-
	359	9	4	8	7	3	7	-	-	-
	360	(15)	-	-	2	Ⓟ	-	-	-	-
	361	4	4	9	3	7	Ⓟ	-	-	-
	363	5	-	-	12	14	-	-	-	-
	367	-	-	-	4	5	11	-	-	-
	369	0	-	-	-	-	-	-	-	-
	371	4	6	-	5	5	5	-	-	-
	372	(16)	-	-	10	9	-	-	-	-
	375	3	-	-	5	-	7	-	-	-
	376	5	0	6	1	-	1	-	-	-
	381	-	-	-	11	0	-	-	-	-
	382	(13)	(11)	-	-	-	-	11	10	-
	385	9	5	(6)	10	12	7	-	-	-
	387	-	-	-	15	-	-	-	-	-
	388	3	4	0	3	3	4	-	-	-
	389	-	-	-	-	-	-	0	-	-
	390	4	3	5	3	5	3	-	-	-
	391	-	-	-	-	-	-	-	-	-
	396	-	-	-	-	-	-	-	-	-
	397	-	-	(14)	11	0	0	-	-	-
	399	5	4	(11)	5	2	4	-	-	-
	400	2	2	(12)	4	5	2	-	-	-
	Totals		83			53			49	

(Continued)

*"0" indicates a plug was observed for a female that was not pregnant.

**"- " indicates a plug was not detected and the female was not pregnant.

†"()" indicates all implants were in early stages of development and impossible to determine if they were live or dead upon gross observation.

††"Ⓟ" indicates female was pregnant but had no live implants.

Table 11
(Continued)

Treatment	F ₁ Male No.	First Breeding (3 Females)			Second Breeding (3 Females)			Third Breeding (3 Females)		
Captan 2500 ppm	430	-**	-	-	12	8	12	-	-	-
	432	4	(17)†	9	11	10	12	-	-	-
	436	9	5	8	11	12	13	-	-	-
	449	-	-	-	-	-	-	-	-	-
	450	0*	1	14	10	13	13	-	-	-
	451	7	0	-	0	0	10	-	-	-
	452	9	10	10	13	11	13	-	-	-
	455	-	-	-	10	12	11	-	-	-
	461	-	-	-	13	11	9	-	-	-
	469	-	-	-	-	-	11	-	-	-
	472	8	10	9	0	11	10	-	-	-
	474	-	-	-	0	-	-	-	-	-
	479	-	-	-	-	-	-	-	-	-
	480	(3)	0	-	(13)	-	8	2	-	-
	484	4	14	11	11	0	12	-	-	-
	489	11	12	6	9	11	14	-	-	-
	496	-	-	-	-	-	-	-	-	-
	518	-	-	-	0	11	13	-	-	-
	523	-	-	-	-	-	-	-	-	-
	526	7	3	(1)	0	5	0	2	1	0
	528	0	-	-	14	-	-	-	-	-
546	-	-	-	-	-	-	9	13	-	
561	-	-	-	-	-	-	15	11	-	
568	8	(9)	(13)	9	11	8	-	-	-	
569	-	-	-	11	-	10	-	-	-	
582	-	-	-	8	13	7	-	-	-	
583	-	-	-	-	-	-	-	-	-	
585	9	10	12	12	13	13	-	-	-	
588	-	-	(12)	9	-	-	11	-	-	
590	8	0	9	0	11	12	-	-	-	
	Totals		30		11		8			
Captan 5000 ppm	601	-	(11)	(8)	11	0	-	-	-	
	604	7	9	9	9	10	14	-	-	
	620	8	-	-	9	-	-	-	-	
	622	-	-	(9)	0	12	-	-	-	
	626	-	-	-	(12)	12	-	-	-	
	627	8	8	8	10	6	(P)††	-	-	
	634	-	-	-	-	-	-	-	-	
	635	-	-	-	-	(P)	-	-	-	

(Continued)

* "0" indicates a plug was observed for a female that was not pregnant.

** "-" indicates a plug was not detected and the female was not pregnant.

† "()" indicates all implants were in early stages of development and impossible to determine if they were live or dead upon gross observation.

†† "(P)" indicates female was pregnant but had no live implants.

Table 11
(Concluded)

Treatment	F1 Male No.	First Breeding (3 Females)			Second Breeding (3 Females)			Third Breeding (3 Females)		
Captan 5000 ppm	638	10	6	6	10	12	11			
	646	10	-**	-	11	13	9			
	653	0*	(13)†	-	14	13	10			
	657	2	5	1	2	1	4			
	660	0	10	8	14	9	12			
	668	-	-	6	9	12	12			
	671	-	-	-	14	11	-			
	672	-	-	-	-	-	14			
	679	6	0	8	7	12	3			
	722	-	-	-	-	-	-	12	-	-
	732	4	-	-	12	-	-			
	733	-	-	-	-	-	-	-	-	-
	734	-	-	-	11	12	-			
	736	-	-	-	-	-	-	0	-	-
	743	0	-	(15)	14	10	10			
	760	7	5	8	1	0	0			
	761	9	-	-	5	13	11			
	765	9	10	-	12	11	9			
	767	10	-	-	12	13	12			
	769	8	0	(12)	12	13	10			
779	2	0	1	0	4	-				
780	4	1	5	0	2	4				
	Totals		30			9			8	
Captan 5000 ppm ^a	805	-	-	-	-	-	2	-	-	-
	806	-	-	-	12	14	10			
	815	6	9	7	13	11	10			
	816	-	(10)	(11)	9	-	-			
	817	-	-	-	7	0	-	11	2	13
	818	10	6	-	11	8	12			
	822	-	-	-	6	-	-	10	12	12
	823	-	-	-	13	-	-			
	837	-	-	-	7	-	-	-	-	-
	841	-	-	-	13	10	-			
846	5	-	-	10	12	12				
847	(14)	-	-	10	10	14				
	Totals		12			4			2	

^a Traumatized F_0 females.

* "0" indicates a plug was observed for a female that was not pregnant.

** "-" indicates a plug was not detected and the female was not pregnant.

† "()" indicates all implants were in early stages of development and impossible to determine if they were live or dead upon gross observation.

Table 12

TRANSLOCATION STUDY OF CAPTAN
CYTOGENETIC EVALUATION OF F₁ MALE MICE

<u>Treatment</u>	<u>F₁ Male No.</u>	<u>Body Weight (g)</u>	<u>Testes Weight (mg)</u>	<u>Classification-Based Upon Breeding Data</u>	<u>Cytogenetic Classification</u>	
Control	16	64.6	265	Partially sterile (questionable)	Normal	
	17	58.5	307	Nonbreeder	Normal	
	65	38.8	237	Partially sterile	Normal	
	72	68.0	232	Presumptive sterile	Normal	
	122	48.7	289	Partially sterile (questionable)	Normal	
	164	55.8	314	Partially sterile (questionable)	Normal	
	189	49.6	245	Partially sterile	Normal	
	190	47.5	222	Partially sterile	Normal	
	TEM	232	54.6	276	Partially sterile	Positive reciprocal translocation
		262	41.4	243	Partially sterile	Positive reciprocal translocation
290		62.6	222	Partially sterile	Positive reciprocal translocation	
345		54.1	294	Partially sterile	Positive reciprocal translocation	
361		50.4	278	Partially sterile	Positive reciprocal translocation	
Captan 2500 ppm	449	60.2	256	Nonbreeder	Normal	
	474	56.5	175	Presumptive sterile	Normal	
	479	65.1	287	Nonbreeder	Normal	
	480	58.7	307	Partially sterile (questionable)	Normal	
	496	60.9	242	Nonbreeder	Normal	
	523	50.7	348	Nonbreeder	Normal	
	526	54.1	295	Partially sterile	Normal	
	583	50.7	297	Nonbreeder	Normal	

Table 12
(Concluded)

<u>Treatment</u>	<u>F1 Male No.</u>	<u>Body Weight (g)</u>	<u>Testes Weight (mg)</u>	<u>Classification-Based Upon Breeding Data</u>	<u>Cytogenetic Classification</u>
Captan	634	57.1	299	Nonbreeder	Normal
5000 ppm	635	56.0	312	Partially sterile	Normal
	657	61.0	231	Partially sterile	Positive reciprocal translocation
	733	62.2	256	Nonbreeder	Normal
	736	57.9	256	Presumptive sterile	Normal
	760	55.9	298	Partially sterile	Normal
	779	54.8	224	Partially sterile	Normal
	780	53.0	295	Partially sterile	Normal
Captan	805	70.0	268	Partially sterile	Normal
5000 ppm ^a	837	56.9	303	Partially sterile (questionable)	Normal

^aTraumatized F₀ females.

Table 13

TRANSLOCATION STUDY OF CAPTAN
IMPLANTATION SUMMARY OF PRESUMPTIVE F₁ MALES

	F ₁ Male Number	Female Number	Control Group			Initial Classification
			Total Implantations	Dead Implantations	Live Implantations	
First breeding	11	1	-**	-	-	Nonbreeder
		2	-	-	-	
		3	-	-	-	
	16	1	-	-	-	Partially sterile (questionable)
		2	-	-	-	
		3	(13)†	?	(13)	
	17	1	-	-	-	Nonbreeder
		2	-	-	-	
		3	-	-	-	
	24	1	-	-	-	Nonbreeder
		2	-	-	-	
		3	-	-	-	
	39	1	9 [*]	0	9	Normal
		2	0 [*]	0	0	
		3	-	-	-	
	43	1	-	-	-	Nonbreeder
		2	-	-	-	
		3	-	-	-	
	50	1	4	0	4	Partially sterile
		2	8	1	7	
		3	7	1	6	
	59	1	0	0	0	Presumptive sterile
		2	-	-	-	
		3	-	-	-	
	65	1	5	1	4	Partially sterile
		2	0	0	0	
		3	5	0	5	
	67	1	-	-	-	Nonbreeder
		2	-	-	-	
		3	-	-	-	
	72	1	-	-	-	Nonbreeder
		2	-	-	-	
		3	-	-	-	
	86	1	-	-	-	Nonbreeder
		2	-	-	-	
		3	-	-	-	
	102	1	-	-	-	Partially sterile (questionable)
		2	-	-	-	
		3	(11)	?	(11)	
	122	1	16	10	6	Normal
		2	4	2	2	
		3	13	0	13	

^{*}"0" indicates a plug was observed for a female that was not pregnant.

^{**}"-" indicates a plug was not detected and the female was not pregnant.

[†]"(")" indicates all implants were in early stages of development, and thus difficult to determine if they were live or dead upon gross observation.

Table 13 (Continued)

TRANSLOCATION STUDY OF CAPTAN
IMPLANTATION SUMMARY OF PRESUMPTIVE F₁ MALES

F ₁ Male Number	Female Number	Total Implantations	Control Group		Initial Classification	
			Dead Implantations	Live Implantations		
First breeding (concl.)	129	1	**	-	-	Nonbreeder
		2	-	-	-	
		3	-	-	-	
	139	1	-	-	-	Nonbreeder
		2	-	-	-	
		3	-	-	-	
	144	1	-	-	-	Nonbreeder
		2	-	-	-	
		3	-	-	-	
	152	1	7	4	3	Partially sterile
		2	12	3	9	
		3	9	0	9	
	164	1	-	-	-	Nonbreeder
		2	-	-	-	
		3	-	-	-	
	179	1	-	-	-	Nonbreeder
		2	-	-	-	
		3	-	-	-	
	189	1	0 ⁺	0	0	Partially sterile
		2	1	0	1	
		3	0	0	0	
	190	1	1	0	1	Partially sterile
		2	4	0	4	
		3	6	0	6	
	192	1	7	0	7	Partially sterile (questionable)
		2	(14) ⁺	?	(14)	
		3	-	-	-	
	200	1	-	-	-	Nonbreeder
		2	-	-	-	
		3	-	-	-	
<u>Final Classification</u>						
Second breeding	11	1	13	0	13	Normal
		2	9	1	8	
		3	(15)	?	(15)	
	16	1	-	-	-	Rebred ^h
		2	-	-	-	
		3	-	-	-	
	17	1	-	-	-	Rebred ^b
		2	-	-	-	
		3	-	-	-	

^{*}"0" indicates a plug was observed for a female that was not pregnant.

^{**}"-" indicates a plug was not detected and the female was not pregnant.

⁺"(+)" indicates all implants were in early stages of development, and thus difficult to determine if they were live or dead upon gross observation.

^bSee third breeding for final classification.

Table 13 (Continued)

TRANSLOCATION STUDY OF CAPTAN
 IMPLANTATION SUMMARY OF PRESUMPTIVE F₁ MALES

	F ₁ Male Number	Female Number	Control Group			Final Classification
			Total Implantations	Dead Implantations	Live Implantations	
Second breeding (cont.)	24	1	0 ⁺	0	0	Normal
		2	12 ^{**}	1	11	
		3	-	-	-	
	39	1	0	0	0	Rebred ^b
		2	-	-	-	
		3	-	-	-	
	43	1	-	-	-	Rebred ^b
		2	-	-	-	
		3	-	-	-	
	50	1	12	0	12	Normal
		2	11	0	11	
		3	12	0	12	
	59	1	10	0	10	Normal
		2	12	3	9	
		3	-	-	-	
	65	1	3	0	3	Partially sterile
		2	6	0	6	
		3	2	0	2	
	67	1	15	1	14	Normal
		2	11	0	11	
		3	-	-	-	
	72	1	0	0	0	Rebred ^b
		2	-	-	-	
		3	-	-	-	
	86	1	(12) ⁺	?	(12)	Rebred ^b
		2	-	-	-	
		3	-	-	-	
	102	1	10	1	9	Normal
		2	6	1	5	
		3	16	0	16	
	122	1	14	7	7	Normal
		2	14	7	7	
		3	2	8	4	
	129	1	-	-	-	Rebred ^b
		2	-	-	-	
		3	-	-	-	
	139	1	15	1	14	Normal
		2	-	-	-	
		3	-	-	-	
	144	1	9	0	9	Normal
		2	10	0	10	
		3	-	-	-	

^{**}"0" indicates a plug was observed for a female that was not pregnant.

^{**}"-" indicates a plug was not detected and the female was not pregnant.

⁺"(") indicates all implants were in early stages of development, and thus difficult to determine if they were live or dead upon gross observation.

^bSee third breeding for final classification.

Table 13 (Continued)

TRANSLOCATION STUDY OF CAPTAN
IMPLANTATION SUMMARY OF PRESUMPTIVE F₁ MALES

	F ₁ Male Number	Female Number	Control Group			Final Classification
			Total Implantations	Dead Implantations	Live Implantations	
Second breeding (concl.)	152	1	11	1	10	Normal
		2	8	0	8	
		3	14	1	13	
	164	1	-**	-	-	Rebred ^b
		2	-	-	-	
		3	(9)†	?	(9)	
	179	1	11	1	10	Normal
		2	-	-	-	
		3	-	-	-	
	189	1	0*	0	0	Rebred ^b
		2	0	0	0	
		3	0	0	0	
	190	1	1	0	1	Rebred ^b
		2	0	0	0	
		3	0	0	0	
192	1	13	1	12	Normal	
	2	12	1	11		
	3	-	-	-		
200	1	13	0	13	Normal	
	2	-	-	-		
	3	-	-	-		
Third breeding	16	1	-	-	-	Partially sterile (questionable)
		2	-	-	-	
		3	-	-	-	
	17	1	-	-	-	Nonbreeder
		2	-	-	-	
		3	-	-	-	
	39	1	12	0	12	Normal
		2	12	1	11	
		3	11	0	11	
	43	1	10	0	10	Normal
		2	-	-	-	
		3	-	-	-	
	72	1	-	-	-	Presumptive sterile
		2	-	-	-	
		3	-	-	-	
86	1	14	2	12	Normal	
	2	14	2	12		
	3	5	1	4		

*"0" indicates a plug was observed for a female that was not pregnant.

**"- " indicates a plug was not detected and the female was not pregnant.

†"()" indicates all implants were in early stages of development, and thus difficult to determine if they were live or dead upon gross observation.

^bSee third breeding for final classification.

Table 13 (Concluded)

TRANSLOCATION STUDY OF CAPTAN
 IMPLANTATION SUMMARY OF PRESUMPTIVE F₁ MALES

	F ₁ Male Number	Female Number	Control Group			Final Classification
			Total Implantations	Dead Implantations	Live Implantations	
Third breeding (concl.)	129	1	14	0	14	Normal
		2	**	-	-	
		3	-	-	-	
	164	1	-	-	-	Partially sterile (questionable)
		2	-	-	-	
		3	-	-	-	
	189	1	0*	0	0	Partially sterile
		2	1	0	1	
		3	-	-	-	
	190	1	0	0	0	Partially sterile
		2	1	0	1	
		3	1	0	1	

*"0" indicates a plug was observed for a female that was not pregnant.

**"-." indicates a plug was not detected and the female was not pregnant.

Table 14

TRANSLOCATION STUDY OF CAPTAN
 IMPLANTATION SUMMARY OF PRESUMPTIVE F₁ MALES

F ₁ Male Number	Female Number	TEM Group			Initial Classification
		Total Implantations	Dead Implantations	Live Implantations	
First breeding	1	0*	0	0	Presumptive sterile
	2	0	0	0	
	3	0	0	0	
215	1	-**	-	-	Nonbreeder
	2	-	-	-	
	3	-	-	-	
216	1	-	-	-	Partially sterile (questionable)
	2	-	-	-	
	3	(9) ⁺	?	(9)	
220	1	-	-	-	Nonbreeder
	2	-	-	-	
	3	-	-	-	
226	1	-	-	-	Nonbreeder
	2	-	-	-	
	3	-	-	-	
227	1	0	0	0	Partially sterile
	2	11	7	4	
	3	11	9	2	
228	1	-	-	-	Partially sterile (questionable)
	2	(9)	?	(9)	
	3	-	-	-	
229	1	8	1	7	Normal
	2	0	0	0	
	3	10	1	9	
230	1	-	-	-	Nonbreeder
	2	-	-	-	
	3	-	-	-	
231	1	0	0	0	Presumptive sterile
	2	-	-	-	
	3	-	-	-	
232	1	12	9	3	Partially sterile
	2	9	5	4	
	3	10	9	1	
233	1	-	-	-	Nonbreeder
	2	-	-	-	
	3	-	-	-	
235	1	15	7	8	Partially sterile
	2	11	4	7	
	3	11	5	6	
237	1	5	0	5	Partially sterile
	2	12	3	9	
	3	12	4	8	

*"0" indicates a plug was observed for a female that was not pregnant.

**"-" indicates a plug was not detected and the female was not pregnant.

*"()" indicates all implants were in early stages of development, and thus difficult to determine if they were live or dead upon gross observation.

Table 14 (Continued)

TRANSLOCATION STUDY OF CAPTAN
IMPLANTATION SUMMARY OF PRESUMPTIVE F₁ MALES

F ₁ Male Number	Female Number	TEM Group			Initial Classification	
		Total Implantations	Dead Implantations	Live Implantations		
First breeding (cont.)	238	1	14	11	3	Partially sterile
		2	0*	0	0	
		3	9	8	1	
239	1	-**	-	-	-	Nonbreeder
		2	-	-	-	
		3	-	-	-	
241	1	-	-	-	-	Nonbreeder
		2	-	-	-	
		3	-	-	-	
244	1	-	-	-	-	Nonbreeder
		2	-	-	-	
		3	-	-	-	
245	1	6	6	0	0	Partially sterile
		2	0	0	0	
		3	4	3	1	
251	1	0	0	0	0	Presumptive sterile
		2	0	0	0	
		3	0	0	0	
253	1	12	9	3	3	Normal
		2	12	2	10	
		3	-	-	-	
254	1	6	2	4	4	Partially sterile
		2	-	-	-	
		3	-	-	-	
256	1	0	0	0	0	Presumptive sterile
		2	-	-	-	
		3	-	-	-	
258	1	0	0	0	0	Presumptive sterile
		2	-	-	-	
		3	-	-	-	
262	1	12	10	2	2	Partially sterile
		2	12	4	8	
		3	15	10	5	
264	1	8	5	3	3	Partially sterile
		2	8	3	5	
		3	-	-	-	
267	1	-	-	-	-	Nonbreeder
		2	-	-	-	
		3	-	-	-	
268	1	-	-	-	-	Nonbreeder
		2	-	-	-	
		3	-	-	-	
269	1	-	-	-	-	Nonbreeder
		2	-	-	-	
		3	-	-	-	

*"0" indicates a plug was observed for a female that was not pregnant.

"-" indicates a plug was not detected and the female was not pregnant.

Table 14 (Continued)

TRANSLOCATION STUDY OF CAPTAN
 IMPLANTATION SUMMARY OF PRESUMPTIVE F₁ MALES

F ₁ Male Number	Female Number	TEM Group			Initial Classification
		Total Implantations	Dead Implantations	Live Implantations	
First breeding (cont.)	270	1	0*	0	Presumptive sterile
		2	-**	-	
		3	-	-	
273	1	-	-	-	Nonbreeder
	2	-	-	-	
	3	-	-	-	
280	1	13	4	9	Partially sterile
	2	17	5	7	
	3	3	1	2	
281	1	4	4	0	Partially sterile
	2	12	6	6	
	3	-	-	-	
283	1	12	3	9	Normal
	2	11	2	9	
	3	9	1	8	
286	1	-	-	-	Nonbreeder
	2	-	-	-	
	3	-	-	-	
288	1	0	0	0	Presumptive sterile
	2	0	0	0	
	3	0	0	0	
290	1	9	4	5	Partially sterile
	2	14	9	5	
	3	11	7	4	
292	1	14	8	6	Partially sterile
	2	10	7	3	
	3	0	0	0	
294	1	8	0	8	Partially sterile (questionable)
	2	(18)†	?	(18)	
	3	11	2	9	
299	1	11	3	8	Normal
	2	-	-	-	
	3	-	-	-	
300	1	9	3	6	Partially sterile (questionable)
	2	(15)	?	(15)	
	3	-	-	-	
302	1	11	2	9	Normal
	2	-	-	-	
	3	-	-	-	
314	1	-	-	-	Nonbreeder
	2	-	-	-	
	3	-	-	-	

*"0" indicates a plug was observed for a female that was not pregnant.

**"-." indicates a plug was not detected and the female was not pregnant.

†"()" indicates all implants were in early stages of development, and thus difficult to determine if they were live or dead upon gross observation.

Table 14 (Continued)

TRANSLOCATION STUDY OF CAPTAN
IMPLANTATION SUMMARY OF PRESUMPTIVE F₁ MALES

F ₁ Male Number	Female Number	TEM Group			Initial Classification	
		Total Implantations	Dead Implantations	Live Implantations		
First breeding (cont.)	315	1	0*	0		
		2	3	0	3	Partially sterile
		3	0	0	0	
317	1	0	0	0		
		2	**	-	-	Presumptive sterile
		3	-	-	-	
318	1	9	5	4		
		2	13	9	4	Partially sterile
		3	13	10	3	
321	1	0	0	0		
		2	0	0	0	Presumptive sterile
		3	-	-	-	
322	1	10	1	9		
		2	-	-	-	Normal
		3	-	-	-	
326	1	-	-	-		
		2	-	-	-	Nonbreeder
		3	-	-	-	
327	1	0	0	0		
		2	-	-	-	Presumptive sterile
		3	-	-	-	
333	1	0	0	0		
		2	11	5	6	Partially sterile
		3	-	-	-	
334	1	-	-	-		
		2	-	-	-	Nonbreeder
		3	-	-	-	
339	1	0	0	0		
		2	-	-	-	Presumptive sterile
		3	-	-	-	
343	1	0	0	0		
		2	0	0	0	Presumptive sterile
		3	0	0	0	
344	1	10	7	3		
		2	12	7	5	Partially sterile
		3	-	-	-	
345	1	14	8	6		
		2	10	6	4	Partially sterile
		3	10	5	5	
346	1	-	-	-		
		2	-	-	-	Nonbreeder
		3	-	-	-	
349	1	-	-	-	Partially sterile	
		2	-	-	(questionable)	
		3	(12)*	?	(12)	

*"0" indicates a plug was observed for a female that was not pregnant.

"-" indicates a plug was not detected and the female was not pregnant.

*"()" indicates all implants were in early stages of development, and thus difficult to determine if they were live or dead upon gross observation.

Table 14 (Continued)

TRANSLOCATION STUDY OF CAPTAN
 IMPLANTATION SUMMARY OF PRESUMPTIVE F₁ MALES

	F ₁ Male Number	Female Number	Total Implantations	TEM Group		Initial Classification
				Dead Implantations	Live Implantations	
First breeding (cont.)	350	1	12	8	4	Partially sterile
		2	11	8	3	
		3	8	6	2	
	355	1	- ^{Am}	-	-	Nonbreeder
		2	-	-	-	
		3	-	-	-	
	356	1	-	-	-	Nonbreeder
		2	-	-	-	
		3	-	-	-	
	359	1	15	6	9	Partially sterile
		2	12	8	4	
		3	16	8	8	
	360	1	(15) ⁺	?	(15)	Partially sterile (questionable)
		2	-	-	-	
		3	-	-	-	
	361	1	12	8	4	Partially sterile
		2	14	10	4	
		3	11	2	9	
	363	1	9	4	5	Partially sterile
		2	-	-	-	
		3	-	-	-	
	367	1	-	-	-	Nonbreeder
		2	-	-	-	
		3	-	-	-	
	369	1	0 ^o	0	0	Presumptive sterile
		2	-	-	-	
		3	-	-	-	
	371	1	10	6	4	Partially sterile
		2	13	7	6	
		3	-	-	-	
	372	1	(16)	?	(16)	Partially sterile (questionable)
		2	-	-	-	
		3	-	-	-	
	375	1	14	11	3	Partially sterile
		2	-	-	-	
		3	-	-	-	
	376	1	11	6	5	Partially sterile
		2	0	0	0	
		3	10	4	6	
	381	1	-	-	-	Nonbreeder
		2	-	-	-	
		3	-	-	-	

^o"0" indicates a plug was observed for a female that was not pregnant.

^{Am}"-" indicates a plug was not detected and the female was not pregnant.

⁺"(") indicates all implants were in early stages of development, and thus difficult to determine if they were live or dead upon gross observation.

Table 14 (Continued)

TRANSLOCATION STUDY OF CAPTAN
 IMPLANTATION SUMMARY OF PRESUMPTIVE F₁ MALES

	F ₁ Male Number	Female Number	Total Implantations	TEM Group		Initial Classification
				Dead Implantations	Live Implantations	
First breeding (concl.)	382	1	(13) ⁺	?	(13)	Partially sterile (questionable)
		2	(11)	?	(11)	
		3	- ^{**}	-	-	
	385	1	9	0	9	Partially sterile
		2	6	1	5	
		3	(6)	?	(6)	
	387	1	-	-	-	Nonbreeder
		2	-	-	-	
		3	-	-	-	
	388	1	14	11	3	Partially sterile
		2	9	5	4	
		3	0 ⁺	0	0	
	389	1	-	-	-	Nonbreeder
		2	-	-	-	
		3	-	-	-	
390	1	11	7	4	Partially sterile	
	2	12	9	3		
	3	14	8	5		
391	1	-	-	-	Nonbreeder	
	2	-	-	-		
	3	-	-	-		
396	1	-	-	-	Nonbreeder	
	2	-	-	-		
	3	-	-	-		
397	1	-	-	-	Partially sterile (questionable)	
	2	-	-	-		
	3	(14)	?	(14)		
399	1	11	6	5	Partially sterile (questionable)	
	2	9	5	4		
	3	(11)	?	(11)		
400	1	12	10	2	Partially sterile (questionable)	
	2	9	7	2		
	3	(12)	?	(12)		
<u>Final Classification</u>						
Second breeding	202	1	0	0	0	Presumptive sterile
		2	0	0	0	
		3	0	0	0	
	215	1	11	4	7	Partially sterile
		2	?	5	4	
		3	-	-	-	

⁺"0" indicates a plug was observed for a female that was not pregnant.

^{**}"-" indicates a plug was not detected and the female was not pregnant.

⁺"(")" indicates all implants were in early stages of development, and thus difficult to determine if they were live or dead upon gross observation.

Table 14 (Continued)

TRANSLOCATION STUDY OF CAPTAN
IMPLANTATION SUMMARY OF PRESUMPTIVE F₁ MALES

F ₁ Male Number	Female Number	TFM Group			Final Classification
		Total Implantations	Dead Implantations	Live Implantations	
Second breeding (cont.)	216	1	-**	-	Rebred ^b
		2	-	-	
		3	-	-	
	220	1	-	-	Rebred ^b
		2	-	-	
		3	-	-	
	226	1	14	2	Normal
		2	5	3	
		3	-	-	
	227	1	12	0	Normal
		2	11	7	
		3	8	0	
	228	1	14	0	Normal
		2	12	0	
		3	12	1	
	229	1	15	1	Normal
		2	14	0	
		3	13	0	
	230	1	11	0	Normal
		2	10	0	
		3	-	-	
	231	1	0*	0	Presumptive sterile
		2	0	0	
		3	0	0	
	232	1	17	9	Partially sterile
		2	12	10	
		3	15	10	
	233	1	11	0	Normal
		2	15	1	
		3	7	1	
	235	1	11	4	Normal
		2	13	6	
		3	14	1	
	237	1	11	1	Normal
		2	14	4	
		3	14	1	
	238	1	9	8	Partially sterile
		2	12	10	
		3	13	13	
	239	1	13	0	Normal
		2	-	-	
		3	-	-	
	241	1	-	-	Rebred ^b
		2	-	-	
		3	-	-	

*"0" indicates a plug was observed for a female that was not pregnant.

"-" indicates a plug was not detected and the female was not pregnant.

^bSee third breeding for final classification.

Table 14 (Continued)

TRANSLOCATION STUDY OF CAPTAN
 IMPLANTATION SUMMARY OF PRESUMPTIVE F₁ MALES

	F ₁ Male Number	Female Number	TEM Group			Final Classification
			Total Implantations	Dead Implantations	Live Implantations	
Second breeding (cont.)	244	1	12	2	10	Normal
		2	13	0	13	
		3	11	0	11	
	245	1	1	0	1	Partially sterile
		2	0*	0	0	
		3	7	4	3	
	251	1	0	0	0	Presumptive sterile
		2	0	0	0	
		3	0	0	0	
	253	1	16	2	14	Normal
		2	11**	1	11	
		3	-**	-	-	
	254	1	13	0	13	Normal
		2	14	0	14	
		3	11	0	11	
	256	1	-	-	-	Rebred ^b
		2	-	-	-	
		3	-	-	-	
	258	1	13	0	13	Normal
		2	0	0	0	
		3	-	-	-	
	262	1	8	3	5	Partially sterile
		2	12	8	4	
		3	10	5	5	
	264	1	9	9	0	Partially sterile
		2	11	6	5	
		3	12	3	9	
	267	1	14	1	13	Normal
		2	1	1	0	
		3	14	2	12	
	268	1	-	-	-	Rebred ^b
		2	-	-	-	
		3	-	-	-	
	269	1	6	0	6	Partially sterile
		2	7	0	7	
		3	-	-	-	
	270	1	8	0	8	Rebred ^b
		2	-	-	-	
		3	-	-	-	
	273	1	0	0	0	Rebred ^b
		2	-	-	-	
		3	-	-	-	
	280	1	11	3	8	Partially sterile
		2	11	3	8	
		3	10	5	5	

*"0" indicates a plug was observed for a female that was not pregnant.

"-" indicates a plug was not detected and the female was not pregnant.

^bSee third breeding for final classification.

Table 14 (Continued)

TRANSLOCATION STUDY OF CAPTAN
 IMPLANTATION SUMMARY OF PRESUMPTIVE P₁ MALES

	F ₁ Male Number	Female Number	TFM Group		Final Classification	
			Total Implantations	Dead Implantations		
Second breeding (cont.)	281	1	11	7	4	Partially sterile
		2	12	11	1	
		3	10	8	2	
	283	1	14	0	14	Normal
		2	0*	0	0	
		3	-**	-	-	
	286	1	-	-	-	Rebred ^b
		2	-	-	-	
		3	-	-	-	
	288	1	0	0	0	Presumptive sterile
		2	0	0	0	
		3	0	0	0	
	290	1	13	6	7	Partially sterile
		2	8	3	5	
		3	0	0	0	
	292	1	9	6	3	Partially sterile
		2	14	11	3	
		3	11	3	8	
	294	1	11	1	10	Normal
		2	12	0	12	
		3	12	1	11	
	299	1	14	5	9	Normal
		2	12	8	4	
		3	12	3	9	
	300	1	13	5	8	Partially sterile (questionable)
		2	9	4	5	
		3	12	10	2	
	302	1	13	0	13	Normal
		2	13	1	12	
		3	15	0	15	
	314	1	7	5	2	Partially sterile
		2	12	11	1	
		3	12	8	4	
	315	1	0	0	0	Partially sterile
		2	0	0	0	
		3	0	0	0	
	317	1	0	0	0	Presumptive sterile
		2	0	0	0	
		3	-	-	-	
	318	1	14	5	9	Normal
		2	9	5	4	
		3	13	2	11	
	321	1	0	0	0	Presumptive sterile
		2	0	0	0	
		3	-	-	-	

*"0" indicates a plug was observed for a female that was not pregnant.

**"- " indicates a plug was not detected and the female was not pregnant.

^b See third breeding for final classification.

Table 14 (Continued)

TRANSLOCATION STUDY OF CAPTAN
 IMPLANTATION SUMMARY OF PRESUMPTIVE F₁ MALES

	F ₁ Male Number	Female Number	TEM Group			Final Classification
			Total Implantations	Dead Implantations	Live Implantations	
Second breeding (cont.)	322	1	11	0	11	Normal
		2	11	0	11	
		3	-**	-	-	
	326	1	0*	0	0	Presumptive sterile
		2	-	-	-	
		3	-	-	-	
	327	1	13	13	0	Rebred ^b
		2	(10)†	?	(10)	
		3	-	-	-	
	333	1	12	2	10	Normal
		2	0	0	0	
		3	-	-	-	
	334	1	-	-	-	Rebred ^b
		2	-	-	-	
		3	-	-	-	
	339	1	0	0	0	Presumptive sterile
		2	0	0	0	
		3	0	0	0	
	343	1	0	0	0	Presumptive sterile
		2	0	0	0	
		3	0	0	0	
	344	1	10	5	5	Partially sterile
		2	10	6	4	
		3	11	6	5	
	345	1	6	2	4	Partially sterile
		2	12	8	4	
		3	4	1	3	
	346	1	12	0	12	Normal
		2	12	1	11	
		3	-	-	-	
	349	1	10	0	10	Normal
		2	11	3	8	
		3	12	2	10	
	350	1	8	8	0	Partially sterile
		2	0	0	0	
		3	10	9	1	
	355	1	13	0	13	Normal
		2	0	0	0	
		3	-	-	-	
	356	1	11	0	11	Normal
		2	-	-	-	
		3	(15)	?	(15)	

*"0" indicates a plug was observed for a female that was not pregnant.

"-" indicates a plug was not detected and the female was not pregnant.

†"()" indicates all implants were in early stages of development, and thus difficult to determine if they were live or dead upon gross observation.

^bSee third breeding for final classification.

Table 14 (Continued)

TRANSLOCATION STUDY OF CAPTAN
 IMPLANTATION SUMMARY OF PRESUMPTIVE F₁ MALES

	F ₁ Male Number	Female Number	Total Implantations	FEM Group		Final Classification
				Dead Implantations	Live Implantations	
Second breeding (cont.)	359	1	10	3	7	Partially sterile
		2	9	6	3	
		3	16	9	7	
	360	1	10	8	2	Partially sterile (questionable)
		2	13	13	0	
		3	**	-	-	
	361	1	13	10	3	Partially sterile
		2	13	4	7	
		3	8	8	0	
	363	1	14	2	12	Normal
		2	14	0	14	
		3	.	-	-	
	367	1	10	6	4	Normal
		2	11	6	5	
		3	11	0	11	
	369	1	-	-	-	Rebred ^b
		2	-	-	-	
		3	-	-	-	
	371	1	10	5	5	Partially sterile
		2	11	6	5	
		3	14	9	5	
	372	1	11	1	10	Normal
		2	9	0	9	
		3	-	-	-	
	375	1	5	0	5	Partially sterile
		2	-	-	-	
		3	11	4	7	
	376	1	1	0	1	Partially sterile
		2	-	-	-	
		3	4	3	1	
	381	1	11	0	11	Normal
		2	0*	0	0	
		3	-	-	-	
	382	1	-	-	-	Rebred ^b
		2	-	-	-	
		3	-	-	-	
	385	1	10	0	10	Normal
		2	12	0	12	
		3	7	0	7	
	387	1	15	0	15	Normal
		2	-	-	-	
		3	-	-	-	
	388	1	12	9	3	Partially sterile
		2	6	3	3	
		3	12	8	4	

*"0" indicates a plug was observed for a female that was not pregnant.

**"- " indicates a plug was not detected and the female was not pregnant.

^bSee third breeding for final classification.

Table 14 (Continued)

TRANSLOCATION STUDY OF CAPTAN
 IMPLANTATION SUMMARY OF PRESUMPTIVE F₁ MALES

	F ₁ Male Number	Female Number	TEM Group			Final Classification
			Total Implantations	Dead Implantations	Live Implantations	
Second breeding (concl.)	389	1	-**	-	-	Rebred ^b
		2	-	-	-	
		3	-	-	-	
	390	1	14	11	2	Partially sterile
		2	12	7	5	
		3	12	9	3	
	391	1	-	-	-	Rebred ^b
		2	-	-	-	
		3	-	-	-	
	396	1	-	-	-	Rebred ^b
		2	-	-	-	
		3	-	-	-	
	397	1	11	0	11	Normal
		2	0*	0	0	
		3	0	0	0	
399	1	14	9	5	Partially sterile (questionable)	
	2	7	5	2		
	3	4	0	4		
400	1	9	5	4	Partially sterile (questionable)	
	2	13	8	5		
	3	3	1	2		
Third breeding	216	1	9	5	4	Partially sterile (questionable)
		2	-	-	-	
		3	-	-	-	
	220	1	-	-	-	Nonbreeder
		2	-	-	-	
		3	-	-	-	
	241	1	-	-	-	Nonbreeder
		2	-	-	-	
		3	-	-	-	
	256	1	-	-	-	Presumptive sterile
		2	-	-	-	
		3	-	-	-	
	268	1	0	0	0	Presumptive sterile
		2	-	-	-	
		3	-	-	-	
	270	1	9	0	9	Normal
		2	-	-	-	
		3	-	-	-	
	273	1	-	-	-	Presumptive sterile
		2	-	-	-	
		3	-	-	-	

*"0" indicates a plug was observed for a female that was not pregnant.

**"-"- indicates a plug was not detected and the female was not pregnant.

^bSee third breeding for final classification.

Table 14 (Concluded)
 TRANSLOCATION STUDY OF CAPTAN
 IMPLANTATION SUMMARY OF PRESUMPTIVE F₁ MALES

	F ₁ Male Number	Female Number	Total Implantations	TEM Group		Final Classification
				Dead Implantations	Live Implantations	
Third breeding (concl.)	286	1	11	1	10	Normal
		2	-**	-	-	
		3	-	-	-	
	327	1	11	10	1	Partially sterile (questionable)
		2	(9)†	?	(9)	
		3	-	-	-	
	334	1	14	2	12	Normal
		2	12	0	12	
		3	-	-	-	
	369	1	-	-	-	Presumptive sterile
		2	-	-	-	
		3	-	-	-	
	382	1	11	0	11	Normal
		2	10	0	10	
		3	-	-	-	
	389	1	0*	0	0	Presumptive sterile
		2	-	-	-	
		3	-	-	-	
	391	1	-	-	-	Nonbreeder
		2	-	-	-	
		3	-	-	-	
	396	1	-	-	-	Nonbreeder
		2	-	-	-	
		3	-	-	-	

*"0" indicates a plug was observed for a female that was not pregnant.

**"- " indicates a plug was not detected and the female was not pregnant.

†"()" indicates all implants were in early stages of development, and thus difficult to determine if they were live or dead upon gross observation.

Table 15

TRANSLOCATION STUDY OF CAPTAN
IMPLANTATION SUMMARY OF PRESUMPTIVE F₁ MALES

F ₁ Male Number	Female Number	2500 ppm Group			Initial Classification
		Total Implantations	Dead Implantations	Live Implantations	
First breeding	430	1	-**	-	Nonbreeder
		2	-	-	
		3	-	-	
432	1	4	4	4	Partially sterile (questionable)
	2	(17)†	?	(17)	
	3	9	9	9	
436	1	14	5	9	Partially sterile
	2	7	2	5	
	3	11	3	8	
449	1	-	-	-	Nonbreeder
	2	-	-	-	
	3	-	-	-	
450	1	0*	0	0	Normal
	2	5	4	1	
	3	14	0	14	
451	1	11	4	7	Normal
	2	0	0	0	
	3	-	-	-	
452	1	10	1	9	Normal
	2	12	2	10	
	3	13	3	10	
455	1	-	-	-	Nonbreeder
	2	-	-	-	
	3	-	-	-	
461	1	-	-	-	Nonbreeder
	2	-	-	-	
	3	-	-	-	
469	1	-	-	-	Nonbreeder
	2	-	-	-	
	3	-	-	-	
472	1	9	1	8	Normal
	2	11	1	10	
	3	12	3	9	
474	1	-	-	-	Nonbreeder
	2	-	-	-	
	3	-	-	-	
479	1	-	-	-	Nonbreeder
	2	-	-	-	
	3	-	-	-	
480	1	(3)	?	(3)	Partially sterile
	2	0	0	0	
	3	-	-	-	
484	1	11	7	4	Normal
	2	16	2	14	
	3	11	0	11	

*"0" indicates a plug was observed for a female that was not pregnant.

**"-." indicates a plug was not detected and the female was not pregnant.

†"()" indicates all implants were in early stages of development, and thus difficult to determine if they were live or dead upon gross observation.

Table 15 (Continued)

TRANSLOCATION STUDY OF CAPTAN
IMPLANTATION SUMMARY OF PRESUMPTIVE F₁ MALES

F ₁ Male Number	Female Number	2500 ppm Group			Initial Classification
		Total Implantations	Dead Implantations	Live Implantations	
First breeding (concl.)	1	14	3	11	Normal
	2	13	1	12	
	3	8	2	6	
496	1	**	-	-	Nonbreeder
	2	-	-	-	
	3	-	-	-	
518	1	-	-	-	Nonbreeder
	2	-	-	-	
	3	-	-	-	
523	1	-	-	-	Nonbreeder
	2	-	-	-	
	3	-	-	-	
526	1	7	0	7	Partially sterile
	2	4	1	3	
	3	(1)†	?	(1)	
528	1	0*	0	0	Presumptive sterile
	2	-	-	-	
	3	-	-	-	
546	1	-	-	-	Nonbreeder
	2	-	-	-	
	3	-	-	-	
561	1	-	-	-	Nonbreeder
	2	-	-	-	
	3	-	-	-	
568	1	12	4	8	Partially sterile (questionable)
	2	(9)	?	(9)	
	3	(13)	?	(13)	
569	1	-	-	-	Nonbreeder
	2	-	-	-	
	3	-	-	-	
582	1	-	-	-	Nonbreeder
	2	-	-	-	
	3	-	-	-	
583	1	-	-	-	Nonbreeder
	2	-	-	-	
	3	-	-	-	
585	1	10	1	9	Normal
	2	11	1	10	
	3	15	3	12	
588	1	-	-	-	Partially sterile (questionable)
	2	-	-	-	
	3	(12)	?	(12)	
590	1	11	3	8	Normal
	2	0	0	0	
	3	9	0	9	

*"0" indicates a plug was observed for a female that was not pregnant.

**"." indicates a plug was not detected and the female was not pregnant.

*"()" indicates all implants were in early stages of development, and thus difficult to determine if they were live or dead upon gross observation.

Table 15 (Continued)

TRANSLOCATION STUDY OF CAPTAN
IMPLANTATION SUMMARY OF PRESUMPTIVE F₁ MALES

		2500 ppm Group				
F ₁ Male Number	Female Number	Total Implantations	Dead Implantations	Live Implantations	Final Classification	
Second breeding	430	1	12	0	12	
		2	8	0	8	Normal
		3	14	2	12	
	432	1	12	1	11	
		2	11	1	10	Normal
		3	13	1	12	
	436	1	12	1	11	
		2	12	0	12	Normal
		3	13	0	13	
	449	1	-**	-	-	
		2	-	-	-	Rebred ^b
		3	-	-	-	
	450	1	11	1	10	
		2	13	0	13	Normal
		3	14	1	13	
	451	1	0*	0	0	
		2	0	0	0	Normal
		3	11	1	10	
	452	1	14	1	13	
		2	11	0	11	Normal
		3	13	0	13	
	455	1	10	0	10	
		2	12	0	12	Normal
		3	12	1	11	
	461	1	13	0	13	
		2	11	0	11	Normal
		3	11	2	9	
	469	1	-	-	-	
		2	-	-	-	Normal
		3	12	1	11	
	472	1	0	0	0	
		2	12	1	11	Normal
		3	11	1	10	
	474	1	0	0	0	
		2	-	-	-	Rebred ^b
		3	-	-	-	
	479	1	-	-	-	
		2	-	-	-	Rebred ^b
		3	-	-	-	
	480	1	(13) [†]	?	(13)	
		2	-	-	-	Rebred ^b
		3	8	0	8	

*"0" indicates a plug was observed for a female that was not pregnant.

**"- " indicates a plug was not detected and the female was not pregnant.

[†]"(")" indicates all implants were in early stages of development, and thus difficult to determine if they were live or dead upon gross observation.

^b See third breeding for final classification.

Table 15 (Continued)

TRANSLOCATION STUDY OF CAPTAN
 IMPLANTATION SUMMARY OF PRESUMPTIVE F₁ MALES

	F ₁ Male Number	Female Number	2500 ppm Group			Final Classification
			Total Implantations	Dead Implantations	Live Implantations	
Second breeding (cont.)	484	1	11	0	11	Normal
		2	0*	0	0	
		3	14	2	12	
	489	1	9	0	9	Normal
		2	13	2	11	
		3	14	0	14	
	496	1	**	-	-	Rebred ^b
		2	-	-	-	
		3	-	-	-	
	518	1	0	0	0	Normal
		2	11	0	11	
		3	13	0	13	
	523	1	-	-	-	Rebred ^b
		2	-	-	-	
		3	-	-	-	
	526	1	0	0	0	Rebred ^b
		2	5	0	5	
		3	0	0	0	
	528	1	15	1	14	Normal
		2	-	-	-	
		3	-	-	-	
	546	1	-	-	-	Rebred ^b
		2	-	-	-	
		3	-	-	-	
	561	1	-	-	-	Rebred ^b
		2	-	-	-	
		3	-	-	-	
	568	1	11	2	9	Normal
		2	11	0	11	
		3	8	0	8	
	569	1	11	0	11	Normal
		2	-	-	-	
		3	10	0	10	
	582	1	7	0	7	Normal
		2	15	2	13	
		3	10	2	8	
	583	1	-	-	-	Rebred ^b
		2	-	-	-	
		3	-	-	-	
	585	1	12	0	12	Normal
		2	13	0	13	
		3	13	0	13	
	588	1	9	0	9	Rebred ^b
		2	-	-	-	
		3	-	-	-	

*"0" indicates a plug was observed for a female that was not pregnant.

**"- " indicates a plug was not detected and the female was not pregnant.

^bSee third breeding for final classification.

Table 15 (Concluded)

TRANSLOCATION STUDY OF CAPTAN
 IMPLANTATION SUMMARY OF PRESUMPTIVE F₁ MALES

	2500 ppm Group					Final Classification
	F ₁ Male Number	Female Number	Total Implantations	Dead Implantations	Live Implantations	
Second breeding (concl.)	590	1	0*	0	0	Normal
		2	13	2	11	
		3	12	0	12	
Third breeding	449	1	-**	-	-	Nonbreeder
		2	-	-	-	
		3	-	-	-	
474		1	-	-	-	Presumptive sterile
		2	-	-	-	
		3	-	-	-	
479		1	-	-	-	Nonbreeder
		2	-	-	-	
		3	-	-	-	
480		1	10	8	2	Partially sterile (questionable)
		2	-	-	-	
		3	-	-	-	
496		1	-	-	-	Nonbreeder
		2	-	-	-	
		3	-	-	-	
523		1	-	-	-	Nonbreeder
		2	-	-	-	
		3	-	-	-	
526		1	2	0	2	Partially sterile
		2	1	0	1	
		3	0	0	0	
546		1	9	0	9	Normal
		2	13	0	13	
		3	-	-	-	
561		1	15	0	15	Normal
		2	11	0	11	
		3	-	-	-	
583		1	-	-	-	Nonbreeder
		2	-	-	-	
		3	-	-	-	
588		1	12	1	11	Normal
		2	-	-	-	
		3	-	-	-	

*"0" indicates a plug was observed for a female that was not pregnant.

**"- " indicates a plug was not detected and the female was not pregnant.

Table 16

TRANSLOCATION STUDY OF CAPTAN
 IMPLANTATION SUMMARY OF PRESUMPTIVE F₁ MALES

	F ₁ Male Number	Female Number	5000 ppm Group			Initial Classification
			Total Implantations	Dead Implantations	Live Implantations	
First breeding	601	1	-**	-	-	Partially sterile (questionable)
		2	(11) [†]	?	(11)	
		3	(8)	?	(8)	
	604	1	7	0	7	Normal
		2	11	2	9	
		3	14	5	9	
	620	1	8	0	8	Normal
		2	-	-	-	
		3	-	-	-	
	622	1	-	-	-	Partially sterile (questionable)
		2	-	-	-	
		3	(9)	?	(9)	
	626	1	-	-	-	Nonbreeder
		2	-	-	-	
		3	-	-	-	
	627	1	9	1	8	Normal
		2	11	3	8	
		3	8	0	8	
	634	1	-	-	-	Nonbreeder
		2	-	-	-	
		3	-	-	-	
	635	1	-	-	-	Nonbreeder
		2	-	-	-	
		3	-	-	-	
	638	1	10	0	10	Normal
		2	6	0	6	
		3	6	0	6	
	646	1	12	2	10	Normal
		2	-	-	-	
		3	-	-	-	
	653	1	0*	0	0	Partially sterile (questionable)
		2	(13)	?	(13)	
		3	-	-	-	
	657	1	9	7	2	Partially sterile
		2	13	8	5	
		3	8	7	1	
	660	1	0	0	0	Normal
		2	11	1	10	
		3	8	0	8	
	668	1	-	-	-	Partially sterile
		2	-	-	-	
		3	7	1	6	

*"0" indicates a plug was observed for a female that was not pregnant.

**"- " indicates a plug was not detected and the female was not pregnant.

[†]"(")" indicates all implants were in early stages of development, and thus difficult to determine if they were live or dead upon gross observation.

Table 16 (Continued)

TRANSLOCATION STUDY OF CAPTAN
 IMPLANTATION SUMMARY OF PRESUMPTIVE F₁ MALES

	F ₁ Male Number	Female Number	5000 ppm Group			Initial Classification
			Total Implantations	Dead Implantations	Live Implantations	
First breeding (cont.)	671	1	-**	-	-	Nonbreeder
		2	-	-	-	
		3	-	-	-	
	672	1	-	-	-	Nonbreeder
		2	-	-	-	
		3	-	-	-	
	679	1	6	0	6	Partially sterile
		2	0*	0	0	
		3	12	4	8	
	722	1	-	-	-	Nonbreeder
		2	-	-	-	
		3	-	-	-	
	732	1	4	0	4	Partially sterile
		2	-	-	-	
		3	-	-	-	
	733	1	-	-	-	Nonbreeder
		2	-	-	-	
		3	-	-	-	
	734	1	-	-	-	Nonbreeder
		2	-	-	-	
		3	-	-	-	
	736	1	-	-	-	Nonbreeder
		2	-	-	-	
		3	-	-	-	
	743	1	0	0	0	Partially sterile (questionable)
		2	-	-	-	
		3	(15)†	?	(15)	
	760	1	8	1	7	Partially sterile
		2	5	0	5	
		3	9	1	8	
	761	1	12	3	9	Normal
		2	-	-	-	
		3	-	-	-	
	765	1	12	3	9	Normal
		2	11	1	10	
		3	-	-	-	
	767	1	12	2	10	Normal
		2	-	-	-	
		3	-	-	-	
	769	1	8	8	8	Partially sterile (questionable)
		2	-	-	-	
		3	(12)	?	(12)	

*"0" indicates a plug was observed for a female that was not pregnant.

*"-." indicates a plug was not detected and the female was not pregnant.

*"()" indicates all implants were in early stages of development, and thus difficult to determine if they were live or dead upon gross observation.

Table 16 (Continued)

TRANSLOCATION STUDY OF CAPIAN
IMPLANTATION SUMMARY OF PRESUMPTIVE F₁ MALES

	F ₁ Male Number	Female Number	5000 ppm Group			Initial Classification
			Total Implantations	Dead Implantations	Live Implantations	
First breeding (concl.)	779	1	2	0	2	Partially sterile
		2	0*	0	0	
		3	1	0	1	
	780	1	4	0	4	Partially sterile
		2	1	0	1	
		3	0	0	5	
						<u>Final Classification</u>
Second breeding	601	1	12	1	11	Normal
		2	0	0	0	
		3	- [†]	-	-	
	604	1	9	0	9	Normal
		2	10	0	10	
		3	15	1	14	
	620	1	10	1	9	Normal
		2	-	-	-	
		3	-	-	-	
	622	1	0	0	0	Normal
		2	12	0	12	
		3	-	-	-	
	626	1	(12) [†]	?	(12)	Normal
		2	12	0	12	
		3	-	-	-	
	627	1	13	3	10	Normal
		2	6	0	6	
		3	1	1	0	
	634	1	-	-	-	Rebred ^b
		2	-	-	-	
		3	-	-	-	
	635	1	-	-	-	Rebred ^b
		2	1	1	0	
		3	-	-	-	
	638	1	10	0	10	Normal
		2	12	0	12	
		3	11	0	11	
	646	1	11	0	11	Normal
		2	13	0	13	
		3	10	1	9	
653	1	14	0	14	Normal	
	2	13	0	13		
	3	10	0	10		

*"0" indicates a plug was observed for a female that was not pregnant.

†"-" indicates a plug was not detected and the female was not pregnant.

†"()" indicates all implants were in early stages of development, and thus difficult to determine if they were live or dead upon gross observation.

^bSee third breeding for final classification.

Table 16 (Continued)

TRANSLOCATION STUDY OF CAPTAN
 IMPLANTATION SUMMARY OF PRESUMPTIVE F₁ MALES

F ₁ Male Number	Female Number	5000 ppm Group			Final Classification	
		Total Implantations	Dead Implantations	Live Implantations		
Second breeding (cont.)	657	1	7	5	2	Partially sterile
		2	9	8	1	
		3	5	1	4	
660	1	14	0	14	Normal	
	2	10	1	9		
	3	12	0	12		
668	1	9	0	9	Normal	
	2	13	1	12		
	3	13	1	12		
671	1	14	0	14	Normal	
	2	11	0	11		
	3	-**	-	-		
672	1	-	-	-	Normal	
	2	-	-	-		
	3	14	0	14		
679	1	7	0	7	Normal	
	2	12	0	12		
	3	4	1	3		
722	1	-	-	-	Rebred ^b	
	2	-	-	-		
	3	-	-	-		
732	1	12	0	12	Normal	
	2	-	-	-		
	3	-	-	-		
733	1	-	-	-	Rebred ^b	
	2	-	-	-		
	3	-	-	-		
734	1	11	0	11	Normal	
	2	12	0	12		
	3	-	-	-		
736	1	-	-	-	Rebred ^b	
	2	-	-	-		
	3	-	-	-		
743	1	14	0	14	Normal	
	2	10	0	10		
	3	11	1	10		
760	1	1	0	1	Partially sterile	
	2	0*	0	0		
	3	0	0	0		
761	1	5	0	5	Normal	
	2	14	1	13		
	3	12	1	11		
765	1	15	3	12	Normal	
	2	11	0	11		
	3	9	0	9		

*"0" indicates a plug was observed for a female that was not pregnant.

**"-." indicates a plug was not detected and the female was not pregnant.

^bSee third breeding for final classification.

Table 16 (Concluded)

TRANSLOCATION STUDY OF CARTAN
IMPLANTATION SUMMARY OF PRESUMPTIVE F₁ MALES

F ₁ Male Number	Female Number	Total Implantations	5000 ppm Group		Final Classification	
			Dead Implantations	Live Implantations		
Second breeding (concl.)	767	1	14	2	12	Normal
		2	13	0	13	
		3	12	0	12	
	769	1	12	0	12	Normal
		2	13	0	13	
		3	10	0	10	
	779	1	0 [*]	0	0	Partially sterile
		2	4	0	4	
		3	-**	-	-	
780	1	0	0	0	Partially sterile	
	2	2	0	2		
	3	5	1	4		
Third breeding	634	1	-	-	-	Nonbreeder
		2	-	-	-	
		3	-	-	-	
	635	1	-	-	-	Partially sterile
		2	-	-	-	
		3	-	-	-	
	722	1	13	1	12	Normal
		2	-	-	-	
		3	-	-	-	
	733	1	-	-	-	Nonbreeder
		2	-	-	-	
		3	-	-	-	
	736	1	0	0	0	Presumptive sterile
		2	-	-	-	
		3	-	-	-	

*"0" indicates a plug was observed for a female that was not pregnant.

"-" indicates a plug was not detected and the female was not pregnant.

Table 17
 TRANSLOCATION STUDY OF CAPTAN
 IMPLANTATION SUMMARY OF PRESUMPTIVE F₁ MALES

		5000 ^a ppm Group				
F ₁ Male Number	Female Number	Total Implantations	Dead Implantations	Live Implantations	Initial Classification	
First breeding	805	1	- ^{**}	-	-	Nonbreeder
		2	-	-	-	
		3	-	-	-	
	806	1	-	-	-	Nonbreeder
		2	-	-	-	
		3	-	-	-	
	815	1	6	0	6	Partially sterile
		2	9	0	9	
		3	7	0	7	
	816	1	-	-	-	Partially sterile (questionable)
		2	(10) [†]	?	(10)	
		3	(11)	?	(11)	
	817	1	-	-	-	Nonbreeder
		2	-	-	-	
		3	-	-	-	
	818	1	10	0	10	Normal
		2	6	0	6	
		3	-	-	-	
822	1	-	-	-	Nonbreeder	
	2	-	-	-		
	3	-	-	-		
823	1	-	-	-	Nonbreeder	
	2	-	-	-		
	3	-	-	-		
837	1	-	-	-	Nonbreeder	
	2	-	-	-		
	3	-	-	-		
841	1	-	-	-	Nonbreeder	
	2	-	-	-		
	3	-	-	-		
846	1	5	0	5	Partially sterile	
	2	-	-	-		
	3	-	-	-		
847	1	(14)	?	(14)	Partially sterile (questionable)	
	2	-	-	-		
	3	-	-	-		
					<u>Final Classification</u>	
Second breeding	805	1	-	-	Rebred ^b	
		2	-	-		
		3	2	0		2

^aIndicates traumatized F₀ females.

^{*}"0" indicates a plug was observed for a female that was not pregnant.

^{**}"-" indicates a plug was not detected and the female was not pregnant.

[†]"()" indicates all implants were in early stages of development, and thus difficult to determine if they were live or dead upon gross observation.

^bSee third breeding for final classification.

Table 17 (Continued)

TRANSLOCATION STUDY OF CAPLAN
IMPLANTATION SUMMARY OF PRESUMPTIVE P₁ MALES

F ₁ Male Number	Female Number	Total Implantations	5000 ^a ppm Group		Final Classification	
			Dead Implantations	Live Implantations		
Second breeding (concl.)	806	1	13	1	12	Normal
		2	14	0	14	
		3	10	0	10	
	815	1	14	1	13	Normal
		2	11	0	11	
		3	13	3	10	
	816	1	11	2	9	Normal
		2	**	-	-	
		3	-	-	-	
	817	1	11	4	7	Rebred ^b
		2	0*	0	0	
		3	-	-	-	
	818	1	11	0	11	Normal
		2	8	0	8	
		3	13	1	12	
	822	1	6	0	6	Rebred ^b
		2	-	-	-	
		3	-	-	-	
	823	1	13	0	13	Normal
		2	-	-	-	
		3	-	-	-	
	837	1	8	1	7	Rebred ^b
		2	-	-	-	
		3	-	-	-	
	841	1	15	2	13	Normal
		2	10	0	10	
		3	-	-	-	
	846	1	10	0	10	Normal
		2	12	0	12	
		3	12	0	12	
	847	1	11	1	10	Normal
		2	10	0	10	
		3	15	0	14	
Third breeding	805	1	-	-	-	Partially sterile
		2	-	-	-	
		3	-	-	-	
	817	1	13	2	11	Normal
		2	9	7	2	
		3	13	0	13	
	822	1	10	0	10	Normal
		2	12	0	12	
		3	12	0	12	

^aIndicates traumatized F₀ females.

*"0" indicates a plug was observed for a female that was not pregnant.

**"-." indicates a plug was not detected and the female was not pregnant.

^bSee third breeding for final classification.

Table 17 (Concluded)

TRANSLOCATION STUDY OF CAPTAN
 IMPLANTATION SUMMARY OF PRESUMPTIVE F₁ MALES

F ₁ Male Number	Female Number	Total Implantations	5000 ^a ppm Group		Final Classification
			Dead Implantations	Live Implantations	
Third breeding (concl.)	1 2 3	-** - -	- - -	- - -	Normal

^aIndicates traumatized F₀ females.

**"- " indicates a plug was not detected and the female was not pregnant.

SHIPPED FROM NIS

APR 15 1980

