

A

Origin and Framework of the Development of Dietary Reference Intakes

This report is one of a series of publications resulting from the comprehensive effort being undertaken by the Food and Nutrition Board's (FNB) Standing Committee on the Scientific Evaluation of Dietary Reference Intakes (DRI Committee) and its panels and subcommittees.

ORIGIN

This initiative began in June 1993, when FNB organized a symposium and public hearing entitled, "Should the Recommended Dietary Allowances Be Revised?" Shortly thereafter, to continue its collaboration with the larger nutrition community on the future of the Recommended Dietary Allowances (RDAs), FNB took two major steps: (1) it prepared, published, and disseminated the concept paper, "How Should the Recommended Dietary Allowances Be Revised?" (IOM, 1994), which invited comments regarding the proposed concept, and (2) it held several symposia at nutrition-focused professional meetings to discuss FNB's tentative plans and to receive responses to the initial concept paper. Many aspects of the conceptual framework of the DRIs came from the United Kingdom's report, *Dietary Reference Values for Food Energy and Nutrients for the United Kingdom* (COMA, 1991).

The five general conclusions presented in FNB's 1994 concept paper were:

1. Sufficient new information has accumulated to support a reassessment of the RDAs.
2. Where sufficient data for efficacy and safety exist, reduction in the risk of chronic degenerative disease is a concept that should be included in the formulation of future recommendations.
3. Upper levels of intake should be established where data exist regarding risk of toxicity.
4. Components of food that may benefit health, although not meeting the traditional concept of a nutrient, should be reviewed, and if adequate data exist, reference intakes should be established.
5. Serious consideration must be given to developing a new format for presenting future recommendations.

Subsequent to the symposium and the release of the concept paper, FNB held workshops at which invited experts discussed many issues related to the development of nutrient-based reference values. (FNB and DRI Committee members have continued to provide updates and engage in discussions at professional meetings.) In addition, FNB gave attention to the international uses of the earlier RDAs and the expectation that the scientific review of nutrient requirements should be similar for comparable populations.

Concurrently, Health Canada and Canadian scientists were reviewing the need for revision of the *Recommended Nutrient Intakes* (RNIs) (Health Canada, 1990b). Consensus following a symposium for Canadian scientists, cosponsored by the Canadian National Institute of Nutrition and Health Canada in April 1995, was that the Canadian government should pursue the extent to which involvement with the developing FNB process would benefit both Canada and the United States in leading toward harmonization.

Based on extensive input and deliberations, FNB initiated action to provide a framework for the development and possible international harmonization of nutrient-based recommendations that would serve, where warranted, for all of North America. To this end, in December 1995, FNB began a close collaboration with the government of Canada and took action to establish the DRI Committee. It is hoped that representatives from Mexico will join in future deliberations.

THE CHARGE TO THE COMMITTEE

In 1995, the DRI Committee was appointed to oversee and conduct this project. It devised a plan involving the work of seven or more expert nutrient group panels and two overarching subcommittees (Figure A-1).

The Subcommittee on Interpretation and Uses of Dietary Reference Intakes (Uses Subcommittee), composed of experts in nutrition, dietetics, statistics, nutritional epidemiology, public health, economics, and consumer perspectives, was to (1) review the scientific literature regarding the uses of dietary reference standards and their applications, (2) provide guidance for the appropriate application of DRIs for specific purposes and identify inappropriate applications, (3) provide guidance for adjustments to be made for potential errors in dietary intake data and the assumptions regarding intake and requirement distributions, (4) provide specific guidance for use of DRI values for individual nutrients, and (5) identify research needed to improve the statistical underpinnings regarding quantitative applications of the DRIs for assessing and planning diets for individuals and for groups.




FIGURE A-1 Dietary Reference Intakes project structure.

This second report from the Uses Subcommittee examines the appropriate uses of each of the DRI values in *planning* nutrient intakes of groups and of individuals; an earlier report presented information on the appropriate uses of specific DRI values in *assessing* diets for groups and for individuals (IOM, 2000a). Each report presents the statistical underpinnings for the various uses of the DRI values and also indicates when specific uses are inappropriate. This report reflects the work of the DRI Committee, the Uses Subcommittee, and the Subcommittee on Upper Reference Levels of Nutrients.

ISSUES OF RELEVANCE FROM PAST DIETARY REFERENCE INTAKE REPORTS

Methodology to Develop Estimated Average Requirements and Recommended Dietary Allowances When Requirements for Nutrients Are Not Normally Distributed

For most of the nutrients for which Estimated Average Requirements (EARs) have been established, the required assumption of distribution of requirements is that of symmetry about the mean. In the case of iron, a nutrient of concern in many subgroups in the population in the United States, Canada, and other areas, requirements are known to follow a nonnormal distribution (IOM, 2001). Thus, a different method was needed to determine the intake of iron at which half of the individuals would be expected to be inadequate in the criterion used to establish adequacy (the EAR), and also to construct an intake level at which only a small percentage of the population would be inadequate (the RDA). Similar adjustments were made for dietary protein (IOM, 2002a).

If the requirement of a nutrient is not normally distributed but can be transformed to normality, its EAR and RDA can be estimated by transforming the data, calculating the 50th and 97.5th percentiles, and transforming these percentiles back into the original units. In this case, the difference between the EAR and the RDA cannot be used to obtain an estimate of the standard deviation or the coefficient of variation because skewing is usually present.

Where factorial modeling is used to estimate the distribution of requirements from the distributions of the individual components of requirement, as was done in the case of iron recommendations (IOM, 2001), it is necessary to add the individual distributions (convolutions). This is easy to do given that the average requirement is simply the sum of the averages of the individual component distri-

butions, and a standard deviation of the combined distribution can be estimated by standard statistical techniques. The 97.5th percentile can then be estimated (for a further elaboration of this method, see Chapter 9 and Appendix I of *Dietary Reference Intakes for Vitamin A, Vitamin K, Arsenic, Boron, Chromium, Copper, Iodine, Iron, Manganese, Molybdenum, Nickel, Silicon, Vanadium, and Zinc* [IOM, 2001]).

If normality cannot be assumed for all of the components of requirement, then Monte Carlo simulation is used for the summation of the components. This approach models the distributions of the individual components and randomly assigns values to a large simulated population. The total requirement is then calculated for each individual and the median and the 97.5th percentile are calculated directly. As was the case for iron (IOM, 2001), the underlying joint distribution is approximated and a large number of individuals (100,000) are randomly generated. Information about the distribution of values for the requirement components is modeled on the basis of known physiology. Monte Carlo approaches may be used in the simulation of the distribution of components, or where large data sets exist for similar populations (data sets such as growth rates in infants), estimates of relative variability may be transferred to the component in the simulated population (Gentle, 1998). At each step the goal is to achieve distribution values for the component that not only reflect known physiology or known direct observations, but also can be transformed into a distribution that can be modeled and used in selecting random members to contribute to the final requirement distribution. When the final distribution representing the convolution of components has been derived, the median and 97.5th percentiles of the distribution can be directly estimated. It is recognized that in its simplest form, the Monte Carlo approach ignores possible correlation among components. In the case of iron, however, expected correlation is built into the modeling of requirement where components are linked to a common variable (e.g., growth rate) so that not all sources of correlation are neglected.

Life Stage Groups

Nutrient intake recommendations are expressed for 22 life stage groups, as listed in Table A-1 and described in more detail elsewhere (IOM, 1997). If data are too sparse to distinguish differences in requirements by life stage and gender, the analysis may be presented for a larger grouping. Differences are indicated by gender when warranted by the data.

TABLE A-1 The 22 Life Stage Groups for Which Dietary Reference Intakes (DRIs) are Given

Life Stage Groups			
Infants	Males	Females	Pregnancy
0-6 mo	9-13 y	9-13 y	18 y
7-12 mo	14-18 y	14-18 y	19-30 y
	19-30 y	19-30 y	31-50 y
Children	31-50 y	31-50 y	
1-3 y	51-70 y	51-70 y	Lactation
4-8 y	> 70 y	> 70 y	18 y
			19-30 y
			31-50 y

NOTE: Differences in DRIs are indicated by gender when warranted by the data.

Reference Heights and Weights Used in Extrapolating Dietary Reference Intakes for Vitamins and Elements

The most up-to-date data providing heights and weights of individuals in the United States and Canada when the DRI process was initiated in 1995 were anthropometric data from the 1988-1994 Third National Health and Nutrition Examination Survey (NHANES III) in the United States, and older data from Canada. Reference values derived from the NHANES III data and used in previous reports are given in Table A-2. The earlier values were obtained as follows: the median heights for the life stage and gender groups through age 30 years were identified, and the median weights for these heights were based on reported median Body Mass Index (BMI) for the same individuals. Since there is no evidence that weight should change as adults age if activity is maintained, the reference weights for adults aged 19 through 30 years were applied to all adult age groups.

The most recent nationally representative data available for Canadians (from the 1970-1972 Nutrition Canada Survey [Demirjian, 1980]) were also reviewed. In general, median heights of children from 1 year of age in the United States were greater by 3 to 8 cm (1 to 2.5 in) than those of children of the same age in Canada measured two decades earlier (Demirjian, 1980). This difference could be partly explained by approximations necessary to compare the two data sets, but more likely by a continuation of the secular

TABLE A-2 Reference Heights and Weights for Children and Adults in the United States Used in the Vitamin and Element Dietary Reference Intake Reports^a through 2001

Sex	Age	Median Body Mass Index (kg/m ²)	Reference Height, cm (in)	Reference Weight ^b kg (lb)
Male, female	2–6 mo	—	64 (25)	7 (16)
	7–12 mo	—	72 (28)	9 (20)
	1–3 y	—	91 (36)	13 (29)
Male	4–8 y	15.8	118 (46)	22 (48)
	9–13 y	18.5	147 (58)	40 (88)
	14–18 y	21.3	174 (68)	64 (142)
	19–30 y	24.4	176 (69)	76 (166)
Female	9–13 y	18.3	148 (58)	40 (88)
	14–18 y	21.3	163 (64)	57 (125)
	19–30 y	22.8	163 (64)	61 (133)

^a IOM (1997, 1998a, 2000b, 2001). Adapted from the Third National Health and Nutrition Examination Survey, 1988–1994.

^b Calculated from body mass index and height for ages 4 through 8 years and older.

trend of increased heights for age noted in the Nutrition Canada Survey when it compared data from that survey with an earlier (1953) national Canadian survey (Pett and Ogilvie, 1956).

Similarly, median weights beyond age 1 year derived from the recent survey in the United States (NHANES III) were also greater than those obtained from the older Canadian survey (Demirjian, 1980). Differences were greatest during adolescence, ranging from 10 to 17 percent higher. The differences probably reflect the secular trend of earlier onset of puberty (Herman-Giddens et al., 1997), rather than differences in populations. Calculations of BMI for young adults (e.g., a median of 22.6 for Canadian women compared to 22.8 for U.S. women) resulted in similar values, thus indicating greater concordance between the two surveys by adulthood.

The reference weights used in the previous DRI reports (IOM, 1997, 1998a, 2000a, 2000b, 2001) were thus based on the most recent data set available from either country, with recognition that earlier surveys in Canada indicated shorter stature and lower weights during adolescence than did surveys in the United States.

New Reference Heights and Weights

As discussed earlier, when the DRI process was undertaken in 1994, the reference heights and weights used were developed based on NHANES III data on BMI for children and young adults (IOM, 1997). Given the increasing prevalence of overweight and obesity in both adults and children (HHS, 1996), use of such population data is of concern. However, recent data providing heights and ideal BMIs for adults (Kuczmarski et al., 2000) and new growth charts for infants and children have allowed the development of new reference heights and weights (Table A-3) that should more closely approximate ideal weights based on low risk of chronic disease and adequate growth for children. These new values were used in the DRI report published in 2002 (IOM, 2002a) and will be used in subsequent DRI reports until they need to be revised based on new data or because of a conceptual need.

TABLE A-3 New Reference Heights and Weights for Children and Adults in the United States

Sex	Age	Previous	New	New	New
		Median Body Mass Index ^a (BMI) (kg/m ²)			
Male, female	2-6 mo	—	—	62 (24)	6 (13)
	7-12 mo	—	—	71 (28)	9 (20)
	1-3 y	—	—	86 (34)	12 (27)
Male	4-8 y	15.8	15.3	115 (45)	20 (44)
	9-13 y	18.5	17.2	144 (57)	36 (79)
	14-18 y	21.3	20.5	174 (68)	61 (134)
	19-30 y	24.4	22.5	177 (70)	70 (154)
Female	9-13 y	18.3	17.4	144 (57)	37 (81)
	14-18 y	21.3	20.4	163 (64)	54 (119)
	19-30 y	22.8	21.5	163 (64)	57 (126)

^a Taken from male and female median BMI and height-for-age data from the Third National Health and Nutrition Examination Survey (NHANES III), 1988-1994; used in earlier DRI reports (IOM 1997, 1998a, 2000b, 2001).

^b Taken from new data on male and female median BMI and height-for-age data from the Centers for Disease Control and Prevention/National Center for Health Statistics Growth Charts (CDC/NCHS, 2000; Kuczmarski et al., 2000).

^c Calculated from CDC/NCHS Growth Charts (CDC/NCHS, 2000; Kuczmarski et al., 2000), median BMI and median height for ages 4 through 19 years.