

Index

A

Acceptable Macronutrient Range, 30, 78, 80

Adequacy of nutrient intake. *See also*

Prevalence of inadequate intakes
assessment of, 95-96, 196; *see also*

Dietary assessment
confidence levels, 84, 94

criteria of, 23, 24, 28, 29, 30, 83, 102, 166

energy intake and, 31-32, 91

in group-level planning, 29, 75, 81, 83, 91, 95-96, 114, 115, 184, 185

in individual-level planning, 26, 37, 45-47, 50, 53

observed difference and, 37

probability of correct conclusion
about, 37, 81

quantitative assessments of, 37

risk-reduction based indicator of, 22, 23

uncertainty in, 29

Adequate Intakes (AIs). *See also specific nutrients*

applicable population, 25

context for use, 4, 23, 39

Daily Values on food labels compared, 52

defined, 3, 24-25, 39

derivation of, 22-23, 25, 39, 83

EARs compared, 4, 25, 154

and food guides, 47, 49

in group-level planning, 9, 11-12, 83-84, 85-87, 94-95, 103

and group mean intake, 83, 84, 85-87

in individual-level planning, 4, 6, 35, 37, 39, 41

menu planning with, 11-12, 83-84, 85-87

and nutrient density approaches, 94, 103

nutrients, by life-stage group, 85-87

and prevalence of inadequate intakes, 84, 94

RDAs compared, 4, 25

replacement with EARs and RDAs, 25, 154

research recommendations, 25, 154

uncertainty in, 4, 39, 133-134

uses, 2, 4, 25

Adjusting intake distributions

backtransformation of adjusted data, 166, 200-201, 203

defining groups for, 204

EAR and, 65-68

EAR cut-point method and, 10, 109, 201

energy, 33, 205

estimation of within- and between-person variance, 33, 63-64, 199-200, 204, 207-208

incomplete data, 202-203
 individual subjects' means, 200
 Iowa State University method, 64, 109, 117, 196-197, 202, 203-204, 205, 206
 limitations of methods, 205-206
 macronutrient composition of diets and, 205
 NRC method, 64, 109, 196, 197-203, 205, 206, 207-208
 ratios, 184, 205
 residual method of energy adjustment, 33
 sample size and, 64, 197, 202-203, 204
 SAS program, 198, 207-208
 Shapiro-Wilk statistic, 198
 SIDE package, 293
 skewed data, 166-167, 184, 188, 201-202
 special considerations, 57, 201-203
 systematic effects on within-person variation and, 33, 204
 transformation of data, 198-199, 200-201, 202, 203
 underlying concepts and assumptions, 197-198, 205-206
 underreporting of intakes and, 32-34, 206
 weighting factors, 99, 101, 204
 zinc example, 70, 198-201

Adolescent boys
 energy intakes, 185
 vitamin A, 130, 131
 vitamin C, 117, 119, 120, 122, 185, 187

Adolescent girls
 iron, 142
 zinc, 10, 60, 62, 70, 95, 123-126

Adults over age 50, physiological changes affecting nutrient requirements, 64, 135, 143-144

Alcohol, 54, 80, 144, 178

Amino acids, indispensable, 137-138

Assisted-living setting, 56, 81, 83, 108-113

Asymmetrical distribution. *See* Skewed distribution

Athletes, 36, 140-141

B

B vitamins, 140
 Basal metabolic rate, 31, 32

Basal requirement, defined, 28
 Beta-carotene, 47, 152
 Beta-carotene:retinol equivalency ratio, 48
 Between-person variance, 63, 64, 75, 98-99, 188
 Bias. *See also* Error sources; Measurement error
 in cut-point method, 74, 75
 in distribution of intakes, 74
 in energy intakes, 33, 92, 105
 in usual nutrient intakes, 92, 105
 Bioavailability of nutrients, 15, 17, 29, 50, 134-135, 136, 137, 143-144, 145
 Biotin, 52, 85
 Body mass index, 24, 41, 105, 168, 169
 Body weight. *See* Weight
 Boston Nutritional Status Survey, 110-111

C

Calcium, 25, 39, 49, 52, 85, 135, 136, 194

Canada
 Consumer Packaging and Labeling Act, 178
 Food and Drugs Act, 178
 Food and Drugs Regulations, 178
Food Guide to Healthy Eating, 7, 12, 35, 44, 45, 46, 172, 173
 Food Inspection Agency, 178
 food labels, 51, 54, 177-179
 fortification of foods, 50, 126, 192, 194
 Guide to Food Labeling and Advertising, 177-178
 Health Canada, 164
 Nutrition Canada Survey, 168-169
 Nutrition Recommendations, 35, 51, 53, 181-182
 Vitality program, 173

Canadian National Institute of Nutrition, 164

Canadian Recommended Nutrient Intakes (RNI), 2, 7, 19, 20, 21, 22, 44-45, 164, 178, 181

Carbohydrate. *See also* Macronutrient distribution
 Acceptable Macronutrient Range, 78
 actual vs. recommended intakes, 49
 usual energy intakes from, 78-79, 80

Carotenoids, 47, 48, 54

Child and Adult Care Feeding Program, 113

Children, ages 1 through 8 years
 EER, 4
 fortified foods, 50, 194
 nutrition programs, 58, 90, 113-116
 physiological considerations in
 planning for, 141
 Choline, 25, 52, 85
 Chromium, 52, 86, 176
 Chronic disease risk reduction
 food health claims, 174, 178-179
 as indicator of nutrient adequacy, 22,
 164, 180
 Cigarette smoking, 144
 Coefficient of variation (CV)
 EAR and RDA, 24, 65
 of nutrient requirement, 166, 184
 Continuing Survey of Food Intakes by
 Individuals (CSFII), 40, 79, 109,
 110-111, 114, 117, 128, 129, 138,
 185
 Copper, 49, 52, 86, 136, 184
 Criterion of nutritional adequacy, 23, 24,
 28, 29, 30, 83, 102, 166
 Cut-point method
 accuracy, 71
 and adjustment of intake distributions,
 10, 109, 201
 assumptions in, 59, 62, 73
 bias in, 74, 75
 correlated intakes and requirements
 and, 17, 41, 61, 75-76, 78
 defined, 59
 distribution of usual intakes, 57, 70,
 72, 96-97
 EAR and, 9-10, 17, 55, 59, 60, 61, 62,
 68-73, 74-76, 94, 96-97, 109, 110-
 111, 123, 154
 prevalence of inadequate intakes, 9-10,
 17, 55, 59, 60, 61, 62, 68-73, 75-76,
 94, 96-97, 109, 110-111, 123, 154
 probability approach compared, 17,
 59, 73-76
 requirement distribution and, 17, 59
 requirement variance relative to intake
 variance, 59
 Cysteine, 137, 138

D

Daily Reference Values, 174-175
 Daily Value (DV), 51, 52, 174, 175-176

Data and database issues. *See also*
 Adjusting intake distributions
 availability of usual intake data, 108-
 109, 149
 food composition databases, 7, 17, 27,
 29, 30, 132, 151, 152, 206
 quality of dietary intake data, 15-16,
 17, 63, 150-152
 research recommendations, 149, 150-
 152
 self reports of dietary intakes, 15-16,
 27, 29, 30-34, 63, 64-65, 104, 151
 supplement intakes, 151
 uncertainty in nutrient requirements,
 27, 29-30
 Defined nutritional states, 83
 Density estimation. *See* Nutrient density
 approaches; Nutrient density
 distribution approach; Simple
 nutrient density approach
 Dental caries, 192
 Diet histories, 31, 37
 Dietary assessment
 applications, 1, 20-21
 dietary planning compared, 1, 20-21,
 22, 27, 62
 error sources, 64, 202
 generalizability of results, 32-33
 individual-level, 37
 Dietary guidance systems, 44
 Dietary guidelines, 2, 7, 35, 51, 53-54, 179-
 182
Dietary Guidelines for Americans, 35, 171,
 180-181
 Dietary intake data. *See also* Data and
 database issues; Dietary survey data;
 Measuring dietary intakes;
 Observed intakes; Usual nutrient
 intakes
 adjustment of, *see* Adjusting intake
 distributions
 collection, 31, 33, 99, 196, 205
 error sources in, 29, 30-34, 37, 64, 70,
 76, 196, 197
 fortified foods, 129, 193-194
 for group-level planning, 129
 habitual intakes, 195
 individual variations in, 16, 33, 63
 quality of, 15-16, 17, 63, 149, 150-152,
 153
 research recommendations, 17, 149,
 150-152, 153

- self reports, 15-16, 27, 29, 30-34, 63, 64-65, 104, 151
- supplements, 17, 151
- from surveys, 9, 10, 13, 34, 64, 84, 205-206
- uncertainty in estimates, 27, 30-34, 37
- underreporting, 15-16, 17, 27, 29, 30-34, 63, 64-65, 104, 206
- Dietary planning. *See also* Group-level planning; Individual-level planning
 - acceptably low probability of nutrient inadequacy and, 26-27
 - activities most relevant for, 2, 21
 - age and physiological considerations, 27, 140, 141-144, 167
 - bioavailability considerations, 15, 17, 29, 50, 134-135, 136, 137, 143-144, 145
 - body size or composition considerations, 140
 - caveats, 27-34, 108-109
 - choice of requirement criterion and, 28
 - context for applications in this report, 1, 20
 - data limitations and, 27
 - dietary assessment compared, 1, 20-21, 22, 27, 62
 - distribution of usual intakes and, 26
 - with EAR, 4, 6, 8, 9-10, 29-30, 37, 38, 60, 72, 96-97, 124
 - energy metabolism considerations, 140-141
 - error sources in, 29-34; *see also* Uncertainties
 - genetic influences, 139-140
 - goals and goal setting, 3, 26, 30
 - hybrid approach, 58
 - implementation steps, 3, 5, 26-27
 - levels of, 1, 20
 - lifestyle factors affecting, 36, 144
 - nutrient density in, 90-91, 183-191
 - nutrient-nutrient interactions and, 135-136
 - nutritional status and, 28-29, 33, 133, 139, 145
 - research recommendations, 16-17, 57, 83, 88, 147-155
 - special considerations, 15-16, 36, 57, 133-146, 201-203
 - for subgroups, 13-14, 50, 53-54, 90, 97-102, 107, 108-152
- Dietary recalls, 31, 37, 63, 109
- Dietary records, 31, 37, 63, 109
- Dietary Reference Intakes (DRIs). *See also* Dietary planning
 - adjustment for specific individuals and populations, 134, 139-140, 144-146, 154
 - age and, 25, 139, 141, 143-144
 - applicable population, 2, 22, 144
 - assessment applications, 1, 20-21, 22, 27, 32-33, 37, 62, 64
 - categories, 21, 37; *see also* Adequate Intakes, Estimated Average Requirements; Recommended Dietary Allowances; Tolerable Upper Intake Levels
 - criteria for, 2, 23, 28
 - data limitations, 27, 133
 - defined, 1, 2, 20, 22
 - describing dietary survey data with, 17
 - energy intake and, 30, 41, 140
 - extrapolation of data, 168-169
 - framework, 20-21, 22, 23, 164, 165-166
 - group-level planning with, 4, 21, 55-132
 - incorporation into food guidance systems, 35, 36, 44-45, 47-49, 171
 - individual-level planning with, 4, 6, 21, 35-54
 - origin, 163-164
 - parameters for, 167-168; *see also* Life stage group; Reference heights and weights
 - physiological considerations, 140, 141-144
 - processes used to establish, 22-23
 - prorating across meals, 115-116
 - RDAs and RNIs contrasted, 2, 19, 22, 24, 47-48, 113, 164
 - uses, 2, 20-21, 37; *see also* Dietary assessment; Dietary planning
- Dietary reference standards, 44-45, 56
- Dietary survey data
 - adjusting intake distributions, 10
 - describing, 17
 - distribution of usual intakes from, 9, 10, 13, 33-34, 64, 70, 84, 109, 110-111, 123, 203-204, 205-206
 - fortification of foods and, 132
- Distribution of observed intakes
 - in group-level planning, 63-64, 78, 112, 196-208
 - statistical adjustment technique, 26, 196, 203-204

- systematic effects on, 203
 - usual intake distribution estimated
 - from, 26, 63-64, 196-208
 - variance of, 199
 - Distribution of requirements. *See also* Joint distribution; Nutrient requirements
 - AIs and, 103
 - and cut-point method, 17, 59
 - EAR and, 166-167
 - factorial modeling, 166-167
 - iron, 10, 24
 - Monte Carlo simulation, 167
 - normal/symmetrical, 23, 24, 27, 73, 184, 201-202
 - physiological considerations and, 140, 167
 - and probability approach, 17, 73-74
 - protein, 166, 184
 - skewed/asymmetrical, 10, 23, 24, 69-70, 73-74, 94, 80, 105, 166-168, 184
 - standard deviation of, 24, 166-167
 - for subgroups, 166, 184
 - and target usual nutrient intake distribution, 73-74, 80
 - variance of, 24, 59, 62, 74-75, 140
 - Distribution of usual intakes. *See also* Adjusting intake distributions; Skewed distribution; Target nutrient density distribution; Target usual nutrient intake distribution
 - baseline, 57, 59-60, 61, 62, 63-65, 99, 110-111, 112
 - characteristics of, 71
 - defined, 26
 - from distribution of observed intakes, 26, 63-64, 196-208
 - EAR cut-point method and, 57, 70, 72, 96-97
 - fortification of foods and, 127, 128-129
 - for group-level planning, 27, 28, 55-56, 57, 59-60, 61, 63-65, 74, 76, 83, 87-88, 97, 98, 110-111, 196-208
 - interventions to change, 16, 57, 60, 72, 87-88, 107, 108, 123-132, 148-149
 - for macronutrients, 205
 - median of, 65, 78, 110, 111, 124
 - as nutrient density, 98, 184-185, 190-191
 - from one day of intake data, 26, 47, 196, 200, 203, 205
 - and prevalence of inadequate intakes, 10, 28, 57, 58-59
 - probability approach and, 17, 74
 - skewed, 69-70, 166-167, 184, 188, 201-202
 - software for estimating, 82
 - stable, 71
 - in subpopulations, 90, 91-92, 94, 97, 107, 109, 110-111, 112, 115, 117, 184
 - supplement use and, 123-124, 125-126
 - from survey data, 9, 10, 13, 33-34, 64, 70, 84, 109, 110-111, 123, 203-204, 205-206
 - variance of, 9, 26, 33, 58, 59, 62, 63, 64, 67, 68, 71, 72, 74-75, 83, 92, 98-99, 109, 188, 197, 199-200, 207-208
 - Doubly labeled water method, 31, 41, 104
- ## E
- Education. *See* Nutrition education
 - EER (estimated energy requirement)
 - average, for group members, 76, 77-78
 - for body weight maintenance, 43
 - calculation for reference person, 12, 76-77
 - defined, 4, 6
 - derivation of, 23-24, 41-42, 105
 - for heterogeneous groups, 14, 76, 77
 - for homogeneous groups, 9
 - for individuals, 4, 41-43
 - physical activity coefficient, 42, 77
 - standard deviation, 41-42
 - uses, 6
 - Elderly people. *See* Adults over age 50
 - Emergency food rations, 81, 148, 150, 183-191
 - End-stage renal disease, 145-146
 - Energy expenditures
 - defined, 76
 - doubly labeled water method, 31, 41, 104
 - energy intakes and, 33, 42, 75-76, 105
 - error in reporting, 92-93, 104-106
 - factorial methods, 31, 33
 - total (TEE), 24, 41, 43, 76, 105
 - Energy intakes
 - and adequacy of nutrient intakes, 31-32, 91
 - adjustment of, 33, 205
 - athletes, 36, 141
 - averaging, 98-99, 100-101

- between-person variance, 98-99
 - bias related to, 33, 92, 105
 - comparison of target median nutrient intake to mean, 93-96, 118-119, 185
 - correlation between energy expenditures and, 33, 42, 75-76, 105
 - correlation between energy requirement, 41, 76, 78, 105, 140
 - correlation between nutrient intakes and, 98-102, 103-104
 - and DRIs, 30, 41, 140
 - and energy expenditures, 33, 42, 75-76, 77, 105
 - error sources, 12, 15, 30-34, 76, 92-93, 104-106
 - from fat, 9, 31, 32, 79-80, 205
 - group-level planning, 9, 12, 64-65, 72, 75-78, 89, 91-92, 93-96, 98-99, 118-119, 183, 185-186
 - individual-level planning, 4, 6, 31, 41-43, 49, 54
 - iron intakes and, 90
 - joint nutrient intake and, 31-32, 98-102, 103-104, 188
 - macronutrient distribution of, 6, 9, 30, 43, 78-80, 205
 - mean of distribution as target, 12, 13, 75-76, 84, 92, 93-96
 - and nutrient density distribution approach, 91-92, 97, 100-101, 103-104
 - prevalence of inadequate intakes, 78-80
 - self-reports, 12, 15, 30-34, 76, 93, 104-106, 151
 - simple nutrient density approach and, 89, 91, 93-96, 118-120
 - target usual nutrient intakes combined with, 96, 97-98, 105
 - underestimation, 31-32, 64-65
 - underreporting, 12, 15, 30-34, 76, 93, 104-106, 151
 - weight as measure of, 41
 - within-person variability, 31, 33, 41-42, 91-92
 - within-subgroup variability, 89, 90, 91-92, 95-96, 97, 98-99, 117
- Energy requirement. *See also* EER
- body size and, 140
 - defined, 76
 - derivation of, 24
 - energy intake correlation, 41, 76, 78, 84, 105, 140
 - mean, 89, 91, 93-96, 118-120
 - simple nutrient density approach and, 89, 91, 93-96, 118-120
- Error sources. *See also* Measurement error
- in dietary assessments, 64, 202
 - in dietary intake estimates, 29, 30-34, 37, 64, 70, 76, 196, 197
 - in energy intake data, 12, 15, 30-34, 76, 92-93, 104-106
 - in group-level planning, 30, 71
 - in individual-level planning, 37
 - in nutrient density approaches, 92-93, 104-106
 - in requirement estimates, 29-30, 105
 - research needs, 106
 - weight and height reports, 93, 106
- Estimated Average Requirements (EARs)
- Als compared, 4, 25, 154
 - criteria of adequacy, 23, 28, 29, 30
 - cut-point method, 9-10, 17, 55, 59, 60, 61, 62, 68-73, 74-76, 94, 96-97, 109, 110-111, 154
 - CV, 24, 65
 - defined, 3, 23-24
 - derivation of, 22-24, 28
 - dietary intake distribution adjustments, 65-68
 - endpoints, 28
 - and fortification of foods, 127
 - in group-level planning, 55-56, 59, 60, 65-68, 76
 - in individual-level planning, 8, 37, 38
 - by nutrient and life-stage group, 10
 - nutrient density approach and, 96-97
 - planning intakes with, 4, 6, 8, 9-10, 29-30, 37, 38, 60, 72, 96-97, 124
 - rationale for term, 23
 - and RDA, 23, 24, 28, 62, 133
 - research recommendations, 17, 153-154
 - risk of inadequacy, 22, 23
 - skewed distribution of requirements and, 166-167
 - standard deviation of intake for individual, 62, 65
 - uncertainty in, 27, 29-30, 133
 - uses, 2, 4, 127
 - variability related to, 133

Examples of planning for groups
 assisted-living facility for seniors, 108-113
 for heterogeneous groups, 116-123
 school nutrition program, 113-116
 Excessive intakes of nutrients, 6, 35, 40-41,
 50-51. *See also* Prevalence of excess
 intakes; Risk; Tolerable Upper
 Intake Levels
 Extrapolation of data, 27, 168-169

F

Factorial modeling, 31, 33, 166-167
 Fat, dietary
 Acceptable Macronutrient Range, 30,
 78, 80
 dietary guidelines, 54
 energy intakes from, 9, 31, 32, 79-80, 205
 nutrient content claims, 176
 nutrient interactions, 136
 underreporting of intakes, 30, 31
 Fiber, dietary, 85
 Fluoride, 52, 85, 192
 Folate, 28, 47, 49, 50, 52, 54, 86, 127, 134,
 135, 140, 142, 144, 152, 153, 193,
 194, 195
 Folic acid, 50
 Food and Agriculture Organization/
 World Health Organization, 28
 Food and Drug Administration, 51, 173, 174
 Food assistance programs, 2, 56-57, 58, 81,
 88, 113-116
 Food composition data, 7, 17, 27, 29, 30,
 44, 64, 132, 134, 151, 152, 194-195,
 206
 Food frequency questionnaires, 31, 37, 63
Food Guide for the Northwest Territories, 172
 Food Guide Pyramid, 7, 12, 35, 44, 45, 47,
 48, 49, 82, 171-173, 180
 Food guides, 7, 12, 17, 35, 45-49, 81, 82,
 152, 153, 171-173
 Food labels. *See* Nutrition labels and
 labeling
 Food offerings
 and group-level planning, 12, 72, 81,
 82-83, 112-113
 menu planning, 12, 72, 81, 82-83, 112-
 113
 and nutrient intakes, 16, 81, 82-83, 84,
 149
 research needs on, 149
 Food preferences, 13, 20, 84
 Food Stamp Program, 56-57, 58, 90, 113
 Food waste, 12, 82
 Fortification of foods, 2
 benefits of, 49-50
 bioavailability of nutrients, 134, 135
 in Canada, 50, 126, 192, 194
 detrimental effects, 126-127, 130-131
 in developing countries, 192
 and distribution of usual nutrient
 intakes, 127, 128-129
 EAR and, 127
 and food composition databases, 132,
 152, 194-195
 group-level planning, 87-88, 126-132
 impact on intakes, 129, 193-194
 individual-level planning, 7, 36, 40-41,
 49-50, 136
 interactions of nutrients and, 136
 mandatory programs, 50, 127, 132,
 148, 192, 194
 market pressures, 193
 modeling and estimating effects of,
 127-132, 193
 nutrition labeling and, 193, 195
 and prevalence of excess intakes, 130-
 131
 and prevalence of inadequate intakes,
 129-130, 132
 regulation, 50, 194
 supplement availability and use and,
 195
 and survey data, 132
 targeting, 126, 193
 technology for, 192-193
 and ULs, 25, 40, 127, 130-131, 136,
 152, 195
 in United States, 50, 126, 192-195
 Vitamin A added to milk, 127-132
 voluntary, 50, 126, 127, 132, 148, 192

G

Group-level planning. *See also* Examples of
 planning for groups;
 Heterogeneous groups;
 Homogeneous groups; Subgroups/
 subpopulations
 adequacy of nutrient intakes in, 29, 75,
 81, 83, 91, 95-96, 114, 115
 AIs in, 9, 11-12, 83-84, 85-87, 94-95, 103

analytic issues, 114-116
 assessment of plan results, 13, 16, 56,
 57, 78, 83, 84, 87, 95-96, 109, 113,
 120
 CV in, 75, 184, 185
 decision tree for, 106
 distribution of observed intakes in, 63-
 64, 78, 112, 196-208
 distribution of usual intakes for, 27,
 28, 55-56, 57, 59-60, 61, 63-65, 74,
 76, 83, 87-88, 97, 98, 110-111, 196-
 208
 EARs and, 55-56, 59, 60, 65-68, 76
 energy intakes, 9, 12, 64-65, 72, 75-78,
 89, 92, 93-96, 98-99, 118-119, 183,
 185-186
 energy requirements, 76, 77-78
 error sources, 30, 71
 food guides and, 17
 food offerings, 12, 72, 81, 82-83, 112-113
 fortified foods and, 87-88, 126-132
 general considerations, 56-58
 goals and goal setting, 3, 8-9, 11-12, 13,
 26, 28, 30, 55, 56-57, 61-62, 81-82,
 84, 95, 96, 99, 102-103; *see also*
 Nutrient density approaches;
 Target usual nutrient intake
 distribution
 hybrid approach, 58
 implementation steps, 7, 8, 16-17, 56,
 60
 interventions to change intake
 distribution, 57, 60, 72, 87-88, 123-
 132
 macronutrient distribution, 78-80
 menu planning, 11-13, 15, 61, 72, 80-
 87, 112-113, 116
 methods, 1, 55, 59-76
 overview, 58-63
 pilot testing approaches to, 16, 147-148
 prevalence of excessive intakes and, 8,
 55-56, 59, 87-88
 prevalence of inadequate intakes and,
 8, 27, 28, 57, 58-59, 60, 65-68, 74-
 75, 87-88, 94, 99, 110, 114-115, 117-
 118, 123, 185
 quantities of foods, 12
 RDAs and, 4, 21, 26-27, 55, 62, 65-68,
 113, 114, 116
 research implications and
 recommendations, 16-17, 57, 147-
 150

sample size considerations, 64, 71
 supplements and, 123-126
 ULs and, 8, 10-11, 55-56, 59, 60, 92,
 119, 120
 underlying principle, 58-59
 uses of DRIs for, 4, 21, 114
 Group mean intakes, 83, 84, 85-87, 200

H

*Handbook for Canada's Physical Activity
 Guide to Healthy Active Living*, 173
 Health Canada, 164
 Health claims, 174, 178-179
 Heterogeneous groups
 average nutrient requirement
 approach, 90
 comparison of target median nutrient
 intake to mean energy intake, 93-96
 EER, 14, 76, 77
 energy intake variability, 89, 91-92
 examples of planning for, 116-123
 goals and goal setting, 96, 99, 102-103
 menu planning for, 116
 nutrient density approaches, 13-15, 16-
 17, 96-103, 183-191
 target usual nutrient intake
 distribution for, 90, 94-95, 96-97
 Homogeneous groups
 dietary planning for, 8-13, 108-113
 EER for, 9
 energy intakes, 72, 76-78
 examples of planning for, 108-116
 food offerings, 12, 82
 goals and goal setting, 8-9, 11-12, 56,
 65-68, 81-82, 110-111; *see also* Target
 usual nutrient intake distribution
 interventions to change intake
 distribution, 57, 87-88, 107, 108,
 148-149
 macronutrient distribution, 78-80
 menu planning, 11-12, 72, 80-87, 112-
 113, 116
 quantities of foods to purchase, offer
 and serve, 12, 82-83
 reference person, 76-77
 target usual nutrient intake
 distribution for, 9-11, 26, 55, 59-76,
 110-111, 114-115
 Hospital patients, 58, 145
 Human milk, 25, 143

I

- Ill people, 36, 58, 139, 144-146
- Inadequacy of nutrient intake. *See*
 - Prevalence of inadequate intakes
- Indicators of nutrient adequacy, risk
 - reduction-based, 22, 23
- Individual-level planning
 - adequacy of nutrient intakes in, 26, 37, 45-47, 50, 53
 - AIs and, 4, 6, 35, 37, 39, 41
 - development of plans, 1, 4, 7, 35-36, 43-54
 - dietary guidelines and, 51, 53-54
 - dietary intake assessments and, 37
 - EAR and, 8, 37, 38
 - EER and, 4, 41-43
 - for energy intakes, 4, 6, 31, 41-43, 49, 54
 - error sources, 37
 - food guides and, 7, 35, 45-49, 172-173
 - food labels and, 51
 - fortified foods and, 7, 36, 40-41, 49-50, 136
 - goals and goal setting, 3, 4-6, 7, 35, 36-41
 - ill people, 36, 58, 139, 144-146
 - implementation steps, 4-7, 36-37
 - for macronutrient distribution, 43, 44, 51, 53
 - menu planning, 47, 48, 49
 - for normal healthy individuals, 36-54
 - nutrient-based food guidance systems
 - and, 7, 35, 43, 44-49, 51, 53-54, 172-173
 - RDAs and, 4, 6, 21, 26, 35, 37-39, 41, 134, 138-139
 - supplements and, 7, 35, 50-51
 - ULs and, 4, 6, 35, 37, 39-41
 - uses of DRIs for, 4, 6, 21, 35-54
- Infants
 - AI derivation for, 25
 - fortified foods, 50
- Institutional food planning, 2, 56, 58, 64, 72, 75, 81, 90, 108-113, 145
- Iodine, 50, 52, 87, 126, 184, 192
- Iodine deficiency diseases, 192
- Iowa State University method, 64, 109, 117, 196-197, 202, 203-204, 205, 206
- Iron, 10, 15, 24, 28, 38-39, 49, 50, 52, 73, 74, 87, 90, 94, 97, 102, 135-137, 139, 141, 142, 166, 167, 184, 188, 190, 192, 194, 195

J

- Joint distribution
 - of intakes and requirements, 75-76, 103, 104, 167, 188
 - of nutrient and energy intakes, 31-32, 98-102, 103-104, 188
- Joint Food and Agriculture Organization/World Health Organization Expert Consultation, 28

L

- Labels. *See* Nutrition labels and labeling
- Life stage group
 - AIs for nutrients by, 85-87
 - categories, 167-168
 - and derivation of DRIs, 167-168
 - food guidance by, 172
- Lifestyle factors, 36, 144
- Likelihood
 - of adequacy, 65, 91
 - of excessive intakes, 120
- Lipoprotein lipase deficiency, 139
- Long-term care facilities, 64
- Lowest-observed-adverse-effect level (LOAEL), 23
- Lysine, 137-138

M

- Macronutrient distribution
 - adjustment of intake data, 33, 205
 - DRVs, 174-175
 - of energy intake, 6, 9, 30, 43, 78-80, 205
 - for heterogeneous groups, 78-80
 - for homogeneous groups, 78-80
 - individual-level planning for, 43, 44, 51, 53
- Magnesium, 49, 52, 85, 135, 141
- Manganese, 87
- Mean intake, 12, 13, 75-76, 84, 89, 91, 92, 93-96, 97, 118-119, 200
- Mean requirement, 89, 90, 91, 93-96, 118-120
- Measurement error
 - in energy intake and energy expenditure, 92-93, 104-106
 - in estimated prevalence of inadequacy, 64-65, 76, 104-106

- in nutrient intakes, 15-16, 29, 30-34, 37, 70, 196, 197
 - in physiological parameters, 92-93, 106
 - in self-reported data, 106
 - Measuring dietary intakes
 - observational, 15-16, 109, 197
 - self-reports, 15-16, 27, 29, 30-34, 63, 64-65, 76, 93, 104-106, 151
 - Men
 - EER, 76-77
 - serving size, 195
 - vitamin C, 117, 119, 120-121, 122
 - Menu planning
 - with AIs, 11-12, 83-84, 85-87
 - assessment of results, 84, 87
 - contexts, 81
 - food offerings, 12, 72, 81, 82-83, 112-113
 - goals, 11-12, 81-82
 - for groups, 11-13, 15, 61, 72, 80-87, 112-113, 116
 - for individuals, 47, 48, 49
 - quantities of foods, 12, 82-83
 - research needs, 83, 153
 - steps, 81-87
 - Methionine, 137, 138
 - Molybdenum, 87, 184
 - Monte Carlo simulation, 167
- N**
- National Health and Nutrition Examination Survey
 - Fourth, 151
 - Third, 10, 28-29, 30, 60, 70, 74, 109, 110-111, 114, 123, 151, 168, 169, 170
 - National Nutrition Monitoring and Related Research Act, 180
 - National Research Council, 64, 109. *See also* NRC method of adjustment
 - National School Lunch Program, 113, 149, 153
 - Neural tube defects, 135, 141-142
 - Niacin, 49, 50, 52, 86, 135, 140, 144, 184, 192
 - Night blindness, 28, 128, 129
 - No-observed-adverse-effect level (NOAEL), 23
 - Nonnormal distribution. *See* Skewed distribution
 - Normal distribution
 - of intakes, 57, 59-60, 61, 62, 65-66, 68-71, 80, 99, 110-111, 112, 198-199
 - of requirements, 23, 24, 27, 73, 184, 201-202
 - and target usual nutrient intake distribution, 57, 59-60, 61, 62, 63-65, 110-111, 112
 - Normative storage requirement level, 28
 - NRC method of adjustment, 64, 109, 196, 197-203, 205, 206, 207-208
 - Nutrient-based food guidance systems. *See also* Food guides
 - in group-level planning, 17
 - incorporation of DRIs in, 35, 36, 44-45, 47-49, 171
 - and individual-level planning, 7, 35, 43, 44-49, 51, 53-54, 172-173
 - research recommendations, 17, 152-153
 - technical tools for professionals, 153
 - United States, 7, 12, 35, 44, 45, 47, 48, 49, 51, 53, 82, 171-173, 180-181
 - Nutrient calculation software, 12, 82
 - Nutrient content claims, 174, 176-177, 178-179
 - Nutrient content of food, 11
 - Nutrient density
 - average intake, 98-99, 121
 - calculation of usual intake distribution as, 98, 184-185, 190-191
 - defined, 13, 91, 98
 - reference (median intake) distribution, 96, 102-103, 119, 187-188
 - use in dietary planning, 90-91, 183-191
 - Nutrient density approaches. *See also* Nutrient density distribution approach; Simple nutrient density approach
 - comparison of, 122-123
 - for heterogeneous groups, 13-15, 16-17, 96-103, 183-191
 - mathematical proof, 188-191
 - and prevalence of inadequate intakes, 94, 96-97, 99, 102, 185-187
 - research recommendations, 16-17, 149-150
 - Nutrient density distribution approach
 - AIs and, 103
 - correlation between nutrient intakes and energy intakes, 98-102, 103-104, 122

derivation of target usual density
 intake distribution, 14-15, 96, 97-102, 121-122, 184-185
 direct interventions used with, 102-103
 effectiveness, 103
 energy intakes and, 91-92, 97, 100-101, 103-104
 error sources, 104-106
 example, 120-123
 identifying reference distribution and setting goals, 102-103, 119-120
 strengths of, 102-103
 target usual nutrient intake
 distribution and, 14-15, 16-17, 92, 96-97, 99-100, 103, 104-105, 117-118, 120-121
 technical considerations, 103-106
 ULs and, 96, 102, 119, 187
 underreporting of energy intakes and, 104-106
 Nutrient intakes. *See* Observed intakes; Usual nutrient intakes
 Nutrient-nutrient interactions, 135-136
 Nutrient requirements, *See also*
 Distribution of requirements;
 Energy requirement; Estimated Average Requirements
 average/mean, 90, 166-167
 correlated with usual intakes, 17, 41, 61, 75-76, 78
 criterion of nutritional adequacy, 28
 CV, 166, 184
 error sources in estimates, 29-30, 105
 physiological considerations, 140-144
 research recommended, 17, 153-156
 special considerations, 139-144
 uncertainties for individuals, 27, 29-30, 38
 variance of, 59, 139-144, 166
 Nutrition Canada Survey, 168-169
 Nutrition education, 15, 35, 44, 87-88, 95, 125, 148, 152-153
 Nutrition Labeling and Education Act, 174
 Nutrition labels and labeling, 2, 7
 Canadian, 51, 54, 177-179
 components, 176, 179
 Daily Values—Reference Daily Intakes, 51, 52, 174, 175-176, 179
 development of, 173-175, 177-178
 fortification of foods and, 193, 195
 and individual-level planning, 51

 nutrient content claims, 174, 176-177, 178-179
 RDAs and AIs compared to DVs on, 52
 research recommendations, 153
 supplements, 17
 United States, 51, 53, 173-175, 176-177
 usefulness in planning, 35
 Nutritional status, and dietary planning, 28-29, 30, 33, 133, 139, 145
 Nutrition surveys. *See* Dietary survey data; *individual surveys*

O

Observed intakes. *See also* Distribution of observed intakes
 and individual-level planning, 37
 mean intakes, 39, 83
 probability of inadequacy or excess, 37
 short-term, 37
 Oral contraceptives, 142

P

Pantothenic acid, 25, 52, 86
 Parenteral nutrition, 145
 Pellagra, 192
 Phenylketonuria, 139
 Phosphorus, 49, 52, 85, 141, 146
 Physical activity, 24, 31, 32, 42, 77, 93, 105, 106, 140-141
 Physiological consideration, 27, 92-93, 106, 140, 141-144, 167
 Phytic acid, 136
 Phytochemicals, 54
 Plasma pyridoxal phosphate levels, 29, 110
 Polyunsaturated fatty acids, 43, 78, 85
 Pregnant and lactating women
 EER, 4
 gestation of multiple fetuses, 142-143
 menu planning, 47, 48, 49
 supplement use, 50, 141-142
 Prevalence of excess intakes
 estimation, 9, 10-11
 fortification of foods and, 130-131
 group-level planning and, 8, 55-56, 59, 87-88
 skewed data and, 201-202
 zinc, 10-11

Prevalence of inadequate intakes

- AIs and, 84, 94
 - bias in, 64-65, 76, 104-106
 - distribution of usual intakes and, 10, 28, 57, 58-59, 87-88, 196
 - EAR cut-point method, 9-10, 17, 55, 59, 60, 61, 62, 68-73, 75-76, 94, 96-97, 109, 110-111, 123, 154
 - energy, 78-80
 - estimation, 9-10, 11-12, 26-27, 55, 59, 71
 - fortification of foods and, 129-130, 132
 - in group-level planning, 8, 27, 28, 57, 58-59, 60, 61, 65-68, 74-75, 87-88, 94, 99, 110, 114-115, 117-118, 123, 185
 - iron, 10
 - low individual vs. low group risk, 65-68
 - median intake equal to RDA, 65-68
 - nutrient density approach and, 94, 96-97, 99, 102, 185-187
 - and nutritional status, 28-29, 30
 - overestimation, 64-65
 - probability approach to estimating, 17, 59, 65, 73-76, 94, 97, 154
 - research recommendations, 17, 154
 - skewed data and, 87-88, 201-202
 - standard deviation of estimates, 166-167
 - statistical approaches, 64, 109, 117, 167, 196-204, 205, 206, 207-208
 - in subpopulations, 117, 123, 187
 - and target usual nutrient intake distribution, 57, 59, 60, 61, 65-68, 71, 73, 110, 114-115
 - uncertainty in, 71
 - underestimation of intakes and, 64-65
 - zinc, 10
- Probability approach
- cut-point method compared, 17, 59, 73-76
 - and distribution of nutrient requirements, 17, 73-74
 - and distribution of usual intakes, 17, 74
 - prevalence of inadequacy from, 17, 59, 65, 73-76, 94, 97, 154
- Professional development and education, 17, 152-153
- Protein, dietary
- Acceptable Macronutrient Range, 78
 - food sources, 47
 - intake, 31-32, 43, 49. *See also* Macronutrient distribution

- percent of total energy intake, 78-80
- physical activity and, 140
- requirement distribution, 166, 184
- vegan diets, 137-138

Provitamin A carotenoids, 47, 48, 152

R

Recommended Dietary Allowances (RDAs)

- AIs compared, 4, 25
 - applicable population, 2, 22
 - context for use, 37-39
 - CV, 24
 - Daily Values on food labels compared, 52
 - defined, 3, 24
 - derivation, 23, 24, 28
 - and EAR, 23, 24, 28, 62, 133
 - and food and nutrition assistance programs, 113, 114, 116
 - and food guides, 44-45, 47-49, 171, 172
 - and food labeling and nutritional marketing, 52
 - group-level planning with, 4, 21, 26-27, 55, 62, 65-68, 113, 114, 116
 - inappropriate use of, 21, 22, 41
 - individual-level planning with, 4, 6, 21, 26, 35, 37-39, 41, 134, 138-139
 - and median target usual nutrient intake distribution, 62, 65-66
 - replacement with DRIs, 2, 19, 22, 24, 47-48, 113, 164
 - research recommendations, 153-154
 - risk of inadequacy, 38
 - target usual nutrient intake distribution and, 62, 65-68
 - uses, 4, 7, 20, 24, 44-45, 164
- Reference Amounts Customarily Consumed Per Eating Occasion, 176
- Reference Daily Intakes (RDIs), 51, 174, 175-176
- Reference heights and weights and, 168-170
- Reference (median intake) distribution, 96, 102-103, 119-120, 187-188
- Reference person
- EER calculation for, 12, 76-77
 - in homogeneous groups, 76-77
 - in subgroups, 95, 102, 109, 119-120, 121-122, 187

Research recommendations
 group-level planning, 16-17, 57, 147-150
 improving requirement data, 17, 153-156
 nutrient-based food guidance systems, 17, 152-153
 quality of dietary intake data, 17, 149, 150-152, 153
 Residual method of energy adjustment, 33
 Retinol activity equivalents, 47-48, 115
 Retinol equivalents, 48
 Riboflavin, 49, 50, 52, 86, 140, 142
 Richard B. Russell National School Lunch Act, 113
 Rickets, 139, 194
 Risk
 of excess, 6, 22, 39-40, 50-51, 135, 136
 of inadequacy, 6, 22, 23, 30, 38, 65-68, 102, 145
 Risk curve, 155
 Risk-reduction indicator of nutrient adequacy, 22, 23

S

Salt, 54
 Sample size considerations, 64, 71, 197, 202-203, 204
 School Breakfast Program, 113
 School nutrition program, 90, 113-116
 Selenium, 52, 85, 135
 Self-reported data
 accuracy of, 15-16, 27, 29, 30-34, 63, 64-65, 104, 151
 alternatives to, 105-106
 energy intakes, 12, 15, 30-34, 76, 93, 104-106, 151
 measurement error in, 106
 nutrient intakes, 15-16, 29, 30-34, 64
 weight and height, 93, 106
 Senate Select Committee on Nutrition and Human Needs, 171, 180
 Sensory neuropathy, 40
 Serum alpha-tocopherol, 30
 Serum ferritin, 139
 Serving sizes, 45, 47, 116, 176, 195
 Shapiro-Wilk statistic, 198
 SIDE package, 293
 Simple nutrient density approach
 accuracy, 93-94
 AIs and, 94, 103
 assessment of results, 92, 120
 comparison of target median nutrient intake and energy intake, 94, 95, 119-120, 184-185
 derivation of nutrient density distribution, 184-185, 190-191 and EARS, 96-97
 error sources, 92-93, 104
 example, 117-120, 122-123
 and mean energy intake or requirement, 89, 91, 93-96, 118-120
 median of target nutrient intake distribution and, 14, 89, 91, 93-96, 103, 117-118, 119-120
 multiple-food basis, 91
 single-food basis, 91, 183-191
 and ULs, 92, 95, 96, 187
 Skewed distribution
 adjusting, 166-167, 184, 188, 198-199, 201-202
 of intakes, 69-70, 84, 166-167, 184, 188, 201-202
 of iron, 10, 73, 94, 166, 184, 188
 of nutrient requirements, 10, 23, 24, 69-70, 73-74, 94, 80, 105, 166-168, 184
 and prevalence of inadequacy or excess, 87-88, 201-202
 of protein, 184
 and target usual nutrient intake distribution, 69-70, 80, 84, 94
 transformation of, 166, 198-199, 200-201, 202, 203
 Sodium, 146
 Special considerations
 adjusting intake distributions for, 57, 201-203
 illness, 144-146
 individual characteristics that influence requirements, 139-144
 lifestyle factors, 36, 144
 source of nutrient, 134-139
 in supplement use, 136, 142, 143, 144, 173
 Standard deviation
 of distribution of requirements, 24, 166-167
 EAR and, 24, 62, 85, 166
 EER, 41-42
 of intakes, 24, 62, 65, 67-68, 69, 71, 72, 166

- in prevalence estimates, 166-167
 - Z values, 69
 - Statistical Analysis System (SAS)
 - programs, 185, 186, 190-191, 198, 207-208
 - Statistical approaches, 64, 109, 117, 167, 196-204, 205, 206, 207-208
 - Subgroups/subpopulations. *See also*
 - Heterogeneous groups;
 - Homogeneous groups; Special considerations
 - adjustment of DRIs for, 134, 139-140, 144-146, 154
 - direct interventions, 15, 50, 102, 126-132
 - distribution of usual intakes in, 90, 91-92, 94, 97, 107, 109, 110-111, 112, 115, 117, 184
 - energy intake variability within, 89, 90, 91-92, 95-96, 97, 98-99, 117
 - planning intakes for, 13-14, 50, 53-54, 90, 97-102, 107, 108-152
 - prevalence of inadequacy in, 117, 123, 187
 - reference intakes, 95, 102, 109, 119-120, 121-122, 187
 - skewed requirements, 166, 184
 - statistical adjustment of intake data, 204
 - and target usual nutrient intake
 - distribution, 13-14
 - variability within, 89, 95-96, 97, 98-99
 - vegetarians, 36, 136-139
 - vulnerable, 13-14, 15, 50, 89, 91, 95
 - Summer Food Service Program, 113
 - Supplement use
 - bioavailability of nutrients, 134, 135
 - disadvantages, 50
 - and distribution of usual intakes, 123-124, 125-126
 - and fortification of foods, 195
 - in group-level planning, 123-126
 - high-dose/high-potency, 40-41
 - individual-level planning, 7, 35, 50-51
 - interactions, 136
 - labeling, 17
 - pregnancy and lactation and, 142
 - quantifying intakes from, 17, 151
 - research recommendations, 17, 151
 - special considerations in, 136, 142, 143, 144, 173
 - targeted, 15, 36, 95, 143
 - and ULs, 120
 - Supplemental Nutrition Program for Women, Infants, and Children, 113, 149
 - Surveys. *See* Dietary survey data; *individual surveys*
- ## T
- Target nutrient density distribution. *See also* Nutrient density distribution
 - approach
 - derivation, 14-15, 96-102, 103, 121-122, 184-185
 - percentile method, 185, 187-188
 - and ULs, 102
 - Target usual nutrient intake distribution
 - asymmetric requirement distribution and, 73-74, 80
 - baseline usual nutrient intake
 - distribution and, 57, 59-60, 61, 62, 63-65, 110-111, 112
 - bias in, 74
 - comparison of median nutrient intake to mean energy intake, 91, 93-96, 97
 - concept, 59-60
 - considerations in planning, 63-76
 - energy intakes and, 12, 13, 78-80, 84, 93-96, 118-119, 185
 - feasibility of obtaining, 71-73
 - for heterogeneous groups, 90, 94-95, 96-97
 - for homogeneous groups, 9-11, 26, 55, 59-76, 110-111, 114-115
 - median of, 10-11, 13, 14-15, 61-63, 65-68, 69, 79-80, 81-83, 89, 91, 93-96, 109, 112-113, 114-115, 117-118, 119-120
 - menu planning to achieve, 11-13, 61, 80-87, 112-113
 - normal distribution of usual intake and, 68-69
 - nutrient density approaches and, 14-15, 16-17, 91, 92, 93-97, 99-100, 103, 104-105, 117-118, 120-121
 - prorating across meals, 115-116
 - RDA and, 62, 65-68
 - research recommendations, 149
 - skewed distribution of usual intake and, 69-70, 80, 84, 94

stable distribution of usual intakes
and, 71
target prevalence of inadequacy and,
57, 59, 60, 61, 65-68, 71, 73, 110,
114-115
usual energy intakes combined with,
96, 97-98, 105
vulnerable subgroup and, 13-14
Z value, 69
Thiamin, 49, 50, 52, 86, 140, 144, 194
Threonine, 137, 138
Tolerable Upper Intake Levels (ULs)
bioavailability considerations, 135, 136
context for use, 39-41
critical adverse effect, 22
defined, 3, 25, 39
derivation of, 25-26
dose-response assessment, 155
and food labels, 153
fortification of foods and, 25, 40, 127,
130-131, 136, 152, 195
in group-level planning, 8, 10-11, 55-
56, 59, 60, 92, 119, 120
in individual-level planning with, 4, 6,
35, 37, 39-41
LOAEL/NOAEL, 23
nutrient density approach and, 92, 95,
96, 102, 119, 120, 187
rationale for term, 25
research recommendations, 153, 154-
155
risk assessment approach, 25
supplement use and, 25, 40
uncertainty levels, 23, 134
uses, 2, 4
Total energy expenditure (TEE), 24, 41,
43, 76, 105
Tryptophan, 137, 138

U

Uncertainties
in adequacy of nutrient intakes, 29
in AIs, 4, 39, 133-134
in dietary intake estimates, 27, 30-34,
37
in EAR, 27, 29-30, 133
in nutrient requirements for
individuals, 27, 29-30, 38
in prevalence of inadequate intakes, 71
in ULs, 23, 134

United States
Department of Agriculture, 113-114,
134, 171, 180, 203
Department of Health and Human
Services, 180
Dietary Goals (1977), 171
dietary guidelines, 35, 51, 53, 180-181
Food Guide Pyramid, 7, 12, 35, 44, 45,
47, 48, 49, 82, 171-173, 180
food labels, 51, 53, 173-175, 176-177
fortification of foods, 50, 126, 192-195
Urinary nitrogen excretion, 31-32
US RDA, 51, 173-174, 175
Usual nutrient intakes. *See also* Dietary
intake data; Target usual nutrient
intake distribution
assessment, 63
availability of data, 108-109, 149
baseline distribution, 57, 59-60, 61, 62,
63-65
between-person variance, 63, 64, 75,
188
correlation between usual energy
intakes and, 98-102, 103-104
correlation of requirements and, 17,
41, 61, 75-76, 78
data sources, 9, 108-109, 114
defined, 6, 26
errors in, 15-16, 29, 30-34, 37, 70, 196,
197
food choices and, 16, 81, 82-83, 84,
149
joint energy intake and, 104
measurement of, 15-16, 27, 29, 30-34,
63, 64-65, 76, 93, 104-106, 109, 151,
197
nutrient content of food and, 11
observed vs. true, 63
self-reports, 15-16, 29, 30-34, 64
short-term intakes and, 26, 29, 47, 196,
200, 203, 205
within-person (day-to-day) variance, 9,
26, 63, 64, 92, 109, 197, 198, 199

V

Variance. *See also* Between-person
variance; Within-person variance in
intakes
EAR-related, 133

in intake distributions, 9, 26, 33, 58, 59, 62, 63, 64, 67, 68, 71, 72, 74-75, 83, 92, 98-99, 109, 188, 197, 199-200, 201, 207-208

in nutrient requirements, 59, 139-144, 166

in requirement distributions, 24, 59, 62, 74-75, 140

true, 199

unexplained, 199

Vegetarian diets, 36, 136-139

Vitamin A, 2, 28, 47-48, 49, 52, 86, 114-115, 127-132, 136, 152, 184, 192, 193, 194, 202

Vitamin B₆, 28-29, 40, 49, 50, 52, 86, 110-113, 142, 144

Vitamin B₁₂, 28, 49, 52, 86, 135, 136, 143-144, 153

Vitamin C, 14, 23, 47, 49, 52, 54, 86, 98, 101, 114-115, 117-122, 135, 136, 140, 144, 185, 186-187, 194

Vitamin D, 39, 50, 52, 86, 135, 136, 139, 194

Vitamin E, 30, 49, 52, 54, 86, 135, 136, 140, 152

Vitamin K, 86, 136, 176

Vulnerable subgroups, 13-14, 15, 50, 89, 91, 95

W

Weight. *See also* Reference heights and weights

accuracy of self-reports, 93, 106

changes, 42, 168

EER and, 43

errors in reporting, 93, 106

maintaining current, 31, 41, 42, 43, 75, 76

as measure of energy intake, 41

monitoring, 43, 78, 143

nutrient density approach and, 93, 106

Wernicke-Korsakoff syndrome, 144

White House Conference on Food, Nutrition, and Health, 173

Within-person variance in intakes

adjusting intake distributions for, 33, 63-64, 199-200, 204, 207-208

energy intakes, 31, 33, 41-42, 92

incomplete data and, 202-203

systematic effects on, 33, 203, 204

in usual intakes, 9, 26, 63, 64, 92, 109, 197, 198, 199

Women. *See also* Pregnant and lactating women

EER, 42, 43

energy intakes, 99, 101, 185

folate, 28, 135, 141-142

fortified food intakes, 135, 194

iron, 38-39, 74, 90, 190

macronutrient distribution for, 44, 78-80

menopause/hormone therapy, 142

serving size, 195

Vitamin A, 127-130, 131

Vitamin C, 101, 117-118, 119, 120, 122, 123, 185, 186-187

World Health Organization, 28

Z

Zinc, 10-11, 40, 49, 52, 60, 62, 70, 87, 95, 114-115, 123-126, 136-137, 198-201